


Mild Conditions for the Synthesis of  
Functionalized Pyrrolidines via  
Pd-Catalyzed Carboamination Reactions

Myra Beaudoin Bertrand, Matthew L. Leathen, and John P. Wolfe\*

Department of Chemistry, University of Michigan, 930 N. University Avenue,  
Ann Arbor, Michigan, 48109-1055  
jpwolfe@umich.edu

Received November 18, 2006


## ABSTRACT



The palladium-catalyzed carboamination of *N*-protected  $\gamma$ -aminoalkenes with aryl bromides and triflates has been achieved under new, mild reaction conditions using the weak base  $\text{Cs}_2\text{CO}_3$  in dioxane solvent. These reactions tolerate a wide variety of functional groups, including enolizable ketones, nitro groups, methyl esters, and acetates, which are not compatible with previously described conditions.

The development of synthetic methods for the construction of substituted pyrrolidines has been of longstanding importance in organic chemistry as a result of the prevalence of this moiety in biologically active molecules and natural products.<sup>1</sup> Over the past several years, the palladium-catalyzed carboamination of  $\gamma$ -aminoalkenes with aryl bromides has emerged as an efficient and stereoselective method for the construction of substituted pyrrolidine derivatives.<sup>2,3</sup> These transformations effect tandem cyclization and coupling in a process that generates a C–N bond, a C–C bond, and up to two stereocenters in one step. For example, treatment of Boc-protected amine **1** with 4-bromoanisole in

the presence of  $\text{NaOtBu}$  and catalytic amounts of  $\text{Pd}_2(\text{dba})_3$  and dppb afforded pyrrolidine **2** in 60% yield with >20:1 dr (eq 1).<sup>2c</sup>



Despite the synthetic utility of these transformations, the reactions are typically conducted in the presence of the strong

Table 1. Optimization Summary<sup>a</sup>

entry	base	Pd	solvent	yield (%)
1	$\text{NaOtBu}$	$\text{Pd}_2(\text{dba})_3$	toluene	81
2	$\text{Cs}_2\text{CO}_3$	$\text{Pd}_2(\text{dba})_3$	toluene	38
3	$\text{Cs}_2\text{CO}_3$	$\text{Pd}(\text{OAc})_2$	toluene	63
4	$\text{Cs}_2\text{CO}_3$	$\text{Pd}(\text{OAc})_2$	dioxane	82 <sup>b</sup>

<sup>a</sup> Conditions: 1.0 equiv substrate, 1.2 equiv ArBr, 2.3 equiv base, 1 mol %  $\text{Pd}_2(\text{dba})_3$  (2 mol % Pd) or 2 mol %  $\text{Pd}(\text{OAc})_2$ , 2 mol % Dpe-phos (with  $\text{Pd}_2(\text{dba})_3$ ) or 4 mol % Dpe-Phos (with  $\text{Pd}(\text{OAc})_2$ ), solvent (0.25 M), 105 °C. <sup>b</sup> The reaction was conducted at 100 °C.

(1) For recent reviews, see: (a) Bellina, F.; Rossi, R. *Tetrahedron* **2006**, 62, 7213–7256. (b) Coldham, I.; Hufton, R. *Chem. Rev.* **2005**, 105, 2765–2810.

(2) (a) Ney, J. E.; Wolfe, J. P. *Angew. Chem., Int. Ed.* **2004**, 43, 3605–3608. (b) Lira, R.; Wolfe, J. P. *J. Am. Chem. Soc.* **2004**, 126, 13906–13907. (c) Bertrand, M. B.; Wolfe, J. P. *Tetrahedron* **2005**, 61, 6447–6459. (d) Ney, J. E.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, 127, 8644–8651. (e) Yang, Q.; Ney, J. E.; Wolfe, J. P. *Org. Lett.* **2005**, 7, 2575–2578. (f) Ney, J. E.; Hay, M. B.; Yang, Q.; Wolfe, J. P. *Adv. Synth. Catal.* **2005**, 347, 1614–1620. (g) Bertrand, M. B.; Wolfe, J. P. *Org. Lett.* **2006**, 8, 2353–2356.

(3) For a review on other Pd-catalyzed alkene carboamination reactions that afford pyrrolidine products, see: (a) Wolfe, J. P. *Eur. J. Org. Chem.* **2007**, in press. See also: (b) Harayama, H.; Abe, A.; Sakado, T.; Kimura, M.; Fugami, K.; Tanaka, S.; Tamaru, Y. *J. Org. Chem.* **1997**, 62, 2113–2122. (c) Scarborough, C. C.; Stahl, S. S. *Org. Lett.* **2006**, 8, 3251–3254. (d) Sherman, E. S.; Chemler, S. R.; Tan, T. B.; Gerlits, O. *Org. Lett.* **2004**, 6, 1573–1575. (e) Larock, R. C.; Yang, H.; Weinreb, S. M.; Herr, R. J. *J. Org. Chem.* **1994**, 59, 4172–4178.

**Table 2.** Palladium-Catalyzed Carboamination of *N*-Protected  $\gamma$ -Aminoalkenes with Functionalized Aryl Bromides<sup>a</sup>

entry	amine	aryl bromide	product	yield (%) <sup>b</sup>	entry	amine	aryl bromide	product	dr	yield (%) <sup>b</sup>
1				75	8				—	88 <sup>e</sup>
2				82	9				12:1	80
3				78 <sup>e</sup>	10				15:1	76
4				76 <sup>e</sup>	11				>20:1	75
5				79	12				>20:1	74
6				76 <sup>c,e</sup>	13				>20:1	71
7				88	14				>20:1	73
				17 <sup>d</sup>						

<sup>a</sup> Conditions: 1.0 equiv amine, 1.2 equiv ArBr, 2.3 equiv Cs<sub>2</sub>CO<sub>3</sub>, 2 mol % Pd(OAc)<sub>2</sub>, 4 mol % Dpe-phos, dioxane (0.2–0.25 M), 100 °C. <sup>b</sup> Yield refers to average isolated yield obtained in two or more experiments. <sup>c</sup> Dppe used in place of Dpe-phos. <sup>d</sup> NaOtBu used in place of Cs<sub>2</sub>CO<sub>3</sub>. <sup>e</sup> The reaction was conducted at 85 °C in DME solvent.

base NaOtBu, which limits the scope of this method. For example, the use of NaOtBu restricts the functional group tolerance of these reactions, and transformations of aryl triflate electrophiles, which decompose in the presence of strong base, have not been reported. Additionally, Cbz protecting groups, which are frequently employed in the synthesis of complex alkaloids, are incompatible with the strongly basic conditions. In this Letter we describe the development of new conditions that replace NaOtBu with weaker bases (Cs<sub>2</sub>CO<sub>3</sub> or K<sub>3</sub>PO<sub>4</sub>), which significantly expands the scope of the carboamination method.

In our preliminary studies on palladium-catalyzed carboamination reactions of  $\gamma$ -(*N*-Boc-amino)- or  $\gamma$ -(*N*-acyl-amino)alkenes, our attempts to conduct the transformations using bases other than NaOtBu were met with limited success.<sup>2c</sup> For example, the Pd<sub>2</sub>(dba)<sub>3</sub>/Dpe-phos<sup>4</sup> catalyzed carboamination of **3** with 4-bromo-*tert*-butylbenzene afforded

**4** in 81% yield when the reaction was conducted in toluene solvent with NaOtBu as base (Table 1, entry 1). However, use of Cs<sub>2</sub>CO<sub>3</sub> in place of NaOtBu provided only a 38% isolated yield of **4** and led to the formation of large amounts of side products (entry 2).<sup>5,6</sup>

To improve the yields obtained in Pd-catalyzed carboamination reactions that employ mild bases, the effect of palladium source and solvent were systematically examined; the key results of these studies are summarized in Table 1. After some experimentation, it was discovered that use of Pd(OAc)<sub>2</sub> in place of Pd<sub>2</sub>(dba)<sub>3</sub> leads to significantly improved yields of **4** (63%, entry 3), and replacement of

(5) The use of Cs<sub>2</sub>CO<sub>3</sub> and other weak bases in Pd-catalyzed *N*-arylation reactions of amines with aryl halides has been reported. For reviews, see: (a) Muci, A. R.; Buchwald, S. L. *Top. Curr. Chem.* **2002**, *219*, 131–209. (b) Hartwig, J. F. In *Modern Arene Chemistry*; Astruc, D., Ed., Wiley-VCH: Weinheim, 2002; pp 107–168. (c) Schlummer, B.; Scholz, U. *Adv. Synth. Catal.* **2004**, *346*, 1599–1626.

(6) Lower yields were obtained with other weak bases including K<sub>2</sub>CO<sub>3</sub> and Et<sub>3</sub>N.

(4) Dpe-phos = bis(2-diphenylphosphino)phenyl ether.

toluene with dioxane as solvent provides optimal results (82%, entry 4).<sup>7,8</sup>


As shown in Table 2, the new reaction conditions described above are effective for the transformation of a number of different substrate combinations. A variety of functional groups are tolerated under these mild conditions, including aldehydes (entry 3), enolizable ketones (entry 4), nitro groups (entries 6 and 11), methyl esters (entries 8 and 14), and alkyl acetates (entry 9). In addition, the carboamination reactions of electron-rich (entry 10), electron-neutral (entries 1, 2, 5, 7, and 13), and heterocyclic (entry 12) aryl bromides proceed with good chemical yields. The mild conditions also are effective for stereoselective reactions and provide selectivities that are comparable to those observed in reactions that use NaOtBu as base. For example, transformations of starting materials **1** and **9**, which bear a substituent adjacent to the nitrogen atom, provide *cis*-2,5-disubstituted products **20** and **21** with excellent (>20:1) diastereoselectivity (entries 11 and 12). Similarly, substrates **7** and **8**, which are substituted at the allylic position, are transformed to *trans*-2,3-disubstituted products **18** and **19** with good stereocontrol (12–15:1).

In addition to providing increased tolerance of base-sensitive functional groups, the new reaction conditions also allow the efficient carboamination of substrates bearing Cbz protecting groups. For example, the Pd-catalyzed coupling of **6** with 2-bromonaphthalene using Cs<sub>2</sub>CO<sub>3</sub> as base provided the desired product **16** in 88% isolated yield (entry 7). In contrast, cleavage of the Cbz group from the substrate was problematic when reactions were conducted with NaOtBu as base; these conditions provided only a 17% yield of **16**.

More complex  $\gamma$ -aminoalkene substrates are also efficiently transformed using the new reaction conditions. As shown in Table 2 (entries 13 and 14), Pd-catalyzed reactions of **10** with bromobenzene or methyl-4-bromobenzoate proceeded smoothly to provide **22** and **23** with excellent stereoselectivity. Trisubstituted pyrrolidine **22** has been previously employed as an intermediate in the synthesis of the natural product (+)-preussin.<sup>2g,9</sup>

The high degree of functional group tolerance of this method also allows straightforward access to 1-substituted tetrahydropyrroloisquinolin-5-ones. As shown in Scheme 1, the Pd-catalyzed reaction of **8** with methyl-2-bromobenzoate afforded pyrrolidine **24** in 73% yield with 14:1 dr. Treatment of this product with trifluoroacetic acid followed by an alkaline workup gave **25** in 95% yield.

**Scheme 1.** Stereoselective Synthesis of **25**



The main limitations of these new reaction conditions involve transformations of sterically encumbered substrate combinations.<sup>10</sup> For example, attempts to convert substrates bearing internal alkenes to pyrrolidines were unsuccessful under these conditions. In addition, the reaction of methyl 2-bromobenzoate with **1**, which bears a substituent on C-1 (adjacent to the nitrogen atom), was not effective. However, as noted above, this *ortho*-substituted aryl bromide was effectively coupled with the less hindered carbamate **8** (Scheme 1).

**Table 3.** Palladium-Catalyzed Carboamination of Aryl Triflates<sup>a</sup>

entry	amine	aryl triflate	product	dr	yield (%) <sup>b</sup>
1				—	67 0 <sup>c</sup>
2				—	74
3				12:1	70
4				>20:1	64

<sup>a</sup> Conditions: 1.0 equiv amine, 1.2 equiv ArOTf, 2.3 equiv K<sub>3</sub>PO<sub>4</sub>, 4 mol % Pd(OAc)<sub>2</sub>, 8 mol % Dpe-phos, dioxane (0.25 M), 100 °C. <sup>b</sup> Yield refers to average isolated yield obtained in two or more experiments. <sup>c</sup> NaOtBu used in place of K<sub>3</sub>PO<sub>4</sub>.

In addition to greatly expanding the scope of Pd-catalyzed carboamination reactions involving aryl bromide substrates, the use of mildly basic reaction conditions also allows the first Pd-catalyzed carboamination reactions with aryl triflates.

(10) These conditions were also less effective for carboamination reactions of  $\gamma$ -(*N*-arylamino)alkenes; the desired *N*-arylpiperidine products were obtained in low to moderate yield (35–55%).

(7) In some cases use of DME as solvent provided results comparable to those obtained with dioxane (Table 2, entries 3, 4, 6, and 8.)

(8) The differences in reactivity between Pd<sub>2</sub>(dba)<sub>3</sub> and Pd(OAc)<sub>2</sub> are not fully understood, but may result from coordination of dba (dibenzylideneacetone) to the metal at key stages in the catalytic cycle, or through the formation of dba-ligated complexes that lie outside of the catalytic cycle. Dba has previously been demonstrated to have a large impact on the rates of Pd-catalyzed or -mediated process. For lead references, see: (a) Shekhar, S.; Ryberg, P.; Hartwig, J. F.; Mathew, J. S.; Blackmond, D. G.; Strieter, E. R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2006**, *128*, 3584–3591. (b) Amatore, C.; Broeker, G.; Jutand, A.; Khalil, F. *J. Am. Chem. Soc.* **1997**, *119*, 5176–5185.

(9) (a) Huang, P.-Q.; Wu, T.-J.; Ruan, Y.-P. *Org. Lett.* **2003**, *5*, 4341–4344. (b) Kadota, I.; Saya, S.; Yamamoto, Y. *Heterocycles* **1997**, *46*, 335–348. (c) Yoda, H.; Yamazaki, H.; Takabe, K. *Tetrahedron: Asymmetry* **1996**, *7*, 373–374.

Our preliminary efforts to conduct these transformations with the strong base NaOtBu were unsuccessful as a result of competing cleavage of the trifluoromethanesulfonate ester, which resulted in conversion of the aryl triflate to the corresponding phenol. For example, treatment of **3** with 4-formylphenyl triflate in the presence of catalytic Pd(OAc)<sub>2</sub>/Dpe-phos and stoichiometric NaOtBu failed to generate the desired pyrrolidine product **12**. However, subsequent experiments demonstrated that use of K<sub>3</sub>PO<sub>4</sub> as base provides the desired pyrrolidine **12** in 67% yield (Table 3, entry 1). These conditions are effective with both Boc- and Cbz-protected substrates, and diastereoselectivities are similar to those obtained in related reactions with aryl bromide electrophiles (entries 3 and 4).

In conclusion, we have developed new conditions for palladium-catalyzed carboamination reactions of *N*-protected  $\gamma$ -aminoalkenes with aryl bromides and triflates. These conditions, which use Cs<sub>2</sub>CO<sub>3</sub> or K<sub>3</sub>PO<sub>4</sub> in place of the strong

base NaOtBu, tolerate the presence of a broad array of functional groups and significantly expand the scope of this method. Applications of these new conditions to the synthesis of complex pyrrolidine alkaloids are currently being pursued.

**Acknowledgment.** The authors thank the NIH-NIGMS (GM071650) for financial support of this work. Additional support was provided by the Camille and Henry Dreyfus Foundation (New Faculty Award, Camille Dreyfus Teacher Scholar Award), Research Corporation (Innovation Award), Eli Lilly, Amgen, and 3M.

**Supporting Information Available:** Experimental procedures, spectroscopic data, and copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds reported in the text. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL062808F