


SmCl₃-Catalyzed C-Acylation of
1,3-Dicarbonyl Compounds and
MalononitrileQuansheng Shen,[†] Wen Huang,[†] Jialiang Wang,[†] and Xigeng Zhou^{*,†,‡}

Department of Chemistry, Shanghai Key Laboratory of Molecular Catalysis and Innovative Materials, Fudan University, Shanghai 200433, People's Republic of China, and State Key Laboratory of Organometallic Chemistry, Shanghai 200032, People's Republic of China

xgzhou@fudan.edu.cn

Received August 10, 2007

ABSTRACT



A recyclable, convenient, and efficient catalytic system for C-acylation of 1,3-dicarbonyl compounds and malononitrile with acid chlorides has been developed, giving moderate to excellent yields under mild conditions. This is the first catalytic example of such reactions. In addition, by applying this protocol as the key step, 3,5-disubstituted-1*H*-pyrazole-4-carboxylate can easily be synthesized in high yields in a one-pot procedure.

1,3,3'-Triketones are useful intermediates in organic synthesis, especially for the preparation of some biologically active compounds such as SR141716, phloroglucinols, and 5-deazaaminopterin.^{1,2} Surprisingly, only a limited number of procedures for the synthesis of 1,3,3'-triketones have been developed. In most cases, 1,3,3'-triketones are prepared by C-acylation of 1,3-dicarbonyl compounds.^{3–8} Nevertheless,

these methods usually require the use of stoichiometric amounts of strong bases such as EtONa,⁴ BaH₂,⁵ EtMgBr,⁶ *n*-BuLi,⁷ or powerful reductive metals like Na,⁸ which are not suitable for sensitive substrates. Notably, Rathke and Cowan developed a MgCl₂-promoted C-acylation of 1,3-dicarbonyl compounds.⁹ Although these possess many potential advantages, the main limitation of this strategy is that a stoichiometric amount of Lewis acid and 2 equiv of tertiary amine additive are required. This is because an acylation product **3** is always a stronger acid than the corresponding dicarbonyl precursor **2** and thus may neutralize a portion of the β-diketone intermediate (Scheme 1, route c). Therefore, the development of an alternative catalytic method for obtaining 1,3,3'-triketones represents a challenging but attractive subject from the viewpoints of operational simplicity, economy, and environmental impact.

[†] Fudan University.[‡] State Key Laboratory of Organometallic Chemistry.

(1) (a) Francisco, M. E.; Seltzman, H. H.; Gilliam, A. F.; Mitchell, R. A.; Rider, S. L.; Pertwee, R. G.; Stevenson, L. A.; Thomas, B. F. *J. Med. Chem.* **2002**, *45*, 2708. (b) Su, T. L.; Huang, J. T.; Chou, T. C.; Otter, G. M.; Sirotinak, F. M.; Watanabe, K. A. *J. Med. Chem.* **1988**, *31*, 1209.

(2) (a) Terada, M.; Nakano, M.; Ube, H. *J. Am. Chem. Soc.* **2006**, *128*, 16044. (b) Jung, M. E.; Min, S. J.; Houk, K. N.; Ess, D. *J. Org. Chem.* **2004**, *69*, 9085. (c) Lin, C. T.; Shih, J. H.; Chen, C. L.; Yang, D. Y. *Tetrahedron Lett.* **2005**, *46*, 5033.

(3) (a) Miyashita, A.; Suzuki, Y.; Nagasaki, I.; Ishiguro, C.; Iwamoto, K. I.; Higashino, T. *Chem. Pharm. Bull.* **1997**, *45*, 1254. (b) Sekine, M.; Kume, A.; Nakajima, M.; Hata, T. *Chem. Lett.* **1981**, 1087. (c) Tarbell, D. S.; Price, J. R. *J. Org. Chem.* **1956**, *21*, 144.

(4) Lawesson, S. O.; Busch, T. *Acta Chem. Scand.* **1989**, *8*, 1717.

(5) Lim, S.; Min, Y.; Choi, B.; Kim, D.; Lee, S. S.; Lee, I. M. *Tetrahedron Lett.* **2001**, *42*, 7645.


(6) Skarzewski, J. *Tetrahedron* **1989**, *14*, 4593.

(7) Jung, J. C.; Watkins, E. B.; Avery, M. A. *Synth. Commun.* **2002**, *32*, 3767.

(8) Peter, M.; Christo, I.; Marin, A. *Chem. Ber.* **1964**, *97*, 2987.

(9) (a) Rathke, M. W.; Cowan, P. J. *J. Org. Chem.* **1985**, *50*, 2622. (b) Little, R. D.; Russu, W. A. *J. Org. Chem.* **2000**, *65*, 8096.


Scheme 1. Mechanistic Hypothesis



Taking into account those results, we considered that one way to avoid the competitive deprotonation of the acylation product would be to make the coordination between metal and β -diketonate more stable than that of the tricarbonyl counterpart. Lanthanide trichlorides are known to activate β -diketone in the presence of bases, forming lanthanide β -diketonate complexes.¹⁰ Given the facts that the lanthanide–ligand bond strength predominantly depends on the electrostatic interaction and the degree of steric saturation, and that the introduction of an electron-withdrawing substituent could lead to decreased stabilization of lanthanide tri(β -diketonates),¹¹ we reasoned that it should be possible to control the preferential deprotonation of dicarbonyl compounds and intercept the resulting enolate intermediate with acyl chloride as an electrophile (Scheme 1, routes a and b). Herein, we wish to report a general, highly efficient, and catalytic method for C-acylation of 1,3-dicarbonyl compounds and malononitrile with acid chlorides.

Initially, we sought an effective catalytic system for the C-acylation of active methylene compounds, guided by the template reaction between benzoyl chloride **1a** and diethyl malonate **2a**. A range of reaction conditions were tested, and some of the results are listed in Table 1. As expected, lanthanide trichlorides proved to be efficient catalysts for the C-acylation of **1a** with **2a**. However, under the same conditions, for the other Lewis acids only very low yields of **3aa** were obtained even with longer reaction time (Table 1, entries 1–7). Among lanthanide trichlorides employed (Table 1, entries 8–14), the medium-sized SmCl_3 catalyzed the formation of **3aa** most efficiently. This is consistent with

Table 1. Optimization of the Reaction Conditions for Acylation of Diethyl Malonate with Benzoyl Chloride^a



entry	catalysts	solvent	time (h)	yield ^b (%)
1	MgCl_2	toluene	12	20
2	AlCl_3	toluene	10	24
3	FeCl_3	toluene	10	20
4	BiCl_3	toluene	12	18
5	InCl_3	toluene	12	32
6	H_3BO_3	toluene	12	21
7	$\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$	toluene	10	16
8	LaCl_3	toluene	5	52
9	NdCl_3	toluene	5	56
10	SmCl_3	toluene	4	87
11 ^c	SmCl_3	toluene	4	84
12	DyCl_3	toluene	5	78
13	ErCl_3	toluene	5	65
14	YbCl_3	toluene	5	50
15	$\text{Yb}(\text{OTf})_3$	toluene	12	55
16	none	toluene	12	0
17	SmCl_3	THF	4	57
18	SmCl_3	$n\text{-C}_6\text{H}_{14}$	4	68
19	SmCl_3	CH_3CN	4	trace
20	SmCl_3	$\text{ClCH}_2\text{CH}_2\text{Cl}$	4	58
21 ^d	SmCl_3	toluene	5	62

^a Reaction conditions: diethyl malonate (1.0 mmol), benzoyl chloride (1.1 mmol), catalyst (10 mol %), solvent (4 mL), Et_3N (1.2 mmol), rt. ^b LC yield based on 1,3-dicarbonyl compound. ^c Catalyst (5 mol %). ^d Pyridine as base (1.2 mmol).

the upward trend of the stability of lanthanide β -diketonate complexes,¹¹ demonstrating that the coordination of the diketone to lanthanide may play a key role in the equilibrium of the deprotonation of the acylation product and the substrate. No C-acylation reaction occurred in the absence of any catalyst (Table 1, entry 16).

A screening of solvents for the SmCl_3 system revealed that toluene is a suitable solvent, and triketone **3aa** was formed in 87% yield (Table 1, entry 10). Less activity in CH_3CN and with pyridine as a base indicate that increased basicity of solvent or base may lead to preferential coordination of these reagents over β -diketonates, which resulted in lower yields (entries 19 and 21). In the MgCl_2 catalytic system,⁹ where a stoichiometric amount of MgCl_2 and 2 equiv of tertiary amine additive were required, it was observed in the present case that 1 equiv of Et_3N was enough to effect complete conversion of the substrate in the presence of a catalytic amount of SmCl_3 .


Having determined the optimum reaction conditions, we investigated the generality of this process. As can be seen from Table 2, a variety of 1,3-dicarbonyl compounds can be C-acylated by **1a** to give the corresponding products in moderate to excellent yields, depending on the steric hindrance at the methylene carbon and the nature of carbonyl functional groups (Table 2, entries 1–8). Interestingly, cyclic 1,3-diketones can also be employed in this SmCl_3 -catalyzed

(10) Xu, G.; Wang, Z. M.; He, Z.; Lue, Z.; Liao, C. S.; Yan, C. H. *Inorg. Chem.* **2002**, *41*, 6802.

(11) (a) Tsukube, H.; Shinoda, S. *Chem. Rev.* **2002**, *102*, 2389. (b) Tsukube, H.; Shinoda, S.; Tamiaki, H. *Coord. Chem. Rev.* **2002**, *226*, 227. (c) Marks, T. J.; Ernst, R. D. In *Comprehensive Organometallic Chemistry*; Wilkinson, G., Stone, F. G., Abel, A. E. W., Eds.; Pergamon Press: Oxford, UK, 1982; Vol. 3, pp 173–270. (d) Evans, W. J. *Adv. Organomet. Chem.* **1985**, *24*, 77. (e) Chen, Y. L.; Li, X. S.; Zhang, H.; Ou, Y. M. *J. Rare Earth* **2004**, *25*, 87.

Until now, no example of the Lewis acid-catalyzed coupling of malononitrile with acid chlorides has been reported. After having established that SmCl_3 is an efficient catalyst for the inter- and intramolecular acylations of 1,3-dicarbonyl compounds, we were keen to explore the reaction of malononitrile with benzoyl chloride. SmCl_3 indeed led to the desired product **7** in good yield (Scheme 4), indicating

Scheme 4. SmCl_3 -Catalyzed Acylation of Malononitrile with Benzoyl Chloride



that SmCl_3 is also highly effective in the catalytic acylation of malononitrile with acid chlorides as substrates.

Pyrazole-4-carboxylate derivatives possess important pharmacological properties including analgesic, antipyretic, and antiinflammatory properties.¹³ Several methods to synthesize these compounds from 1,3-diketones have been reported in the literature.¹⁴ However, they usually involve multistep processes and give the products only in poor to moderate yields. Having successfully developed an efficient acylation of 1,3-dicarbonyl compounds with acid chlorides, we finally turned our attention to the application of this method to the one-pot synthesis of highly substituted pyrazole-4-carboxylate building blocks, which can be further functionalized. Treatment of 1,3-dicarbonyl compounds **2** with acid chlorides **1** in the presence of 5 mol % of SmCl_3 and 1.2 equiv of Et_3N followed by reacting with hydrazine allowed the isolation of multisubstituted pyrazoles **8–11** in 72–81%


(12) (a) Liao, Y. X.; Kuo, P. Y.; Yang, D. Y. *Tetrahedron Lett.* **2003**, 44, 1599. (b) Mao, P. C. M.; Mouscadet, J. F.; Leh, H.; Auclair, C.; Hsu, L. Y. *Chem. Pharm. Bull.* **2002**, 50, 1634. (c) Staunton, J. In *Comprehensive Organic Chemistry*; Sammes, P. G., Ed.; Pergamon: Oxford, UK, 1979; Vol. 4.

(13) Moro, S.; Varanob, F.; Cozzac, G.; Paganoc, M. A.; Zagotto, G.; Chilina, A.; Guiotto, A.; Catarzib, D.; Calottab, V.; Pinnac, L. A.; Meggioc, F. *Lett. Drug. Des. Discovery* **2006**, 3, 281.

(14) (a) Morelli, C. F.; Saladino, A.; Speranza, G.; Manitto, P. *Eur. J. Org. Chem.* **2005**, 21, 4621. (b) Vanier, C.; Wagner, A.; Mioskowski, C. *J. Comb. Chem.* **2004**, 6, 846.

yields. This provides a mild and straightforward route to pyrazole-4-carboxylate derivatives (Scheme 5).

Scheme 5. SmCl_3 -Catalyzed One-Pot Synthesis of Pyrazole-4-carboxylate



Finally, we checked the reusability of the SmCl_3 catalyst in the reaction of 8.8 mmol of **1a** with 8 mmol of **2c** in toluene at room temperature. After the first experiment, the insoluble catalyst was readily recovered by filtrating the reaction mixture and then removing amine salt washed by acetonitrile for the next use. The results of seven runs showed that the recovered catalyst retains its activity in terms of yields (86%, 82%, 83%, 85%, 84%, 85%, and 82%, respectively) and rates.

In summary, a general and highly efficient LnCl_3 -catalyzed intra- and intermolecular *C*-acylation of 1,3-dicarbonyl compounds and malononitrile with acid chlorides has been developed. This method provides significant advantages in cost, tolerance of functional groups, and simplicity of operation by obviating the use of strong bases and stoichiometric amounts of Lewis acids. Furthermore, the present catalyst is readily recovered and reused at least seven times for such reaction without visible loss of catalytic activity. Further investigations on the mechanism details and synthetic applications of this method are underway in our lab.

Acknowledgment. We thank the National Natural Science Foundation of China and the Natural Science Foundation of Shanghai.

Supporting Information Available: Experimental details and characterization of the compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL701961Z