

Enantioselective Synthesis of Aryl Sulfoxides via Palladium-Catalyzed Arylation of Sulfenate Anions


Guillaume Maitro, Sophie Vogel, Mounel Sadaoui, Guillaume Prestat, David Madec,* and Giovanni Poli*

Université Pierre et Marie Curie–Paris 6, Laboratoire de Chimie Organique (UMR CNRS 7611), Institut de Chimie Moléculaire (FR 2769), Case 183, 4 Place Jussieu, F-75252 Paris Cedex 05, France

giovanni.poli@upmc.fr; madec@ccr.jussieu.fr

Received October 7, 2007

ABSTRACT



Arylation of various sulfenate anions generated from β -sulfinyl esters by retro-Michael reaction in the presence of palladium(0) and enantiopure ligands gave the corresponding aryl sulfoxides in enantio-enriched form. The Josiphos-type ligand (*R*)-(*S*)-PPF-*t*-Bu₂ turned out to be the best ligand tested, allowing ee's up to 83% in a predictable sense.

Enantiopure sulfoxides represent an important class of compounds because of their usefulness as chiral auxiliaries or as chiral ligands in asymmetric catalysis.¹ Moreover, a number of sulfoxides are known in the field of pharmaceutical and medicinal chemistry.² As a consequence, the synthesis of chiral nonracemic sulfoxides has been of high interest for several decades.³ Two different methods are commonly employed for the preparation of this type of compounds: (a) the addition of organometallic nucleophiles, such as Grignard reagents, to enantiopure sulfonates having a stereogenic sulfur atom,^{4,5} and (b) the enzymatic or catalytic asymmetric

oxidation of unsymmetrical thioethers,⁶ such as the modified Sharpless procedures described by Kagan or Modena's groups.⁷ Despite their value, these two methods suffer from some limitations such as the use of a stoichiometric amount of chiral auxiliary in the preparation of enantiopure sulfinate precursors and high substrate-dependency of the asymmetric oxidation.

On the other hand, palladium-catalyzed arylation represents a primary synthetic tool to generate carbon-heteroatom bonds.⁸ In particular, the use of sulfur-based nucleophiles⁹

(1) Fernandez, I.; Khiar, N. *Chem. Rev.* **2003**, *103*, 3651–3705.

(2) For reviews, see: (a) Carrenõ, M. C. *Chem. Rev.* **1995**, *95*, 1717–1760. (b) Legros, J.; Dehli, J. R.; Bolm, C. *Adv. Synth. Catal.* **2005**, *347*, 19–31. (c) Bentley, R. *Chem. Soc. Rev.* **2005**, *34*, 609–624.

(3) Khiar, N.; Fernandez, I.; Alcudia, A.; Alcudia, F. In *Advances in Sulfur Chemistry*; Rayner, C. M., Ed.; JAI Press, Inc.: Stanford, 2000; Vol. 2, p 57.

(4) (a) Andersen, K. K. *Tetrahedron Lett.* **1962**, *3*, 93–95. (b) Andersen, K. K.; Gafield, W.; Papanikolaou, N. E.; Foley, J. W.; Perkins, J. I. *J. Am. Chem. Soc.* **1964**, *86*, 5637–5646. (c) Mioskowski, C.; Solladié, G. *Tetrahedron* **1980**, *35*, 227–236. (d) Solladié, G.; Hutt, J.; Girardin, A. *Synthesis* **1987**, 173.

(5) For selected recent examples, see: (a) Khiar, N.; Araujo, C. S.; Alcudia, F.; Fernández, I. *J. Org. Chem.* **2002**, *67*, 345–356. (b) Delouvrié, B.; Fensterbank, L.; Nájera, F.; Malacria, M. *Eur. J. Org. Chem.* **2002**, 3507–3525. (c) Ruano, J. L. G.; Alemparte, C.; Aranda, M. T.; Zarzuelo, M. M. *Org. Lett.* **2003**, *5*, 75–78. (d) Lu, B. Z.; Jin, F.; Zhang, Y.; Wu, X.; Wald, S. A.; Senanayake, C. H. *Org. Lett.* **2005**, *7*, 1465–1468.


(6) For selected recent examples, see: (a) Kowalski, P.; Mitka, K.; Ossowska, K.; Kolarska, Z. *Tetrahedron* **2005**, *61*, 1933–1953. (b) Kaczorowska, K.; Kolarska, Z.; Mitka, K.; Kowalski, P. *Tetrahedron* **2005**, *61*, 8315–8327. (c) Madec, D.; Mingoia, F.; Macovei, C.; Maitro, G.; Giambastiani, G.; Poli, G. *Eur. J. Org. Chem.* **2005**, 552–557. (d) Scarso, A.; Strukul, G. *Adv. Synth. Catal.* **2005**, *347*, 1227–1234. (e) Drago, C.; Caggiano, L.; Jackson, R. F. W. *Angew. Chem., Int. Ed.* **2005**, *44*, 7221–7223. (f) Egami, H.; Katsuki, T. *J. Am. Chem. Soc.* **2007**, *129*, 8940–8941. (g) Yamaguchi, T.; Matsumoto, K.; Saito, B.; Katsuki, T. *Angew. Chem., Int. Ed.* **2007**, *46*, 4729–4731. (h) Sun, J.; Zhu, C.; Dai, Z.; Yang, M.; Pan, Y.; Hu, H. *J. Org. Chem.* **2004**, *69*, 8500–8503. (i) del Río, R. E.; Wang, B.; Achab, S.; Bohé, L. *Org. Lett.* **2007**, *9*, 2265–2268.

(7) (a) Pitchen, P.; Dunach, E.; Deshmukh, M. N.; Kagan, H. B. *J. Am. Chem. Soc.* **1984**, *106*, 8188–8193. (b) Pitchen, P.; Kagan, H. B. *Tetrahedron Lett.* **1984**, *25*, 1049–1052. (c) DiFuria, F.; Modena, G.; Seraglia, R. *Synthesis* **1984**, 325–326. (d) Brunel, J. M.; Duetsch, M.; Kagan, H. B. *J. Org. Chem.* **1995**, *60*, 8086–8088. (e) Brunel, J. M.; Kagan, H. B. *Synlett* **1996**, 404–406. (d) Kagan, H. B. In *Catalytic Asymmetric Synthesis*, 2nd ed.; Ojima, I., Ed.; Wiley-VCH: New-York, 2000; Chapter 6c.

such as sulfonates¹⁰ and thiolates¹¹ allows an easy generation of aryl sulfones and aryl sulfides, respectively. In this context, we recently described the synthesis of aryl sulfoxides¹² by palladium-catalyzed arylation of sulfenate anions.¹³


This original coupling reaction has been successfully achieved by generation of the desired sulfenate¹⁴ anion via β -sulfinyl esters enolate elimination¹⁵ under specifically developed biphasic conditions (Scheme 1).¹⁶

Scheme 1



From this result, we reasoned that, if suitable enantiopure ligands were used for the coupling, the prostereogenic nature of the sulfenate anion could disclose a new route toward enantiomerically enriched aryl sulfoxides (Scheme 2).¹⁷ Yet,

Scheme 2



such a strategy appeared non-obvious, as our previous study demonstrated that achiral Xantphos was the only ligand able to promote the coupling reaction.¹²

The coupling reaction between the in situ generated *p*-tolyl sulfenate anion and *p*-iodoanisole was chosen as the model reaction, using toluene/H₂O biphasic conditions (Table 1).

Table 1. Optimization of Reaction Conditions^a

entry	ligand	base	solvent	temp (°C)	yield (%) ^b	% ee ^c (config)
1	Xantphos ^d	KOH	tol/H ₂ O	70	84	
2	Binap ^e	KOH	tol/H ₂ O	70	0	
3	(+)-Diop ^f	KOH	tol/H ₂ O	70	42	8 (R)
4	A	KOH	tol/H ₂ O	70	13	29 (R)
5	B	KOH	tol/H ₂ O	70	55	83 (R)
6	C	KOH	tol/H ₂ O	70	0	
7	D	KOH	tol/H ₂ O	70	0	
8	E	KOH	tol/H ₂ O	70	0	
9	F	KOH	tol/H ₂ O	70	54	25 (S)
10	G	KOH	tol/H ₂ O	70	30	12 (S)
11	H	KOH	tol/H ₂ O	70	0	
12	B	Cs ₂ CO ₃	tol	110	83	73 (R)

^a Reagents and reaction conditions: *p*-iodoanisole (1.2 equiv), β -sulfinylester, Pd₂dba₃ (1 mol %), ligand (2 mol %), KOH (50% aqueous solution, 20 equiv), 1:1 toluene/H₂O, 70 °C or Cs₂CO₃ (4 equiv), at reflux of toluene. ^b Yields are given for isolated products. ^c Enantiomeric excess was determined by chiral supercritical fluid chromatography. ^d Xantphos: 4,5-bis(phenylphosphino)-9,9-dimethylxanthene. ^e Binap: 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl. ^f (+)-Diop: (+)-O-isopropylidene-*trans*-2,3-dihydroxy-1,4-bis(diphenyl-phosphino)butane.

Preliminary experiments performed with Binap and Diop ligands were disappointing. Indeed, with the former ligand no sulfoxide was observed (entry 2), whereas Diop allowed generation of the expected coupling product in a moderate (42%) yield and a poor (8%) ee (entry 3). After the beginning of our project, Hartwig and co-workers reported the palladium-catalyzed coupling reaction of aryl halides with thiols using enantiopure Josiphos **A** as the ligand.^{11c} Although in Hartwig's paper asymmetric catalysis was not addressed, this result prompted us to test Josiphos-type ligands in our coupling reactions.

Encouragingly, use of Josiphos **A** as the ligand allowed access to *p*-tolylsulfinyl anisole with 29% ee albeit with a very poor yield (13%) (entry 4).

On the other hand, use of ligand **B** allowed generation of the corresponding sulfoxide¹⁸ in 55% yield and a satisfactory 83% ee (entry 5). Replacement of Josiphos with the Josiphos-type ligands **C**–**E** or Mandiphos-, Taniaphos-, and Walphos-type ligands **F**–**H**, respectively (Figure 1), increased neither the yield nor the enantioselectivity (entries 6–11). Finally, optimized conditions were set with ligand **B** (Cs₂CO₃ as base

(8) Prim, D.; Campagne, J.-M.; Joseph, D.; Andrioletti, B. *Tetrahedron* **2002**, *58*, 2041–2075.

(9) Kondo, T.; Mitsudo, T.-A. *Chem. Rev.* **2000**, *100*, 3205–3220.

(10) (a) Cacchi, S.; Fabrizi, G.; Goggiani, A.; Parisi, L. M.; Bernini, R. *J. Org. Chem.* **2004**, *69*, 5608–5614. (b) Cacchi, S.; Fabrizi, G.; Goggiani, A.; Parisi, L. M. *Org. Lett.* **2002**, *4*, 4719–4721.

(11) For pioneering works, see: (a) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1389. (b) Foà, M.; Santi, R.; Garavalia, F. *J. Organomet. Chem.* **1981**, *206*, C29–C32. For more recent examples, see: (c) Fernández-Rodríguez, M. A.; Shen, Q.; Hartwig, J. F. *J. Am. Chem. Soc.* **2006**, *128*, 2180–2181. (d) Fernández-Rodríguez, M. A.; Shen, Q.; Hartwig, J. F. *Chem. Eur. J.* **2006**, *12*, 7582–7596. (e) Itoh, T.; Mase, T. *J. Org. Chem.* **2006**, *71*, 2203–2206. (f) Moreau, X.; Campagne, J.-M.; Meyer, G.; Jutand, A. *Eur. J. Org. Chem.* **2005**, 3749–3760. (g) Mispelaere-Canivet, C.; Spindler, J.-F.; Perrio, S.; Beslin, P. *Tetrahedron* **2005**, *61*, 5253–5259. (h) Itoh, T.; Mase, T. *Org. Lett.* **2004**, *6*, 4587–4590. (i) Murata, M.; Buchwald, S. L. *Tetrahedron* **2004**, *60*, 7397–7403.

(12) (a) Maitro, G.; Vogel, S.; Prestat, G.; Madec, D.; Poli, G. *Org. Lett.* **2006**, *8*, 5951–5954. See also: (b) Colobert, F.; Ballesteros-Garrido, R.; Leroux, F. R.; Ballesteros, R.; Belén, A. *Tetrahedron Lett.* **2007**, *48*, 6896–6899.

(13) For reviews on the chemistry of sulfenate anions, see: (a) O'Donnell, J. S.; Schwan, A. L. *J. Sulfur Chem.* **2004**, *25*, 183–211. (b) Hogg, D. R. *Chemistry of Sulfenic Acids and Esters. The Chemistry of Sulfenic Acids and their Derivatives*; Wiley-Interscience: Toronto, 1990; pp 361–402. (c) Drabowicz, J.; Kielbasinski, P.; Mikolajczyk, M. *The Chemistry of Sulfenic Acids, Esters and their Derivatives*; Patai, S., Ed.; John Wiley & Sons: New York, 1990; pp 351–429.

in 75% or 93% yields and in 72% or 65% ee's, respectively (entries 7 and 8). Finally, benzylsulfenate precursor **1c** gave the aryl benzylsulfoxides **2c**²³ or **3c**²⁴ in 67% or 71% yields and 40% or 47% ee's, respectively (entries 9 and 10). It is noteworthy for predictive assignment that ligand **B** constantly favors sulfur-carbon bond formation via the *pro*-S sulfenate anion lone pair (Scheme 2).

In conclusion, we have reported the first asymmetric transition-metal-catalyzed arylation of sulfenate anions, thereby disclosing a new synthetic route toward enantio-merically enriched aryl sulfoxides. The Josiphos-type ligand

(23) Donnoli, M. I.; Superchi, S.; Rosini, C. *J. Org. Chem.* **1998**, 63, 9392–9395.

(24) Kelly, P.; Lawrence, S. E.; Maguire, A. R. *Eur. J. Org. Chem.* **2006**, 4500–4509.

B turned out to be the best ligand tested, allowing ee's up to 83% and in a predictable sense. Studies to elucidate the details of reaction mechanism as well as the origin of the enantioselectivity are currently in progress.

Acknowledgment. We dedicate this paper in memory of Dr. Charles Mioskowski. CNRS and UPMC are acknowledged for financial support. The sponsorship of COST Action D40 "Innovative Catalysis: New Processes and Selectivities" is kindly acknowledged.

Supporting Information Available: General procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL702343G