

Highly Ortho-Selective Cross-Coupling of Dichlorobenzene Derivatives with Grignard Reagents


Shunpei Ishikawa and Kei Manabe*

RIKEN, 2-1 Hirosawa, Wako 351-0198, Japan

keimanabe@riken.jp

Received November 1, 2007

ABSTRACT



Highly ortho-selective cross-coupling of dichlorobenzene derivatives with Grignard reagents was realized using a combination of Pd₂(dba)₃ and PCy₃. Use of hydroxylated terphenylphosphines further improved the reactions of dichlorophenol and dichloroaniline.

Cross-coupling of haloarenes with organometallic reagents constitutes one of the most important and practical reactions among transition-metal-catalyzed C–C bond formations.¹ However, when arene molecules have more than one substituent of the same halogen atom, transition-metal-catalyzed site-selective cross-coupling involving site-selective conversion of one of the halogen atoms to another group remains relatively unexplored. For dihalogenated heteroarenes, many examples of site-selective cross-coupling exist.² In contrast, only a few examples have been reported for dihalogenated benzene derivatives.³ Although the reactions reported can be useful, developing other strategies for site selection is of great interest.

Recently, we reported ortho-selective cross-coupling of dibromophenols or dibromoanilines with Grignard reagents in the presence of Pd catalysts.⁴ Reaction selectivity is controlled by catalysts generated from hydroxylated terphenylphosphine ligands such as **1** and **2**.^{5,6} Other phosphines did not promote the ortho-selective reactions. It is worth noting that these reactions occurred at positions ortho to highly electron-donating groups such as -OMgBr and

-NHMgBr. This ortho-selectivity was not expected from the previous examples of site-selective cross-coupling.^{7,8} To expand the scope of coupling, we next investigated reactions of dichlorobenzene derivatives and found that the dichloro derivatives showed behavior different from dibromo deriva-

(3) (a) Singh, R.; Just, G. *J. Org. Chem.* **1989**, *54*, 4453. (b) Ishii, Y.; Chatani, N.; Yorimitsu, S.; Murai, S. *Chem. Lett.* **1998**, 27, 157. (c) Dirk, S. M.; Price, D. W., Jr.; Chanteau, S.; Kosynkin, D. V.; Tour, J. M. *Tetrahedron* **2001**, *57*, 5109. (d) Schaub, T.; Backes, M.; Radius, U. *J. Am. Chem. Soc.* **2006**, *128*, 15964. (e) Ackermann, L.; Althammer, A. *Angew. Chem., Int. Ed.* **2007**, *46*, 1627. (f) Houpi, I. N.; Hoeck, J.-P. V.; Tilstam, U. *Synlett* **2007**, 2179. MeO as a leaving group: (g) Kakiuchi, F.; Usui, M.; Ueno, S.; Chatani, N.; Murai, S. *J. Am. Chem. Soc.* **2004**, *126*, 2706. For an example of selective cross-coupling of 1,2-dihaloalkenes, see: (h) Bellina, F.; Anselmi, C.; Viel, S.; Mannina, L.; Rossi, R. *Tetrahedron* **2001**, *57*, 9997. For early examples of selective cross-coupling of 1,1-dihaloalkenes, see: (i) Minato, A.; Suzuki, K.; Tamao, K. *J. Am. Chem. Soc.* **1987**, *109*, 1257. (j) Roush, W. R.; Brown, B. B.; Drozda, S. E. *Tetrahedron Lett.* **1988**, 29, 3541. For enantioselective cross-coupling, see: (k) Hayashi, T.; Niizuma, S.; Kamikawa, T.; Suzuki, N.; Uozumi, Y. *J. Am. Chem. Soc.* **1995**, *117*, 9101. Cu-catalyzed intramolecular etherification of bisvinylbromides: (l) Fang, Y.; Li, C. *J. Am. Chem. Soc.* **2007**, *129*, 8092.

(4) Ishikawa, S.; Manabe, K. *Chem. Lett.* **2007**, 36, 1304.


(5) (a) Ishikawa, S.; Manabe, K. *Chem. Lett.* **2007**, 36, 1302. Ligand **1** was purified as its HBF₄ salt, and the salt was used directly in coupling reactions in which the free phosphine was liberated. For an HBF₄ salt, see: (b) Netherton, M. R.; Fu, G. C. *Org. Lett.* **2001**, 3, 4295.

(6) Ligands **1** and **2** were designed on the basis of biphenylphosphines such as **11** developed by Buchwald et al. See: Wolfe, J. P.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **1999**, *38*, 2413.

(1) *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed.; de Meijere, A., Diederich, F., Eds.; Wiley-VCH: Weinheim, 2004.

(2) A review: Schröter, S.; Stock, C.; Bach, T. *Tetrahedron* **2005**, *61*, 2245.

tives. Here, we reveal a new type of site-selective cross coupling in which complete ortho-selectivity was realized even with a simple phosphine, tricyclohexylphosphine (PCy₃). Moreover, use of **1** or **2** further improved reactions of dichlorophenol and dichloroaniline.



The reaction of 2,4-dichlorophenol (**4**) with 4-methoxyphenyl Grignard reagent was examined first.⁹ Remarkably, the catalyst generated from Pd₂(dba)₃ and PCy₃ afforded the ortho-arylated compound in good yield (Table 1, entries 2

Table 1. Ortho-Selective Cross-Coupling of Dihalobenzene Derivatives

entry	dihalobenzene derivatives	Grignard reagent	reaction time (h)	yield (%)		
				ortho	isomer	di
1 ^a	3	4-MeOC ₆ H ₄ MgBr	24	21	14	38
2	4	4-MeOC ₆ H ₄ MgBr	4	80	0	0
3	4	4-MeOC ₆ H ₄ MgBr	18	91	0	0
4	4	PhMgCl	4	85	0	0
5	4	2-ThienylMgBr ^b	20	52	0	0
6 ^c	5	4-MeOC ₆ H ₄ MgBr	18	95	0	0
7	6	4-MeOC ₆ H ₄ MgBr	4	1	0	0
8	7	4-MeOC ₆ H ₄ MgBr	18	99	0	0
9 ^c	8	4-MeOC ₆ H ₄ MgBr	18	87	0	0
10	9	4-MeOC ₆ H ₄ MgBr	4	71	0	0
11 ^c	10	4-MeOC ₆ H ₄ MgBr	8	55	0	0

^a At 25 °C with 4 equiv of the Grignard reagent. ^b 4 equiv. ^c Pd₂(dba)₃ (2 mol %) and PCy₃ (4.8 mol %) were used.

and **3**). Neither the isomeric monoarylated compound (**isomer**) nor the diarylated compound (**di**) was obtained. This


(7) In most of the reported examples, the reactions were mainly controlled by the electronic property of the substrates: the reactions occur preferentially at less electronically negative carbons. For example, selective Sonogashira coupling of dibromoanilines, which has an electron-donating NH₂ group, occurred preferentially at the meta-position over the ortho- or para-position.^{3a} See also: Fahey, D. R. *J. Am. Chem. Soc.* **1970**, 92, 402.

(8) Electron-withdrawn groups such as nitro and carboxylate groups are known to facilitate cross-coupling at the ortho-position. This effect has been attributed to coordination of these groups to Pd in the oxidative addition of the *o*-C–X bond: (a) Kim, Y. M.; Yu, S. *J. Am. Chem. Soc.* **2003**, 125, 1696. (b) Bahmanyar, S.; Borer, B. C.; Kim, Y. M.; Kurtz, D. M.; Yu, S. *Org. Lett.* **2005**, 75, 1011. (c) Widdowson, D. A.; Wilhelm, R. *Chem. Commun.* **2003**, 578. See also ref 3b.


(9) Reviews on cross-coupling with Grignard reagents: (a) Tsuji, J. *Palladium Reagents and Catalysts*; John Wiley & Sons: West Sussex, 2004; p 335. (b) Cepanec, I. *Synthesis of Biaryls*; Elsevier: Oxford, 2004, p 83.

result is contrary to the reaction of dibromophenol, which produced significant amounts of **isomer** and **di** (entry 1).⁴ Formation of diarylated compounds often are unavoidable in cross-coupling reactions,¹⁰ and even in our previous ortho-selective reactions of dibromobenzene derivatives using **1**,⁴ complete suppression of diarylation could not be achieved. Thus, the absence of **di** is worth noting. The complete ortho-selectivity was also observed for compound **5** (entry 6). The result is worth mentioning because cross-coupling occurred at more electronically negative and more sterically hindered carbons. Compound **6** was not reactive under these conditions, probably because of the steric hindrance of *o*-Cl (entry 7). Not only the OH group but also other directing groups such as CH₂OH, NH₂, NHAc, and NHBoc worked well to produce ortho-arylated products (entries 8–11).¹¹

The reaction of 2,4-dichloroanisole under the same conditions did not give the cross-coupled products as shown in eq 1. This result strongly suggests that the presence of a



protic group in the substrates is essential for rate acceleration at the ortho-position. The requirement of a protic group seems counterintuitive, because the presence of an anionic substituent formed by deprotonation with a Grignard reagent should retard oxidative addition due to the strong electron-donating ability.¹² To explain the acceleration at the ortho-position, we propose transition state **A**, in which Lewis acidic



Mg facilitates oxidative addition of the *o*-C–Cl bond to Pd.¹³ On the other hand, we cannot exclude the possibility that

(10) (a) Tamao, K.; Sumitani, K.; Kiso, Y.; Zembayashi, M.; Fujioka, A.; Kodama, S.-i.; Nakajima, I.; Minato, A.; Kumada, M. *Bull. Chem. Soc. Jpn.* **1976**, 49, 1958. See also: (b) Sinclair, D. J.; Sherburn, M. S. *J. Org. Chem.* **2005**, 70, 3730. (c) Dong, C.-G.; Hu, Q.-S. *J. Am. Chem. Soc.* **2005**, 127, 10006.

(11) Ortho-selective Suzuki–Miyaura coupling of these substrates was not successful. For example, a reaction of **4** with 4-MeOC₆H₄B(OH)₂ in the presence of Pd₂(dba)₃ (2 mol %), PCy₃ (4.8 mol %), and KF (3 equiv) in THF at 70 °C for 21 h gave **ortho** (32%), **isomer** (14%), and **di** (12%). For Suzuki–Miyaura coupling using these conditions, see: Littke, A. F.; Dai, C.; Fu, G. C. *J. Am. Chem. Soc.* **2000**, 122, 4020.


(12) Examples of cross-coupling of halophenols with Grignard reagents: (a) Jendralla, H.; Chen, L.-J. *Synthesis* **1990**, 827. (b) Bumagin, N. A.; Luzikova, E. V. *J. Organomet. Chem.* **1997**, 532, 271. (c) Huang, J.; Nolan, S. P. *J. Am. Chem. Soc.* **1999**, 121, 9889. An example of Suzuki–Miyaura coupling of 2-halophenols: (d) Wawrzyniak, P.; Heinicke, J. *Tetrahedron Lett.* **2006**, 47, 8921.


(13) Activation of C–X bonds by coordination to Mg was reported by Nakamura et al. in their Ni-catalyzed cross coupling of ArF or ArCl with Grignard reagents: Yoshikai, N.; Mashima, H.; Nakamura, E. *J. Am. Chem. Soc.* **2005**, 127, 17978.

coordination of the anionic directing groups toward Pd stabilizes transition states of ortho-oxidative addition,⁸ especially for **7**, **9**, and **10**, which might form five- or six-membered rings by coordination of the anionic oxygen atom to Pd.


To further improve the catalytic system, we used ligands **1** and **2** instead of PCy₃ (Table 2).¹⁴ As expected, **1** and **2**

Table 2. Ortho-Selective Cross-Coupling Using **1**, **2**, and Reference Ligands

					
entry	Y	Grignard reagent	ligand	reaction time (h)	yield (%)
1	OH (4)	4-MeOC ₆ H ₄ MgBr	1	2	88
2	OH (4)	4-MeOC ₆ H ₄ MgBr	1	4	99
3	OH (4)	4-MeOC ₆ H ₄ MgBr	2	2	84
4	OH (4)	4-MeOC ₆ H ₄ MgBr	11	4	16
5	OH (4)	4-MeOC ₆ H ₄ MgBr	12	4	16
6	NH ₂ (8)	4-MeOC ₆ H ₄ MgBr	1	4	93
7	OH (4)	2-ThienylMgBr ^a	1	20	73
8	OH (4)	4-ClC ₆ H ₄ MgBr ^b	1	13	63
9 ^c	OH (4)	Me ₂ C=CHMgBr	1	24	66
10	OH (4)	BuMgCl	1	24	0



11



12


^a 4 equiv. ^b A solution in Et₂O was used. ^c Pd₂(dba)₃ (2 mol %) and **1** (4.8 mol %) were used.

accelerated ortho-selective reaction compared with PCy₃ (entries 1–3 vs Table 1, entry 2). Similar ligands such as **11**⁶ and **12**^{5a} did not exhibit high activity (entries 4 and 5), indicating that the presence and the position of the OH group of **1** were important for the high activity. Use of **1** was also beneficial for reaction of dichloroaniline **8** (entry 6 vs Table 1, entry 9). Other aryl and alkenyl Grignard reagents also worked well (entries 7–9). Neither para-arylated isomers nor the diarylated compounds were obtained in any of the reactions. Note that a Grignard reagent with a chloro substituent afforded the desired product in an acceptable yield (entry 8), while small amounts of oligomeric byproducts were formed. Alkyl Grignard reagents such as BuMgCl did not afford the desired product (entry 10); instead, a large amount


(14) Nakamura et al. also used a phosphine with a hydroxy group for Ni-catalyzed cross-coupling. See ref. 13.

of 4-chlorophenol was produced. For substrates **7**, **9**, and **10**, the acceleration effect of **1** was not observed (data not shown), with PCy₃ better than **1**.

Although the mechanism of acceleration by **1** and **2** is unclear, we speculate formation of transition state **B**, in which coordination of the oxido group of **1** or **2** to the Mg cation places the Pd atom close to the *o*-C–Cl bond. Since **7**, **9**, and **10** were poor substrates and **12** was a poor ligand, subtle changes in the position of Mg and the oxido group may be detrimental to the proper arrangement of the reacting species.



Finally, we demonstrated effectiveness of the hydroxylated phosphines for reaction of 4-bromo-2-chlorophenol (eq 2). Surprisingly, the Cl group at the ortho-position reacted preferentially over the Br group. This unusual selectivity was not observed in the reaction with PCy₃.



In summary, we have developed a method for achieving ortho-selective cross-coupling of dichlorobenzene derivatives with Grignard reagents. Combining use of Pd(0) and PCy₃ realized high selectivity for substrates with directing groups such as OH, CH₂OH, NH₂, NHAc, and NHBoc. Furthermore, ligands **1** and **2** were more effective than PCy₃ for dichlorophenol and dichloroaniline. Although the mechanism of selectivity remains unknown, the reactions described here provide new routes to synthesize multisubstituted benzene derivatives.

Acknowledgment. This work was partly supported by a Grant-in-Aid for Scientific Research on Priority Areas “Advanced Molecular Transformations of Carbon Resources” from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Supporting Information Available: Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL702646S