


Hemi-Phorboxazole A: Structure
Confirmation, Analogue Design and
Biological EvaluationAmos B. Smith III,^{*,†} Zhuqing Liu,[†] Anne-Marie L. Hogan,[†] Doralyn S. Dalisay,[‡]
and Tadeusz F. Molinski[‡]

Department of Chemistry, Laboratory for Research on the Structure of Matter,
and Monell Chemical Senses Center, University of Pennsylvania,
Philadelphia, Pennsylvania 19104, and Department of Chemical and Biochemistry and
Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California,
San Diego, 9500 Gilman Drive, MC0358, La Jolla, California 92093

smithab@sas.upenn.edu

Received June 24, 2009

ABSTRACT



A synthesis providing totally synthetic (+)-hemi-phorboxazole A (1), proceeding in two steps (85% yield) from known vinyl iodide precursor (+)-2, has been achieved in conjunction with the design, synthesis, and biological evaluation of two hemi-phorboxazole analogues [(+)-3 and (–)-4] featuring ring replacements inscribed within the macrolide. Although hemi-phorboxazole A (1) displayed no activity when tested against *Candida albicans* and two human cancer cell lines, analogue (–)-4 exhibited significant tumor cell growth inhibitory activity in the nanomolar range against HCT-116 (colon) and SK-BR-3 (breast), while (+)-3 displayed promising antifungal activity against *C. albicans*.

In April 2009, Dalisay and Molinski reported¹ the isolation of hemi-phorboxazole A (1), a diminutive form of (+)-phorboxazole A (5), derived from the same marine sponge (*Phorbas* sp.) that not only provides phorboxazoles A and B (5 and 6)² but also the unrelated phorbosides A–F (7–12)³

and very recently muironolide A (13)⁴ (Figure 1). Remarkably, the complete structure including absolute configuration was assigned with only 16.5 μ g (!) of natural material, exploiting state-of-the-art microcryoprobe NMR, in conjunction with HRMS and circular dichroism (CD) studies. The truncated hemi-phorboxazole A (1), the first reported example of a natural phorboxazole variant since the original Molinski et al. disclosure of the phorboxazoles in 1995,² was found to be ~ 10000 times less abundant than phorboxazoles A and B (cf. 0.07 vs 400 ppm, respectively). Not surprisingly, the limited availability of the natural material precluded any significant evaluation of the biological activity. Thus, to confirm the assigned structure of 1, as well as to provide

[†] University of Pennsylvania.[‡] University of California.(1) Dalisay, D. S.; Molinski, T. F. *Org. Lett.* **2009**, *11*, 1967.(2) (a) Searle, P. A.; Molinski, T. F. *J. Am. Chem. Soc.* **1995**, *117*, 8126.(b) Searle, P. A.; Molinski, T. F.; Brzezinski, L. J.; Leahy, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 9422. (c) Molinski, T. F. *Tetrahedron Lett.* **1996**, *37*, 7879.(3) (a) MacMillan, J. B.; Xiong-Zhou, G.; Skepper, C. K.; Molinski, T. F. *J. Org. Chem.* **2008**, *73*, 3699. (b) Skepper, C. K.; MacMillan, J. B.; Zhou, G.-X.; Masuno, M. N.; Molinski, T. F. *J. Am. Chem. Soc.* **2007**, *129*, 4150. (c) Dalisay, D. S.; Molinski, T. F. *J. Nat. Prod.* **2009**, *72*, 739.(4) Dalisay, D. S.; Morinaka, B. I.; Skepper, C. K.; Molinski, T. F. *J. Am. Chem. Soc.* **2009**, *131*, 7552.




Figure 1. Natural products isolated from *Phorbas* sp.


material to define the biological properties, recourse to total synthesis was clearly required for forward progress.

Given that our laboratory has had a long-standing interest in the phorboxazole area, with first- and second-generation total syntheses of (+)-phorboxazole A (**5**) recorded in 2001⁵ and 2005,⁶ we were intrigued with the diminutive form of hemi-phorboxazole A (**1**). Indeed, a synthetic program directed at hemi-phorboxazole A could take ready advantage of the Petasis–Ferrier union/rearrangement,⁷ an effective tactic specifically designed and developed for the stereocontrolled construction of the **A** and **C** *cis*-tetrahydropyran rings inscribed within the parent phorboxazole macrolide. Equally exciting was the potential for the design, synthesis, and biological evaluation of hemi-phorboxazole A analogues, a program that would fit nicely with our ongoing phorboxazole analogue venture.⁸

With this as background, we envisioned that (+)-hemi-phorboxazole A (**1**) could be readily elaborated by exploiting the same late-stage vinyl iodide macrolide (+)-**2**, which was constructed via a longest linear sequence of 20 steps (20% yield) exploiting our now scalable second-generation phorboxazole A synthesis.⁶ Indeed, removal of the TBS group at C(13) by treatment with tetrabutylammonium fluoride (TBAF), followed by a palladium-catalyzed cyanation,⁹ employing tributyltin cyanide, provided totally synthetic (+)-

hemi-phorboxazole A (**1**) in 85% yield for the two steps (Scheme 1). Importantly, the spectral data of synthetic (+)-**1** [cf. 1D and 2D NMR, UV, circular dichromism (CD), and HRMS] were identical in all respects with the data derived from natural hemi-phorboxazole A (**1**), thereby confirming both the complete relative stereochemistry and assigned absolute configuration.

Scheme 1



With synthetic (+)-hemi-phorboxazole A (**1**) in hand, we turned to the design, elaboration, and biological evaluation of analogues, with the important goal of introducing synthetic simplicity while maintaining the potentially important macrolide ring conformation of the natural congener. At the outset, we selected two analogues featuring macrolide ring replacements on the basis of our earlier analogue studies of (+)-phorboxazole A (**5**).⁸ In particular, we proposed: (i) replacement of the **C**-ring tetrahydropyran with a cyclic acetal and (ii) a bis-ring replacement comprising a similar acetal for the **C**-ring tetrahydropyran and a phenyl ring for the **B**-ring oxazole (Figure 2). Replacement of the tetrahy-




Figure 2. Phorboxazole analogues.

dropyran ring with a synthetically simplified but geometrically similar cyclic acetal (i.e., 1,3-dioxane) comprises a tactic that was introduced and elegantly employed by Wender et al. in their bryostatin analogue program.¹⁰ This concept was also utilized with some success in our (+)-

(5) Smith, A. B., III; Verhoest, P. R.; Minbirole, K. P.; Schelhaas, M. *J. Am. Chem. Soc.* **2001**, *123*, 4834.

(6) (a) Smith, A. B., III; Razler, T. M.; Ciavarri, J. P.; Hirose, T.; Ishikawa, T. *Org. Lett.* **2005**, *7*, 4399. (b) Smith, A. B., III; Razler, T. M.; Ciavarri, J. P.; Hirose, T.; Ishikawa, T.; Meis, R. M. *J. Org. Chem.* **2008**, *73*, 1192.


(7) Smith, A. B., III; Fox, R. J.; Razler, T. M. *Acc. Chem. Res.* **2008**, *41*, 675.

(8) (a) Smith, A. B., III; Razler, T. M.; Pettit, G. R.; Chapuis, J.-C. *Org. Lett.* **2005**, *7*, 4403. (b) Smith, A. B., III; Razler, T. M.; Meis, R. M.; Pettit, G. R. *Org. Lett.* **2006**, *8*, 797. (c) Smith, A. B., III; Razler, T. M.; Meis, R. M.; Pettit, G. R. *J. Org. Chem.* **2008**, *73*, 1201.

phorboxazole A (**5**) SAR study to furnish a novel, fully endowed phorboxazole analogue (**14**) possessing the macrolide ring, complete with an acetylenic side chain. Importantly, only moderate loss (ca. 10-fold) in tumor cell growth inhibition activity was observed when compared to (+)-phorboxazole A (**5**).^{8b}


Toward this end, the C-ring replacement congener of (+)-hemi-phorboxazole A (cf. **3**) was constructed via a palladium-catalyzed cyanation of vinyl iodide (+)-**15** (Scheme 2). Vinyl iodide (+)-**15** was prepared in accordance with our reported synthetic route to (+)-**14**.^{8b}

Scheme 2




The second analogue, hemi-phorboxazole congener **4**, possessing B- and C-ring replacements, was envisioned to be constructed in an analogous manner to that of (+)-**1**, exploiting two late-stage intermediates, vinyl iodide (+)-**17**^{8a} and diol (–)-**18**,^{8b} readily available from our second-generation phorboxazole A (**5**) total synthesis (Scheme 3). Construction of the key C(11–15) acetal would then result via union of diol (–)-**18** and aldehyde **16**, the latter prepared in two steps from commercially available 1,3-benzenedimethanol (see the Supporting Information).

Scheme 3



We began with the condensation of diol (–)-**18** and aldehyde **16** (Scheme 4), employing dehydrating conditions, to provide *cis*-acetal (–)-**19** in 94% yield (dr > 20:1). Removal of the PMB group at C(19) and exposure of the resultant primary alcohol to CCl₄ and PPh₃ furnished chloride (–)-**20** in 88% yield for the two steps.⁵ Next, conversion of

Scheme 4



chloride (–)-**20** to the corresponding tributyl phosphonium salt, followed by Wittig reaction with aldehyde (+)-**17**, provided the C(3–28) carbon skeleton of analogue **4**. The Wittig reaction, which employed a stabilized phosphonium ylide, proceeded in 94% yield with modest selectivity (ca. 4:1) favoring the *E*-isomer. Hydroxy aldehyde (–)-**22** was then obtained via a three-step sequence: removal of the silyl ether at C(3), Dess–Martin oxidation of the resultant primary alcohol to the corresponding aldehyde, and DDQ-mediated generation of the secondary hydroxyl at C(24). Macrocyclization of (–)-**22** to furnish the desired *Z*-macrolide (–)-**24** was then achieved employing a Still–Genari modified Horner–Emmons olefination¹¹ similar to that employed for the construction of (+)-**2**. Thus, generation of the phosphonate ester by reaction at the C(24) hydroxyl of (–)-**22** with 2-[bis(2,2,2-trifluoroethoxy)phosphoryl]acetic acid **23** and subsequent exposure to K₂CO₃/18-crown-6 in toluene afforded (–)-**24** in 44% overall yield for the two steps. The *E*-macrolide was also obtained in 15% yield. Finally, palladium-catalyzed cyanation at C(28) of vinyl iodide (–)-**24** provided hemi-phorboxazole congener (–)-**4** in 93% yield.

Synthetic (+)-hemi-phorboxazole A (**1**) and congeners (+)-**3** and (–)-**4** were assayed for tumor cell growth inhibitory activity, in parallel with (+)-phorboxazole A (**5**), against two human cancer cell lines (HCT-116, colon and

(9) For related transformations, see: (a) Duffey, M. O.; LeTiran, A.; Morken, J. P. *J. Am. Chem. Soc.* **2003**, *125*, 1458. (b) Nicolaou, K. C.; Pratt, B. A.; Arseniyadis, S.; Wartmann, M.; O'Brate, A.; Giannakakou, P. *ChemMedChem* **2006**, *1*, 41.

(10) Wender, P. A.; Baryza, J. L.; Bennett, C. E.; Bi, F. C.; Brenner, S. E.; Clarke, M. O.; Horan, J. C.; Kan, C.; Lacôte, E.; Lippa, B.; Nell, P. G.; Turner, T. M. *J. Am. Chem. Soc.* **2002**, *124*, 13648.

(11) Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, *24*, 4405.

Table 1. IC₅₀ Values for (+)-Hemi-phorboxazole A and Analogues against Human Cancer Cell Lines¹³

compd	HCT-116IC ₅₀ (ng/mL)	SKBR-3IC ₅₀ (ng/mL)
(+)-phorboxazole A (5)	0.71	2.0
(+)-hemi-phorboxazole A (1)	>6200	>6200
(+)- 3	>6200	>6200
(-)- 4	207	258

SK-BR-3, breast) (Table 1). Neither (+)-hemi-phorboxazole A (**1**) nor congener (+)-**3**, involving replacement of the C-ring tetrahydropyran with an acetal, displayed activity against either cell line. This result is perhaps not surprising given earlier reports that truncation of the phorboxazole side chain results in complete loss of potency.¹² Pleasingly, however, analogue (-)-**4**, which entailed replacement of two rings, displayed activity at the nanomolar level against both HCT-116 and SK-BR-3 cells.

Hemi-phorboxazole A (+)-**1** and analogue (+)-**3** were also assayed for antifungal activity against a panel of *C. albicans* strains. Although (+)-hemi-phorboxazole A (**1**) was again inactive, analogue (+)-**3** recorded a minimum inhibitory concentration (MIC) of 16 µg/mL (Table 2).

In summary, the first synthesis of (+)-hemi-phorboxazole A (**1**) has been achieved in two steps from our previously reported macrolide (+)-**2**, thereby confirming the assigned structure and complete stereochemistry. Two analogues of (+)-hemi-phorboxazole A, featuring macrolide ring replacements, were also designed and constructed. Biological evaluations of (+)-hemi-phorboxazole A (**1**) and analogues (+)-**3** and (-)-**4** against two human cancer cell lines (HCT-116, colon and SK-BR-3, breast) and *C. albicans* strains were conducted with the synthetic material. Although (+)-hemi-phorboxazole A (**1**) proved to be inactive in all cases, two analogues, (+)-**3** and (-)-**4**, respectively, displayed promising antifungal activity and significant anticancer activity against the two human cell lines. Although disappointed with

Table 2. Minimum Inhibitory Concentration (MIC, µg/mL) of (+)-Hemi-phorboxazole A and Analogue (+)-**3** against Pathogenic *C. albicans* Strains¹⁴

compd	<i>C. albicans</i> (µg/mL)		
	ATCC14503	UCD-FR1 ^a	96-489 ^a
(+)-phorboxazole A (5)	1.0	2.0	1.0
(+)-hemiphorboxazole A (1)	>64	>64	>64
(+)- 3	16.0	16.0	>64

^a Clinical isolate, fluconazole-resistant (MIC > 64 µg/mL).

our initial biological evaluation of (+)-hemi-phorboxazole A (**1**), the biological results revealed by analogues (+)-**3** and (-)-**4** warrant further investigation and thus will constitute part of our continuing phorboxazole synthetic program.

Acknowledgment. Support was provided by the National Institute of Health (National Cancer Institute) through grants CA-19033 (to A.B.S.) and CA-122256 (to T.F.M.) and the University of Pennsylvania. We thank Dr. Thomas M. Razler and Ms. Regina M. Meis (University of Pennsylvania) for some of the advanced intermediates prepared in connection with our phorboxazole synthetic program. We also thank Dr. Furst, Dr. Gu, and Dr. Kohli at the University of Pennsylvania for assistance in obtaining NMR and high-resolution mass spectra, respectively.

Supporting Information Available: Experimental procedures and full spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL9014317

(12) Uckan, F. M.; Forsyth, C. J. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1181.

(13) Cory, A. H.; Owen, T. C.; Barltrop, J. A.; Cory, J. G. *Cancer Commun.* **1991**, *7*, 207.

(14) National Committee for Clinical Laboratory Standards, 2002. Reference method for broth dilution antifungal susceptibility testing of yeast, 2nd ed. Approved standard M27-A2. National Committee for Clinical Laboratory Standards, Wayne, PA.