

Palladium-Catalyzed Desulfitative Cross-Coupling Reaction of Sodium Sulfinates with Benzyl Chlorides


Feng Zhao, Qi Tan, Fuhong Xiao, Shufeng Zhang, and Guo-Jun Deng*

Key Laboratory of Environmentally Friendly Chemistry and Application of Ministry of Education, College of Chemistry, Xiangtan University, Xiangtan 411105, China

gjdeng@xtu.edu.cn

Received February 1, 2013

ABSTRACT



A palladium-catalyzed approach for the synthesis of diarylmethanes from sodium sulfinates and benzyl chlorides is described. Various aromatic sodium sulfinates were used as aryl sources via extrusion of SO₂ and gave the diarylmethanes in moderate to good yields.

Diarylmethanes are versatile intermediates in the synthesis of pharmaceuticals and biologically active compounds,¹ and they are also employed as subunits in the design of supramolecular structures.² Consequently, development of efficient methods for construction of diarylmethanes has stimulated considerable interest. Traditionally, diarylmethanes are synthesized via Friedel–Crafts benzylation of benzylic electrophiles in the presence of Lewis acid.³ However, this approach is mainly suitable for electron-rich arenes and often suffers from low regioselectivity. The reduction of diaryl ketones could provide an alternative approach for preparation of diarylmethanes.⁴ Recently,

the transition-metal-catalyzed cross-coupling reactions of benzylic halides with arylmetals (or benzyldmetals with aryl halides)^{5,6} have been proven to be very powerful for construction of these compounds. Among them, the most promising transition-metal-catalyzed methods for diarylmethane synthesis have been the Suzuki–Miyaura-type coupling of benzylic electrophiles with arylboronic acids.^{7,8}

Metal-catalyzed C–H bond activation and subsequent C–C bond-forming reaction can provide an efficient synthesis with a reduced number of synthetic operations.⁹

(1) For selected examples, see: (a) McPhail, K. L.; Rivett, D. E. A.; Lack, D. E.; Davis-Coleman, M. T. *Tetrahedron* **2000**, *56*, 9391. (b) Graffner-Nordberg, M.; Kolmodin, K.; Aqvist, J.; Queener, S. F.; Hallberg, A. *J. Med. Chem.* **2001**, *44*, 2391. (c) Forsch, R. A.; Queener, S. F.; Rosowsky, A. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1811. (d) Howell, A.; Dowsett, M. *Breast Cancer Res.* **2004**, *6*, 269. (e) Bentley, K. W. *Nat. Prod. Rep.* **2005**, *22*, 249. (f) Mertins, K.; Iovel, I.; Kischel, J.; Zapf, A.; Beller, M. *Adv. Synth. Catal.* **2006**, *348*, 691.

(2) (a) Philp, D.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **1996**, *35*, 1154. (b) Ma, J. C.; Dougherty, D. A. *Chem. Rev.* **1997**, *97*, 1303. (c) Conn, M. M.; Rebek, J. *Chem. Rev.* **1997**, *97*, 1647. (d) Jasat, A.; Sherman, J. C. *Chem. Rev.* **1999**, *99*, 931.

(3) (a) Olah, G. A. *Friedel–Crafts and Related Reactions*; Wiley-Interscience: New York, 1964; Vol. II, Part 1. (b) Roberts, R. M.; Khalaf, A. A. *Friedel–Crafts Alkylation Chemistry. A Century of Discovery*; Marcel Dekker: New York, 1984. (c) Bandini, M.; Melloni, A.; Umani-Ronchi, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 550.

(4) (a) Olah, G. A.; Arvanaghi, M.; Ohannesian, L. *Synthesis* **1986**, 770. (b) Popielarz, R.; Arnold, D. R. *J. Am. Chem. Soc.* **1990**, *112*, 3068. (c) Esaki, H.; Aoki, F.; Umemura, M.; Kato, M.; Maegawa, T.; Mouguchi, Y.; Sajiki, H. *Chem.—Eur. J.* **2007**, *13*, 4052.

(5) (a) Liegault, B.; Renaud, J. L.; Bruneau, C. *Chem. Soc. Rev.* **2008**, *37*, 290. (b) Kuwano, R. *Synthesis* **2009**, *7*, 1049. (c) *Metal Catalyzed Cross-Coupling Reactions*; Diederich, F., Stang, P. J., Eds.; Wiley-VCH: Weinheim, 1998. (d) *Metal Catalyzed Cross-Coupling Reactions*, 2nd ed.; de Meijere, A., Diederich, F., Eds.; Wiley-VCH: Weinheim, 2004.

(6) For an example on selective diarylmethane formation using Grignard reagents with benzyl alcohols, see: Yu, D. G.; Wang, X.; Zhu, R. Y.; Luo, S.; Zhang, X. B.; Wang, B. Q.; Wang, L.; Shi, Z. J. *J. Am. Chem. Soc.* **2012**, *134*, 14638.

(7) For recent examples, see: (a) Singh, R.; Viciu, M. S.; Kramareva, N.; Navarro, O.; Nolan, S. P. *Org. Lett.* **2005**, *7*, 1829. (b) Molander, G. A.; Elia, M. D. *J. Org. Chem.* **2006**, *71*, 9198. (c) Burns, M. J.; Fairlamb, I. J. S.; Kapdi, A. R.; Sehna, P.; Taylor, R. J. K. *Org. Lett.* **2007**, *9*, 5397. (d) Srimani, D.; Sarkar, A. *Tetrahedron Lett.* **2008**, *49*, 6304. (e) Inés, B.; Moreno, I.; SanMartin, R.; Dominguez, E. *J. Org. Chem.* **2008**, *73*, 8448. (f) Inés, B.; SanMartin, R.; Moure, M.; Dominguez, E. *Adv. Synth. Catal.* **2009**, *351*, 2124. (g) Bedford, R. B.; Gower, N. J.; Haddow, M. F.; Harvey, J. N.; Nunn, J.; Okopie, R. A.; Sankey, R. F. *Angew. Chem., Int. Ed.* **2012**, *51*, 5435. (h) Yu, A. J.; Li, X. D.; Peng, D. P.; Wu, Y. J.; Chang, J. B. *Appl. Organometal. Chem.* **2012**, *26*, 301. (i) Maity, P.; Shacklady-McAtee, D. M.; Yap, G. P. A.; Sirianni, E. R.; Watson, M. P. *J. Am. Chem. Soc.* **2013**, *135*, 280.

(8) For a reaction using zinc in water, see: Duplais, C.; Krasovskiy, A.; Wattenberg, A.; Lipshutz, B. H. *Chem. Commun.* **2010**, *46*, 562.

Various methods have been developed using direct benzylation of heteroarenes and directing-group containing arenes under transition-metal catalysis.¹⁰ The Goossen group and others developed various methods for decarboxylative coupling reactions using inexpensive and readily available aromatic carboxylic acids as substrates.¹¹ The decarboxylative cross-coupling reactions with aryl halides have proved to be very effective and powerful for biaryl synthesis. However, this strategy requires *ortho*-functionalized benzoic substrates and is not suitable for cross-coupling reactions with benzyl halides due to the easy formation of very stable benzyl benzoate derivatives.¹² Meanwhile, we and others developed various palladium-catalyzed desulfative Heck-type reactions,¹³ addition reactions,¹⁴ homocoupling reactions,¹⁵ and cross-coupling

reactions with C–H bonds¹⁶ using sodium sulfonates as substrates. Unlike the decarboxylative coupling reactions, no electron-withdrawing or donating group *ortho* to the sulfinic acids group is necessary. We also found that the reaction of benzylic halides with sodium sulfonates generates stilbene derivatives in the absence of transition metals. The formation of benzyl sulfone intermediate plays an important role in this transformation.¹⁷ In continuing with our interest in using aryl sodium sulfonates as aryl sources, herein we describe a palladium-catalyzed desulfative cross-coupling reactions of sodium sulfonates with benzyl chlorides, affording diarylmethanes in moderate to good yields.¹⁸

(9) For recent reviews on C–H functionalization, see: (a) Dyker, G. *Handbook of C–H Transformations: Applications in Organic Synthesis*; Wiley-VCH: Weinheim, 2005. (b) Yu, J.; Shi, Z. *C–H Activation*; Springer: Berlin, 2010. (c) Goldberg, K. I.; Goldman, A. S. *Activation and Functionalization of C–H Bond*; ACS Symposium Series 885; American Chemical Society: Washington, DC, 2004. (d) Diaz-Requejo, M.; Pérez, P. *Chem. Rev.* **2008**, *108*, 3379. (e) Hartwig, J. *Nature* **2008**, *455*, 314. (f) Giri, R.; Shi, B.; Engle, K.; Maugel, N.; Yu, J. Q. *Chem. Soc. Rev.* **2009**, *38*, 3242. (g) Li, C. J. *Acc. Chem. Res.* **2009**, *42*, 335. (h) Copéret, C. *Chem. Rev.* **2010**, *110*, 656. (i) Mkhallid, I.; Barnard, J.; Marder, T.; Murphy, J.; Hartwig, J. *Chem. Rev.* **2010**, *110*, 890. (j) Colby, D. A.; Bergman, R. G.; Ellman, J. A. *Chem. Rev.* **2010**, *110*, 624. (k) Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147.

(10) (a) Arockiam, P. B.; Bruneau, C.; Dixneuf, P. H. *Chem. Rev.* **2012**, *112*, 5879. (b) Jaouhari, R.; Dixneuf, P. H. *Inorg. Chim. Acta* **1988**, *145*, 179. (c) Kuwano, R.; Kondo, Y.; Matsuyama, Y. *J. Am. Chem. Soc.* **2003**, *125*, 12104. (d) Mertins, K.; Iovel, I.; Kischel, J.; Zapf, A.; Beller, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 238. (e) Dong, C. G.; Hu, Q. S. *Angew. Chem., Int. Ed.* **2006**, *45*, 2289. (f) Ren, H. J.; Knochel, P. *Angew. Chem., Int. Ed.* **2006**, *45*, 3462. (g) Kuwano, R.; Shige, T. *J. Am. Chem. Soc.* **2007**, *129*, 3802. (h) Yu, J. Y.; Kuwano, R. *Org. Lett.* **2008**, *10*, 973. (i) Hwang, S. J.; Cho, S. H.; Chang, S. J. *Am. Chem. Soc.* **2008**, *130*, 16158. (j) Lapointe, D.; Fagnou, K. *Org. Lett.* **2009**, *11*, 4160. (k) Ackermann, L.; Novák, P. *Org. Lett.* **2009**, *11*, 4966. (l) Verrier, C.; Hoarau, C.; Marsair, F. *Org. Biomol. Chem.* **2009**, *7*, 647. (m) Ueno, S.; Ohtsubo, M.; Kuwano, R. *J. Am. Chem. Soc.* **2009**, *131*, 12904. (n) Ackermann, L.; Novák, P.; Vicente, R.; Hofmann, N. *Angew. Chem., Int. Ed.* **2009**, *48*, 6045. (o) Deng, G. J.; Li, C. J. *Org. Lett.* **2009**, *11*, 1171. (p) Mukai, T.; Hirno, K.; Satoh, T.; Miura, M. *Org. Lett.* **2010**, *12*, 1360. (q) Ackermann, L.; Füssler, S.; Pospech, J. *Org. Lett.* **2010**, *12*, 724. (r) Yi, C. S.; Lee, D. W. *Organometallics* **2010**, *29*, 1883. (s) Ackermann, L.; Hofmann, N.; Vicente, R. *Org. Lett.* **2011**, *13*, 1875.

(11) For recent reviews, see: (a) Goossen, L. J.; Rodríguez, N.; Goossen, K. *Angew. Chem., Int. Ed.* **2008**, *47*, 3100. (b) Rodríguez, N.; Goossen, L. J. *Chem. Soc. Rev.* **2011**, *40*, 5030. For selected examples on decarboxylative cross-coupling reactions, see: (c) Goossen, L. J.; Deng, G. J.; Levy, L. M. *Science* **2006**, *313*, 662. (d) Goossen, L. J.; Rodríguez, N.; Bettina, M.; Linder, C.; Deng, G. J.; Levy, L. M. *J. Am. Chem. Soc.* **2007**, *129*, 4824. (e) Forgione, P.; Brochu, M.; St-Onge, M.; Thesen, K.; Bailey, M.; Bilodeau, F. *J. Am. Chem. Soc.* **2006**, *128*, 11350.

(12) Otera, J.; Nishikido, J. *Esterification: Methods, Reactions and Applications*; Wiley-VCH: Weinheim, 2010.


(13) (a) Zhou, X. Y.; Luo, J. Y.; Liu, J.; Peng, S. M.; Deng, G. J. *Org. Lett.* **2011**, *13*, 1432. (b) Wang, G.; Miao, T. *Chem.—Eur. J.* **2011**, *17*, 5787.

(14) (a) Liu, J.; Zhou, X.; Rao, H.; Xiao, F.; Li, C. J.; Deng, G. J. *Chem.—Eur. J.* **2011**, *17*, 7996. (b) Yao, H.; Yang, L.; Shuai, Q.; Li, C. J. *Adv. Synth. Catal.* **2011**, *353*, 1701. (c) Miao, T.; Wang, G. W. *Chem. Commun.* **2011**, *47*, 9501. (d) Behrends, M.; Sävmarker, J.; Sjöberg, P.; Larhed, M. *ACS Catal.* **2011**, *1*, 1455. (e) Wang, H.; Li, Y.; Zhang, R.; Jin, K.; Zhao, D.; Duan, C. Y. *J. Org. Chem.* **2012**, *77*, 4849. (f) Chen, W.; Zhou, X.; Xiao, F.; Luo, J.; Deng, G. J. *Tetrahedron Lett.* **2012**, *53*, 4347.

(15) Rao, B.; Zhang, W.; Hu, L.; Luo, M. M. *Green Chem.* **2012**, *14*, 3436.

(16) (a) Chen, R.; Liu, S.; Liu, X.; Yang, L.; Deng, G. J. *Org. Biomol. Chem.* **2011**, *9*, 7675. (b) Wu, M.; Luo, J.; Xiao, F.; Zhang, S.; Deng, G. J.; Luo, H. A. *Adv. Synth. Catal.* **2012**, *354*, 335. (c) Liu, B.; Guo, Q.; Cheng, Y.; Lan, J.; You, J. S. *Chem.—Eur. J.* **2011**, *17*, 13415. (d) Wang, M.; Li, D.; Zhou, W.; Wang, L. *Tetrahedron* **2012**, *68*, 1926.

Table 1. Optimization of the Reaction Conditions^a

					
entry	catalyst	ligand	base	solvent	yield ^b (%)
1	Pd(OAc) ₂	PPh ₃	Na ₂ CO ₃	dioxane	45
2	PdCl ₂	PPh ₃	Na ₂ CO ₃	dioxane	40
3	PdBr ₂	PPh ₃	Na ₂ CO ₃	dioxane	24
4	Pd(OH) ₂	PPh ₃	Na ₂ CO ₃	dioxane	trace
5	Pd(acac) ₂	PPh ₃	Na ₂ CO ₃	dioxane	25
6	Pd(COD)Cl ₂	PPh ₃	Na ₂ CO ₃	dioxane	44
7	Pd(TFA) ₂	PPh ₃	Na ₂ CO ₃	dioxane	20
8	Pd(OAc) ₂	dppe	Na ₂ CO ₃	dioxane	50
9	Pd(OAc) ₂	DPEphos	Na ₂ CO ₃	dioxane	21
10	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	Na ₂ CO ₃	dioxane	58
11	Pd(OAc) ₂	<i>o</i> -Tol ₃ P	Na ₂ CO ₃	dioxane	trace
12	Pd(OAc) ₂	<i>m</i> -Tol ₃ P	Na ₂ CO ₃	dioxane	57
13	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	NaHCO ₃	dioxane	60
14	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	<i>t</i> -BuOK	dioxane	62
15	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	dioxane	65
16	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	anisole	66
17	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	toluene	64
18	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	DMSO	trace
19	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	cyclohexane	72
20 ^c	Pd(OAc) ₂	<i>p</i> -Tol ₃ P	CH ₃ ONa	cyclohexane	88

^a Conditions: **1a** (0.2 mmol), **2a** (0.2 mmol), catalyst (5 mol %), ligand (10 mol %), base (0.3 mmol), solvent (0.6 mL), 120 °C, 4 h under air unless otherwise noted. ^b GC yield. ^c 160 °C.

We began our study by examining the reaction of benzyl chloride (**1a**) with *p*-toluenesulfinic acid sodium salt (**2a**) in dioxane by using Pd(OAc)₂/PPh₃ as catalyst and Na₂CO₃ as base. When benzyl chloride reacted with an equal amount of **2a** under air, the desired product was obtained in 45% yield as detected by GC–MS and ¹H NMR methods (Table 1, entry 1). Then various palladium salts were investigated for this reaction under similar reaction

(17) Zhao, F.; Luo, J. Y.; Tan, Q.; Liao, Y. F.; Peng, S. M.; Deng, G. J. *Adv. Synth. Catal.* **2012**, *354*, 1914.

(18) Diarylmethanes can be formed using sulfonyl chlorides with organometals: (a) Dubbaka, S. R.; Vogel, P. *J. Am. Chem. Soc.* **2003**, *125*, 15292. (b) Dubbaka, S. R.; Vogel, P. *Org. Lett.* **2004**, *6*, 95. (c) Dubbaka, S.; Vogel, P. *Angew. Chem., Int. Ed.* **2005**, *44*, 7674. (d) Rao Volla, C.; Vogel, P. *Angew. Chem., Int. Ed.* **2008**, *47*, 1305.

conditions. Similar yields were observed when PdCl₂ and Pd(COD)Cl₂ were used as the catalysts (entries 2 and 6). Other palladium salts such as PdBr₂, Pd(OH)₂, Pd(acac)₂, and Pd(TFA)₂ were less effective (entries 3–5 and 7). With Pd(OAc)₂ as the catalyst, the effect of ligand on the reaction yield was investigated by screening various phosphine ligands. Among them, *p*-Tol₃P (tri-*p*-toluenephosphine) showed the best reactivity, and the desired product was obtained in 58% yield (entry 10), whereas the use of *o*-Tol₃P completely inhibited the reaction (entry 11). Base also affected the reaction yields slightly. The use of CH₃ONa can further improve the reaction yield to 65% (entry 15). The choice of solvents was crucial for this reaction. The reaction in DMSO did not give any desired product (entry 18). Other solvents such as anisole, toluene, and cyclohexane were proved to be also good reaction media (entries 16, 17 and 19). When the reaction temperature increased to 160 °C, the reaction yield was improved to 88% using cyclohexane as solvent (entry 20).

Table 2. Reaction of **2a** with Various Benzyl Chlorides^a

entry	benzyl chloride		product	yield (%) ^b
1		1a		80
2		1b		72
3		1c		72
4		1d		70
5		1h		50
6		1e		63
7		1f		66
8		1g		51

^a Conditions: **1** (0.2 mmol), **2a** (0.2 mmol), Pd(OAc)₂ (5 mol %), *p*-Tol₃P (10 mol %), CH₃ONa (0.3 mmol), cyclohexane (0.6 mL), 160 °C, 4 h under air. ^b Isolated yield.

With the optimized reaction conditions established, the scope of the reaction with respect to *p*-toluenesulfinic acid sodium salt (**2a**) and various benzyl chlorides (**1**) was investigated (Table 2). The reactions with benzyl chlorides bearing electron-donating groups (entries 2–5) and electron-withdrawing substituents (entries 6–8) at the aromatic ring

proceeded smoothly to give the desired products in moderate to good yields. The position of the substituents did not affect the reaction yields, and *o*-, *m*-, and *p*-methylbenzyl chlorides all gave the desired products in good yields (entries 2–4). However, the use of 4-methoxybenzyl chloride and 4-nitrobenzyl chloride both decreased the reaction yields significantly. Notably, replacement of benzyl chloride with benzyl bromide dramatically decreased the reaction yield under similar reaction conditions partly due to the much higher reactivity of benzyl bromide.


Table 3. Reaction of **1d** with Various Sodium Sulfonates (**2**)^a

entry	sodium sulfonate		product	yield (%) ^b
1		2b		70
2		2c		72
3		2d		71
4		2e		73
5		2f		52
6		2g		72
7		2h		66
8		2i		81
9		2j		trace

^a Conditions: **1d** (0.2 mmol), **2** (0.2 mmol), Pd(OAc)₂ (5 mol %), *p*-Tol₃P (10 mol %), CH₃ONa (0.3 mmol), cyclohexane (0.6 mL), 160 °C, 4 h under air. ^b Isolated yield.

Under the optimized reaction conditions, the substituent effect on the reaction yield was investigated and the results are presented in Table 3. In general, alkyl substituents on the para position of sulfinic acid group did not affect the reaction yield significantly (entries 2–4). However, a methoxy substituent profoundly decreased the reaction yield, and the corresponding product **3m** was obtained in 52% yield (entry 5). Halogen substituents such as fluoro, chloro, and trifluoromethyl were tolerated under the

Scheme 1. Control Experiments



optimal reaction conditions, and the desired products were obtained in moderate to good yields (entries 6–8). An electron-withdrawing group such as nitro on the aromatic ring of sulfinic acid dramatically decreased the reaction yield, and only a trace amount of product was observed when (4-nitrophenyl)sulfinic acid sodium salt (**2j**) reacted with **1d** (entry 9). In all cases, cross-coupled diarylmethanes were the major products as determined by ^1H NMR method.¹⁹


To get more information about the reaction mechanism, several control experiments were set up under the standard conditions. When the reaction of **1a** with **2a** was set up at 60°C under air, 1-(benzylsulfonyl)-4-methylbenzene (**4a**) was obtained in 83% yield, and no corresponding ester (**5a**) was observed (scheme 1).¹⁷ Treatment of **4a** under the standard reaction conditions did not afford the final product **3a**. This means the product was not generated from **4a** or **5a** via extrusion of SO_2 . Based on these observations, a tentative mechanism to rationalize the transformation is illustrated in Scheme 2. Oxidative addition of $\text{Pd}(0)$ species with **1a** generates an intermediate Bn-Pd(II)-Cl (**A**).^{20,21,5a} The replacement of chloride with arylsulfinic acid sodium salt forms an intermediate **B**, which can be further converted

(19) The use of *o*-toluenesulfinic acid sodium salt significantly decreased the reaction yield to 20% as determined by GC.

(20) For an early example on insertion of $\text{Pd}(0)$ into a benzylic carbon–halogen bond, see: Fitton, P.; McKeon, J. E.; Ream, B. C. *J. Chem. Soc. D*. **1969**, 370.

(21) For an example of palladium-catalyzed cross-coupling of benzyl chlorides with olefins involving radical mechanism, see: Pan, Y.; Zhang, Z. Y.; Hu, H. W. *Synth. Commun.* **1992**, 22, 2019.

Scheme 2. Proposed Mechanism



into intermediate **C** via extrusion of SO_2 . A reductive elimination of **C** affords the final product and regenerates $\text{Pd}(0)$ species, thus closing the catalytic cycle.

In conclusion, we have demonstrated a palladium-catalyzed approach for the synthesis of diarylmethanes via desulfative benzylation of aromatic sulfinic acid sodium salts with benzyl chlorides. The reaction showed good selectivity and tolerated various functional groups. This method provides an alternative route for the synthesis of diarylmethanes from benzyl chlorides. The scope, mechanism, and synthetic applications of this reaction are being explored in our laboratory.

Acknowledgment. This work was supported by the National Natural Science Foundation of China (21172185), the Hunan Provincial Natural Science Foundation of China (11JJ1003, 12JJ7002), the New Century Excellent Talents in University from Ministry of Education of China (NCET-11-0974), and the Hunan Provincial Innovative Foundation for Postgraduate (CX2012B250) and Undergraduate.

Supporting Information Available. General experimental procedure and characterization data of the products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.