

Metal-Catalyzed Cascade Rearrangements of 3-Alkynyl Flavone Ethers


Yuan Xiong, Scott E. Schaus, and John A. Porco, Jr.*

Center for Chemical Methodology and Library Development, Boston University,
590 Commonwealth Avenue, Boston, Massachusetts 02215, United States

porco@bu.edu

Received March 9, 2013

ABSTRACT



Metal-mediated rearrangements of 3-alkynyl flavone ethers are reported. The overall process involves 5-*endo* enyne cyclization to a platinum-containing spiro-oxocarbenium intermediate which may be trapped with methanol to produce spirodihydrofurans or further rearranged to afford either allenyl chromanediones or benzofuranones.

As part of our interest in the chemical reactivity of 3-flavone ethers, we have reported the scandium triflate catalyzed Claisen rearrangement of 3-allyloxyflavone ethers such as **1** to access 3,4-chromanediones **2**.¹ In the current study, we investigated the possibility for metal-catalyzed alkynyl-Claisen (Saucy–Marbet) rearrangements of the corresponding 3-alkynyl flavone ethers **3** to access allenyl-chromanediones **4**. Use of transition metal catalysts such as Au(I),² Ag(I),³ Pd(II),⁴ Rh(I),⁵

(1) Marie, J.-C.; Xiong, Y.; Min, G. K.; Yeager, A. R.; Taniguchi, T.; Berova, N.; Schaus, S. E.; Porco, J. A., Jr. *J. Org. Chem.* **2010**, *75*, 4584–4590.

(2) (a) Sherry, B. D.; Toste, F. D. *J. Am. Chem. Soc.* **2004**, *126*, 15978–15979. (b) Suhre, M. H.; Reif, M.; Kirsch, S. F. *Org. Lett.* **2005**, *7*, 3925–3927. (c) Sherry, B. D.; Maus, L.; Laforteza, B. N.; Toste, F. D. *J. Am. Chem. Soc.* **2006**, *128*, 8132–8133. (d) Mauleón, P.; Krinsky, J. L.; Toste, F. D. *J. Am. Chem. Soc.* **2009**, *131*, 4513–4520.

(3) (a) Grissom, J. W.; Klingberg, D.; Huang, D.; Slattery, B. J. *J. Org. Chem.* **1997**, *62*, 603–626. (b) Binder, J. T.; Kirsch, S. F. *Org. Lett.* **2006**, *8*, 2151–2153. (c) Menz, H.; Kirsch, S. F. *Org. Lett.* **2006**, *8*, 4795–4797. (d) Harschneck, T.; Kirsch, S. F. *J. Org. Chem.* **2011**, *76*, 2145–2156.

(4) (a) Nevado, C.; Echavarren, A. M. *Tetrahedron* **2004**, *60*, 9735–9744. (b) Huang, H.; Jiang, H.; Cao, H.; Zhao, J.; Shi, D. *Tetrahedron* **2012**, *68*, 3135–3144.

(5) (a) Saito, A.; Kanno, A.; Hanzawa, Y. *Angew. Chem., Int. Ed.* **2007**, *46*, 3931–3933. (b) Okazaki, E.; Okamoto, R.; Shibata, Y.; Noguchi, K.; Tanaka, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 6722–6727.

(6) Cao, H.; Jiang, H.; Yao, W.; Liu, X. *Org. Lett.* **2009**, *11*, 1931–1933.





Figure 1. Metal-catalyzed rearrangement of 3-flavone ethers.

Cu(I),⁶ and Fe(III)⁷ for alkynyl-Claisen rearrangements has been reported. Subsequent one-pot cascade reactions to access heterocycles including furans, pyrroles, 2*H*-pyrans, and dihydropyridines have also been reported by the Kirsch^{2b,3b–3d} and Jiang laboratories.^{4b,6,7} Moreover, Kozłowski and co-workers have reported the first asymmetric Saucy–Marbet rearrangement of propargyl-substituted indoles to afford allenyl oxindoles or spirolactones

(7) Jiang, H.; Yao, W.; Cao, H.; Huang, H.; Cao, D. *J. Org. Chem.* **2010**, *75*, 5347–5350.

Table 1. Metal Catalyst Screening


entry	metal	t (°C)	time (h)	yield		
				7	8	5
1	AuCl ₃	60	36	53%	14%	33%
2	NaAuCl ₄ ·2H ₂ O	60	36	50%	trace	—
3	NaAuCl ₄ ·2H ₂ O	80	60	74%	trace	—
4	PtCl ₂	60	36	21%	—	60%
5	PtCl ₄	60	36	40%	28%	24%
6	PtCl ₂	80	60	—	53% ^a	—
7	PtCl ₄	80	60	—	44% ^a	—
8	Rh ₂ (tfa) ₄	80	36	—	24% ^a	—

^a The remaining mass balance is the cleavage product 3-hydroxyflavone.

catalyzed by Pd(II)/BINAP.⁸ In this Communication, we report metal-catalyzed rearrangements of 3-alkynyl flavone ethers which led to the identification of an unexpected cascade rearrangement pathway.



We began our evaluation of the alkynyl Claisen rearrangement of 3-alkynyl flavone ether **5** with a series of alkynophilic metals (Table 1). Among a panel of catalysts screened, AuCl₃ was found to catalyze the reaction to produce allenyl 3,4-chromanedione **6** which was condensed with 1,2-phenylenediamine¹ to provide chromenoquinoxaline **7**. Aurate salts proved to be efficient catalysts; in particular, 5 mol % of NaAuCl₄·2H₂O at elevated temperatures cleanly catalyzed the rearrangement of substrate **5** which was followed by diamine condensation to afford **7** in 74% yield (entry 3). In addition to the allene **7**, a small amount of furanyl benzofuranone **8** (*vide infra*) was also observed in the AuCl₃-catalyzed reaction (entry 1). The structure of **8** was confirmed by X-ray crystal structure analysis⁹ (Table 1). Use of platinum catalysis also led to the formation of **8**. At 60 °C, PtCl₄ afforded both **7** (40%) and **8** (28%) (entry 5).¹⁰ At a higher reaction temperature, PtCl₂ was highly efficient in producing benzofuranone **8**.

(8) (a) Cao, T.; Deitch, J.; Linton, E. C.; Kozlowski, M. C. *Angew. Chem., Int. Ed.* **2012**, *51*, 2448–2451. (b) Cao, T.; Linton, E. C.; Deitch, J.; Bertritt, S.; Kozlowski, M. C. *J. Org. Chem.* **2012**, *77*, 11034–11055.

(9) See Supporting Information for complete experimental details.

(10) For PtCl₂- and PtCl₄-catalyzed hydroarylation of alkynes, see: Pastine, S. J.; Youn, S. W.; Sames, D. *Org. Lett.* **2003**, *5*, 1055–1058.

(11) Gainer, M. J.; Bennett, N. R.; Takahashi, Y.; Looper, R. E. *Angew. Chem., Int. Ed.* **2011**, *50*, 684–687.

Scheme 1. Crossover Experiments**Scheme 2.** Intermolecular Trapping of an Oxocarbenium Intermediate

Rh₂(tfa)₄¹¹ also promoted the rearrangement to afford **8** in 24% yield along with a significant amount of propargyl ether cleavage to yield 3-hydroxyflavone as a side product.⁹ Rare earth triflate catalysts (Sc(OTf)₃ and La(OTf)₃) which were found to be workable for 3-allyloxyflavone ethers¹ were not effective for substrates such as **5**.⁹


In order to probe the reaction mechanism, crossover experiments were conducted to exclude the possibility of an intermolecular reaction pathway (Scheme 1). In the event, rearrangement of an equimolar mixture of butynyl ether **5** and pentynyl ether **9** in the presence of metal catalysts (PtCl₂, PtCl₄, or NaAuCl₄, 60 °C) resulted in the sole production of chromenoquinoxalines **7** and **10** after condensation with 1,2-phenylenediamine. The absence of crossover products by mass spectrometric⁹ analysis of the crude reaction mixture strongly suggests an intramolecular pathway as opposed to an ion pair mechanism.^{1,12}

Further mechanistic insights were gained by treatment of substrate **5** with Au(III) or Pt catalysts in the presence of methanol which afforded spirodihydrofuran **11**⁹ as a single diastereomer (36–58% yield, Scheme 2). The formation

(12) For a representative crossover study, see: Burger, E. C.; Tunge, J. A. *Org. Lett.* **2004**, *6*, 2603–2605.

(13) For transition-metal-catalyzed 5-endo cyclization of alkynes and 1,5-enynes, see: (a) Staben, S. T.; Kennedy-Smith, J. J.; Toste, F. D. *Angew. Chem., Int. Ed.* **2004**, *43*, 5350–5352. (b) Zhang, L.; Kozmin, S. A. *J. Am. Chem. Soc.* **2005**, *127*, 6962–6963. (c) Staben, S. T.; Kennedy-Smith, J. J.; Huang, D.; Corkey, B. K.; LaLonde, R. L.; Toste, F. D. *Angew. Chem., Int. Ed.* **2006**, *45*, 5991–5994. (d) Buzas, A. K.; Istrate, F. M.; Gagosz, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 1141–1144. (e) Brazeau, J.-F.; Zhang, S.; Colomer, I.; Corkey, B. K.; Toste, F. D. *J. Am. Chem. Soc.* **2012**, *134*, 2742–2749.

Scheme 3. Intramolecular Trapping of an Oxocarbenium Intermediate




of product **11** indicates that the metal catalysts preferentially activate the alkyne of flavone ether substrates such as **5** toward 5-*endo*¹³ and not 6-*endo* enyne cyclization. In the case of PtCl_4 additional bidentate coordination of the metal catalyst to the C-3 flavone ether and C-4 carbonyl oxygens¹⁴ may contribute to lower nucleophilicity at C-2, thereby rendering attack at C-3 more preferable to produce platinum-containing oxocarbenium intermediate **A**¹⁵ which afforded spirodihydrofuran **11** after reaction with methanol. The observed diastereoselectivity may be explained by steric shielding of the β face of the oxonium intermediate by the vinyl methyl group, leading to diastereoselective addition of methoxide to the α face.⁹


We also evaluated a 3-alkynyl flavone ether substrate bearing an internal nucleophile (Scheme 3). Treatment of 2'-hydroxyflavone ether **12** with PtCl_2 afforded allene **13** (45%) along with a trace amount of the bridged dihydropyran product **14**, the structure of which was confirmed by X-ray crystal structure analysis.⁹ Both **13** and **14** are presumably derived from intramolecular trapping of oxocarbenium **B** (*vide infra*) by the adjacent 2'-hydroxyl group. Use of PtCl_4 and $\text{NaAuCl}_4 \cdot \text{H}_2\text{O}$ as catalysts favored the formation of **14**. Eventually, we found that addition of 1,1,1,2,2,2-hexafluoroisopropanol (HFIP) as the proton source¹⁶ provided tetracycle **14** in 44% yield, conceivably by promoting protodemetalation (Scheme 3, pathway **b**) versus elimination (pathway **a**).

Additional mechanistic information was also obtained from allenyl chromanedione substrate **15**, derived from alkynyl flavone ether **16** using Au(III)-catalyzed rearrangement. When **15** was resubjected to rearrangement conditions at 80 °C, benzofuranone **17** was isolated in 60–80% yield (Scheme 4), suggesting that allenyl chromanedione substrates may also participate in the metal-

Scheme 4. Preparation and Rearrangement of an Allenyl Chromanedione



Scheme 5. Proposed Mechanistic Pathway



catalyzed cycloisomerization cascade to produce benzofuranone products (Scheme 5).¹⁷

Based on our experimental results, we propose the overall mechanistic pathway shown in Scheme 5. Following initial 5-*endo* enyne cyclization¹⁸ of alkynyl flavone ether **18**, the resulting platinum-containing spiro-oxocarbenium intermediate **C** may undergo pinacol-type 1,2-rearrangement¹⁹ to form dipole **D**,²⁰ possibly via a cyclopropyl platinacarbene intermediate **E**.²¹ Dipole **D** may further fragment to yield allenyl chromanedione **19** through platinum dissociation

(18) For reviews on metal-catalyzed enyne cycloisomerization, see: (a) Zhang, L.; Sun, J.; Kozmin, S. A. *Adv. Synth. Catal.* **2006**, *348*, 2271–2296. (b) Nieto-Oberhuber, C.; López, S.; Jiménez-Núñez, E.; Echavarren, A. M. *Chem.—Eur. J.* **2006**, *12*, 5916–5923. (c) Shen, H. C. *Tetrahedron* **2008**, *64*, 7847–7870. (d) Jimenez-Nunez, E.; Echavarren, A. M. *Chem. Rev.* **2008**, *108*, 3326–3350. (e) Michelet, V.; Toullec, P. Y.; Genêt, J.-P. *Angew. Chem., Int. Ed.* **2008**, *47*, 4268–4315.

(19) For examples of metal-catalyzed cycloisomerization involving pinacol-type rearrangement, see: (a) Kirsch, S. F.; Binder, J. T.; Crone, B.; Duschek, A.; Haug, T. T.; Liébert, C.; Menz, H. *Angew. Chem., Int. Ed.* **2007**, *46*, 2310–2313. (b) Huang, X.; Zhang, L. *J. Am. Chem. Soc.* **2007**, *129*, 6398–6399. (c) Tang, J.; Bhunia, S.; Soheli, S. M. A.; Lin, M.; Liao, H.; Datta, S.; Das, A.; Liu, R. *J. Am. Chem. Soc.* **2007**, *129*, 15677–15683. (d) Crone, B.; Kirsch, S. F. *Chem.—Eur. J.* **2008**, *14*, 3514–3522. (e) Shu, X.-Z.; Liu, X.-Y.; Ji, K. G.; Xiao, H.-Q.; Liang, Y.-M. *Chem.—Eur. J.* **2008**, *14*, 5282–5289.

(20) Kusama, H.; Suzuki, Y.; Takaya, J.; Iwasawa, N. *Org. Lett.* **2006**, *8*, 895–897.

(21) Martín-Matute, B.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. *J. Am. Chem. Soc.* **2003**, *125*, 5757–5766.

(14) Tang, H.; Wang, X.; Yang, S.; Wang, L. *Rare Met.* **2004**, *23*, 38–42.

(15) Li, G.; Huang, X.; Zhang, L. *J. Am. Chem. Soc.* **2008**, *130*, 6944–6945.

(16) Roth, K. E.; Blum, S. A. *Organometallics* **2010**, *29*, 1712–1716.

(17) For platinum- and gold-catalyzed 6-*endo* cycloisomerization of allenes, see: (a) Gockel, B.; Krause, N. *Org. Lett.* **2006**, *8*, 4485–4488.

(b) Kong, W.; Fu, C.; Ma, S. *Chem. Commun.* **2009**, *0*, 4572–4574.

Table 2. Substrate Scope

entry	substrate	product	yield%
1			PtCl ₂ , 20% PtCl ₄ , 10%
2			PtCl ₂ , 43% PtCl ₄ , 31%
3			PtCl ₂ , 31% PtCl ₄ , 43%
4			PtCl ₂ , 70% PtCl ₄ , 65%
5			PtCl ₂ , 80% PtCl ₄ , 64% (12 h)
6			PtCl ₂ , 18% PtCl ₄ , 12% (12 h)
7			PtCl ₂ 34: NR 36: 53% (48 h) 17: 67% (24 h)
	33 R = 2'-CF ₃ 35 R = 2'-MeO 16 R = 2',4'-MeO	34 R = 2'-CF ₃ 36 R = 2'-MeO 17 R = 2',4'-MeO	

^aProducts obtained after condensation with 1.5 equiv of 1,2-phenylenediamine.

(Scheme 5, blue arrow). At elevated temperatures, intermediate **C** may rearrange *via* a formal 1,2-acyl migration to obtain oxocarbenium **F**²² which after protodemetalation and aromatization may afford **20** as the thermodynamic product (Scheme 5, red arrow). Conceivably, allenyl chromanediene **19** may also undergo metal catalyzed 6-*endo* cycloisomerization¹⁷ to yield oxocarbenium intermediates **D**

(22) For formal 1,2-acyl migration in a Pt(II)-catalyzed cycloisomerization, see: (a) Li, G.; Huang, X.; Zhang, L. *Angew. Chem., Int. Ed.* **2008**, *47*, 346–349. For acid-catalyzed 1,2-acyl migration, see: (b) Bach, R. D.; Domagala, J. M. *J. Org. Chem.* **1984**, *49*, 4181–4188. (c) Bach, R. D.; Klix, R. C. *J. Org. Chem.* **1985**, *50*, 5438–5440. (d) Klix, R. C.; Bach, R. D. *J. Org. Chem.* **1987**, *52*, 580–586. (e) Crane, S. N.; Burnell, D. J. *J. Org. Chem.* **1998**, *63*, 5708–5710.

and **C**, which following similar pathways may produce benzofuranone **20** (Scheme 5, red arrow).

Regarding substrate scope, substitutions at the terminal position of the alkyne such as hydrogen as well as alkyl and phenyl groups (Table 2, **21**, **23**, **25**, entries 1–3) were tolerated, although substrates with terminal alkynes suffered low yields from side reactions presumably due to metal vinylidene formation. Flavone ether **27** derived from a secondary propargylic alcohol was also reactive affording the 2,3,5-trisubstituted furanyl benzofuranone **28** in high yield (entry 4). 5-Methoxy substrate **29**, with increased electron density on the flavone A-ring, underwent smooth acyl migration to afford benzofuranone **30** in high yield (entry 5). In contrast, electron-withdrawing groups found in substrates such as **31** disfavored acyl migration. In this instance, only allenyl chromenoquinoxaline **32** was isolated following diamine condensation even at an elevated reaction temperature (entry 6). Electron-rich C-rings found in substrates **16** and **35** favored the cascade process producing benzofuranones by increasing nucleophilicity at C-3 and stabilizing the forming positive charge in the proposed spiro-oxocarbenium intermediate **C** (entry 7, *cf.* Scheme 5). In contrast, the electron-poor 2'-trifluoromethyl flavone ether **33** was not a useable substrate and did not afford appreciable amounts of benzofuranone **34** (entry 7).

In summary, evaluation of the metal-catalyzed alkynyl Claisen (Saucy–Marbet) rearrangements of 3-alkynyl flavone ethers led to the discovery of a 1,2-acyl migration cascade which afforded novel furanyl benzofuranones. Mechanistic studies have revealed that the rearrangement is likely initiated by 5-*endo* enyne cyclization to a platinum-containing spiro-oxocarbenium intermediate, which may be intercepted by methanol to produce spirodihydrofurans or further rearranged to afford allenyl chromanediens at 60 °C and benzofuranones at elevated reaction temperatures. Further studies including asymmetric reaction development and diverse nucleophilic trapping of spiro-oxocarbenium intermediates are currently in progress and will be reported in future publications.

Acknowledgment. Financial support from the National Institutes of Health (P50 GM067041 and CA 099920) is gratefully acknowledged. We thank Prof. John Snyder and Drs. Alexander Grenning and Munmun Mukerjee (Boston University) for helpful discussions and Dr. Jeffrey Bacon (Boston University) for X-ray crystal structure analyses.

Supporting Information Available. Experimental procedure and characterization of compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.