

Et₃B-Mediated Radical-Polar Crossover Reaction for Single-Step Coupling of O, Te-Acetal, α,β -Unsaturated Ketones, and Aldehydes/Ketones


Daigo Kamimura, Daisuke Urabe, Masanori Nagatomo, and Masayuki Inoue*

Graduate School of Pharmaceutical Sciences, The University of Tokyo, Hongo,
Bunkyo-ku, Tokyo 113-0033, Japan

inoue@mol.f.u-tokyo.ac.jp

Received September 5, 2013

ABSTRACT



Et₃B-mediated three-component coupling reactions between O,Te-acetal, α,β -unsaturated ketones, and aldehydes/ketones were developed. Et₃B promoted the generation of the potently reactive bridgehead radical from the O,Te-acetal of the trioxadadamantane structure and converted the α -carbonyl radical of the resultant two-component adduct to the boron enolate, which then underwent a stereoselective aldol reaction with the aldehyde/ketone. This powerful, yet mild, radical-polar crossover reaction efficiently connected the hindered linkages between the three units and selectively introduced three new stereocenters.

One-pot, three-component coupling is a powerful and rapid means for construction of functionalized molecules¹ because it minimizes the number of synthetic operations and maximizes the build up of structural and functional complexity. While these reactions have historically employed either ionic or radical processes, the in situ combination of

radical and ionic reactions, designated as radical-polar crossover reactions, has recently emerged.^{2,3} Since the reactivities of the radical and ionic intermediates are orthogonal, sequential application of these two mechanisms offers great advantages in the assembly of complex molecular architectures. Despite these synthetically valuable features, these reactions have been unexplored in the context of target-oriented synthesis.

Multiply fused and highly oxygenated structures of terpenoids generally present a daunting synthetic challenge (Scheme 1).⁴ We have been interested in developing new three-component coupling reactions for construction of the substructures of bioactive terpenoids, such as cytotoxic trigonellin A⁵ and potassium channel blocker correolide.⁶ These compounds share a common motif, in which the

(1) For recent reviews on the multicomponent reactions, see: (a) *Multicomponent Reactions*; Zhu, J., Bienaymé, H., Eds.; Wiley-VCH: Weinheim, 2005. (b) Touré, B. B.; Hall, D. G. *Chem. Rev.* **2009**, *109*, 4439. (c) de Graaff, C.; Ruijter, E.; Orru, R. V. A. *Chem. Soc. Rev.* **2012**, *41*, 3969. (d) Marson, C. M. *Chem. Soc. Rev.* **2012**, *41*, 7712. (e) Pellissier, H. *Chem. Rev.* **2013**, *113*, 442.

(2) For representative examples, see: (a) Shono, T.; Nishiguchi, I.; Sasaki, M. *J. Am. Chem. Soc.* **1978**, *100*, 4314. (b) Nozaki, K.; Oshima, K.; Utimoto, K. *Tetrahedron Lett.* **1988**, *29*, 1041. (c) Murakami, M.; Kawano, T.; Ito, H.; Ito, Y. *J. Org. Chem.* **1993**, *58*, 1458. (d) Takai, K.; Ueda, T.; Ikeda, N.; Moriwake, T. *J. Org. Chem.* **1996**, *61*, 7990. (e) Takai, K.; Matsukawa, N.; Takahashi, A.; Fujii, T. *Angew. Chem., Int. Ed.* **1998**, *37*, 152. (f) Bazin, S.; Feray, L.; Siri, D.; Naubron, J.-V.; Bertrand, M. P. *Chem. Commun.* **2002**, 2506. (g) Yamamoto, Y.; Nakano, S.; Maekawa, H.; Nishiguchi, I. *Org. Lett.* **2004**, *6*, 799. (h) Godineau, E.; Landais, Y. *J. Am. Chem. Soc.* **2007**, *129*, 12662.

(3) For reviews on the radical-polar crossover reactions, see: (a) Ryu, I. *Chem. Rev.* **2002**, *2*, 249. (b) Murphy, J. A. *The Radical-Polar Crossover Reaction*. In *Radicals in Organic Synthesis*, Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, 2001; Vol. 1, pp 298. (c) Godineau, E.; Landais, Y. *Chem.—Eur. J.* **2009**, *15*, 3044.


(4) For reviews on synthesis of complex terpenoids, see: (a) Nicolaou, K. C.; Vourloumis, D.; Winssinger, N.; Baran, P. S. *Angew. Chem., Int. Ed.* **2000**, *39*, 44. (b) Maimone, T. J.; Baran, P. S. *Nature Chem. Biol.* **2007**, *3*, 396.

(5) Dong, S.-H.; Zhang, C.-R.; Xu, C.-H.; Ding, J.; Yue, J.-M. *J. Nat. Prod.* **2011**, *74*, 1255.

(6) Goetz, M. A.; Hensens, O. D.; Zink, D. L.; Borris, R. P.; Morales, F.; Tamayo-Castillo, G.; Slaughter, R. S.; Felix, J.; Ball, R. G. *Tetrahedron Lett.* **1998**, *39*, 2895.

tetrasubstituted carbon of the highly oxygenated cyclohexanol derivatives is linked to the tertiary carbon of the 2,3-*trans*-disubstituted cycloalkane structures (highlighted in blue and red). Here, we report development of a novel three-component coupling reaction using O,Te-acetal **1**, α,β -unsaturated ketone **2**, and aldehyde/ketone **3**. The single-step reaction proceeded through the radical-polar crossover mechanism under mild conditions and generated the complex ring connection motifs (**4**) of trigohownin A or correolide with stereoselective formation of the one C–O and two C–C bonds.

Scheme 1. Plan of Radical-Polar Crossover Reaction




A possible scenario of the radical-polar reaction is illustrated in Scheme 1. We selected O,Te-acetal **1** with the 2,4,10-trioxadamantane orthoester structure as the radical precursor and Et₃B as the radical initiator and terminator. Ethyl radical, which is generated by the reaction of Et₃B and O₂,⁷ would induce the homolytic cleavage

(7) (a) Nozaki, K.; Oshima, K.; Utimoto, K. *J. Am. Chem. Soc.* **1987**, *109*, 2547. For a review, see: (b) Ollivier, C.; Renaud, P. *Chem. Rev.* **2001**, *101*, 3415.

(8) (a) Barton, D. H. R.; Ozbalik, N.; Sarma, J. C. *Tetrahedron Lett.* **1988**, *29*, 6581. (b) He, W.; Togo, H.; Waki, Y.; Yokoyama, M. *J. Chem. Soc., Perkin. Trans. 1* **1998**, 2425.

Scheme 2. Preparation and Et₃B-Mediated Radical Reaction of O,Se-Acetal **A** and O,Te-Acetal **1**^a



^a EDCI = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide.

of the highly activated C–Te bond of **1** to produce α -alkoxy bridgehead radical **B**.^{8–12} Since the sterically unshielded radical **B** possesses potent reactivity as a nucleophilic radical, a congested C–C bond would be formed through the intermolecular addition of **B** to cycloalkenone **2**, leading to **C**. Radical intermediate **C** would then be converted to anionic intermediate **E** via the reaction with Et₃B and subsequent ejection of ethyl radical.^{2b,13} Finally, aldol reaction of the resultant boron enolate **E** with the carbonyl group of **3** would yield **4**. This attractive sequential reaction can only be realized when the reactivity of each component is completely differentiated and Et₃B enables the radical initiating (**1**→**B**) and trapping processes (**C**→**E**).

We previously reported the radical-based three-component coupling of O,Se-acetal **A**, α,β -unsaturated ketone **2**, and allyl tributyltin (Scheme 1).¹⁰ The C–Se bond of **A** was homolytically cleaved by the action of tributyltin radical at 110 °C, and two subsequent radical reactions led to product **D**. Thus, **A** was first treated with Et₃B, the key ingredient of the radical-polar crossover process (Scheme 2). However, the C–Se bond was found to be inert upon treatment with Et₃B/O₂ at 0 °C, and **A** was almost quantitatively recovered. Therefore, the C–Se bond in **A** was replaced with the weaker C–Te bond in the alternative

(9) For a review on bridgehead radicals, see: Walton, J. C. *Chem. Soc. Rev.* **1992**, *21*, 105.

(10) Urabe, D.; Yamaguchi, H.; Inoue, M. *Org. Lett.* **2011**, *13*, 4778.

(11) Murai, K.; Katoh, S.; Urabe, D.; Inoue, M. *Chem. Sci.* **2013**, *4*, 2364.

(12) Bridgehead radicals have been rarely used for synthetic studies on natural products; for example, see: (a) Kraus, G. A.; Andersh, B.; Su, Q.; Shi, J. *Tetrahedron Lett.* **1993**, *34*, 1741. (b) Kraus, G. A.; Su, Q. *Synlett* **1994**, 237. (c) Kraus, G. A.; Siclovan, T. M.; Watson, B. *Synlett* **1995**, 201. (d) Kim, S.; Cheong, J. H. *Synlett* **1997**, 947.

(13) Nozaki, K.; Oshima, K.; Utimoto, K. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 403.

(14) (a) Curran, D. P.; Martin-Esker, A. A.; Ko, S.-B.; Newcomb, M. *J. Org. Chem.* **1993**, *58*, 4691. (b) Smart, B. A.; Schiesser, C. H. *J. Chem. Soc., Perkin. Trans. 2* **1994**, 2269. (c) Schiesser, C. H.; Smart, B. A. *Tetrahedron* **1995**, *51*, 6051.


radical donor **1**.^{14,15} Condensation of carboxylic acid **5** with 2-mercaptopyridine *N*-oxide using EDCI yielded Barton ester **6**,¹⁶ which was immediately photoirradiated in the presence of (PhTe)₂ to afford the requisite O,Te-acetal **1**.^{17,18} The reaction of **1** and cyclopentenone **2a** (3 equiv) in the presence of Et₃B (3 equiv) under air in CH₂Cl₂ proceeded at 0 °C within 15 min to produce **7** in 85% yield. This facile adduct formation at low temperature without the toxic tin reagent^{19,20} clearly demonstrated the superiority of **1** over **A** as the radical donor. When CH₃OD (10 equiv) was present in the same reaction mixture, deuterated **7-d₁** (85% D) was exclusively obtained, indicating that the boron enolate was indeed formed and quenched by the acidic deuteron of CH₃OD.




Once the radical initiating and trapping ability of Et₃B was verified by the deuteration experiment, benzaldehyde **3a** was applied as the third component together with O,Te-acetal **1** and various cycloalkenones **2a–d** (Table 1). When a mixture of **1**, cyclopentenone **2a** and **3a** was treated with Et₃B/O₂ in CH₂Cl₂ at 0 °C for 15 min, the coupling adduct **4a** was produced through stereoselective installation of the three new stereocenters (89%, 9:1 dr at the benzylic position, entry 1). The radical addition to 4-TBSOxycyclopentenone **2b** proceeded selectively from the opposite face of the C4-substituent, giving rise to **4b** as a single diastereomer (entry 2). Moreover, 2-acetoxycyclopentenone **2c**²¹ underwent the three-component coupling with **1** and **3a** to provide **4c** (entry 3). Not only the cyclopentenone structures but also cyclohexenone **2d** (entry 4) was applicable as the radical acceptors. However, the aldol adduct **4d** was prone to undergo retroaldol reactions to produce large amounts of the two-component adduct **8** under the original conditions (entry 4). To increase the yield of the three-component adduct over the two-component counterpart, **4d** was silylated in the same pot (entry 5). As a result, TMS-ether **4e** was isolated in 73% yield (3:2 dr). It is worth noting that the four contiguous stereocenters in **4c** directly matched those of trigohownin A (circled in Scheme 1), and compound **4e** possessed the bicyclic

substructure of correolide. These results demonstrated the potential utility of the present one-pot reaction in building up the multiply oxygenated natural products.

The stereostructures of the coupling products **4a** and **4c** were established by X-ray crystallography, and the stereochemistries of **4b**, **4d**, and **4e** were confirmed by NMR experiments on derivatized compounds.²² Importantly,

Table 1. Three-Component Coupling of **1**, **2a–d**, and **3a**^a



entry	radical acceptor	electrophile	product	yield
1				89% (9:1) ^b
2	2b : X = H, Y = OTBS	3a	4b : X = H, Y = OTBS	86% ^c
3 ^d	2c : X = OAc, Y = H	3a	4c : X = OAc, Y = H	87% ^e
4 ^e			4d : R = H	27% ^{e,f}
5	2d	3a	4e : R = TMS	73% ^e (3:2)
				

^a Conditions: **1** (1 equiv), **2** (3 equiv), **3a** (3 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), 0 °C, 15 min. ^b **4a** was obtained as the major isomer. ^c Product was obtained as a single isomer. ^d Conditions: **1** (1 equiv), **2c** (2 equiv), **3a** (5 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), rt, 60 min. ^e Conditions: **1** (1 equiv), **2d** (2 equiv), **3a** (5 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), 0 °C, 15 min. ^f Compound **8** was obtained in 65% yield. ^g Conditions: **1** (1 equiv), **2d** (2 equiv), **3a** (5 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), 0 °C, 15 min; TMS-imidazole (10 equiv), DMAP (0.3 equiv), 0 °C, 3 h. Compound **4e** was obtained as the major isomer.

the stereochemical relationship of the newly introduced three centers of **4a–c** was consistent. This remarkable selectivity is explained by the six-membered transition state model depicted in Figure 1.²³ After the addition of bridgehead radical **B** to the radical acceptor and subsequent formation of boron enolate **E**, **3a** can only approach from the face opposite to the bulky trioxadamantane structure, thereby establishing the *trans*-relationship between the radical donor and the electrophile. The chairlike six-membered

(22) See the Supporting Information for details.

(23) (a) Fenzl, W.; Köster, R. *Liebigs Ann. Chem.* **1975**, 1322. (b) Inoue, T.; Mukaiyama, T. *Bull. Chem. Soc. Jpn.* **1980**, 53, 174. (c) Evans, D. A.; Nelson, J. V.; Vogel, E.; Taber, T. R. *J. Am. Chem. Soc.* **1981**, 103, 3099. For reviews on aldol reactions using boron enolate, see: (d) Mukaiyama, T. *Org. React.* **1982**, 28, 203. (e) Cowden, C. J.; Paterson, I. *Org. React.* **1997**, 51, 1.

(15) For reviews on radical reactions using organotellurium compounds, see: (a) Togo, H.; He, W.; Waki, Y.; Yokoyama, M. *Synlett* **1998**, 700. (b) Yamago, S. *Synlett* **2004**, 1875. (c) Petraghani, N.; Stefani, H. A. *Tellurium in Organic Synthesis*, 2nd, updated and enlarged ed.; Elsevier: New York, 2007.

(16) (a) Barton, D. H. R.; Crich, D.; Motherwell, W. B. *J. Chem. Soc., Chem. Commun.* **1983**, 939. For reviews on Barton esters, see: (b) Crich, D.; Quintero, L. *Chem. Rev.* **1989**, 89, 1413. (c) Saraiva, M. F.; Couri, M. R. C.; Hyaric, M. L.; de Almeida, M. V. *Tetrahedron* **2009**, 65, 3563.

(17) Barton, D. H. R.; Bridon, D.; Zard, S. Z. *Tetrahedron Lett.* **1984**, 25, 5777.

(18) For C–C bond formations using O,Te-acetals as radical precursors, see: (a) Barton, D. H. R.; Ramesh, M. *J. Am. Chem. Soc.* **1990**, 112, 891. (b) He, W.; Togo, H.; Yokoyama, M. *Tetrahedron Lett.* **1997**, 38, 5541. (c) Yamago, S.; Miyazoe, H.; Yoshida, J. *Tetrahedron Lett.* **1999**, 40, 2343. (d) Miyazoe, H.; Yamago, S.; Yoshida, J. *Angew. Chem., Int. Ed.* **2000**, 39, 3669. (e) Yamago, S.; Miyazoe, H.; Goto, R.; Hashidume, M.; Sawazaki, T.; Yoshida, J. *J. Am. Chem. Soc.* **2001**, 123, 3697. (f) Yamago, S.; Miyoshi, M.; Miyazoe, H.; Yoshida, J. *Angew. Chem., Int. Ed.* **2002**, 41, 1407. (g) Yamago, S.; Iida, K.; Yoshida, J. *J. Am. Chem. Soc.* **2002**, 124, 2874. See also ref 8b.

(19) Boyer, I. J. *Toxicology* **1989**, 55, 253.

(20) For a review on tin-free radical reactions, see: Baguley, P. A.; Walton, J. C. *Angew. Chem., Int. Ed.* **1998**, 37, 3072.

(21) (a) Maignan, C.; Rouessac, F. *Bull. Soc. Chim. Fr.* **1971**, 3041. (b) Lange, G. L.; Campbell, H. M.; Neidert, E. *J. Org. Chem.* **1973**, 38, 2117.




Figure 1. Rationale of the stereochemical outcome.

Table 2. Three-Component Couplings of **1**, **2a**, and **3b–l**^a

entry	radical acceptor	electrophile	product	yield ^b
1 ^c	2a	3b : R ¹ = Me	4f : R ¹ = Me	86%
2	2a	3c : R ¹ = <i>n</i> -C ₆ H ₁₃	4g : R ¹ = C ₆ H ₁₃	77%
3	2a	3d : R ¹ = cyclohexyl	4h : R ¹ = cyclohexyl	99%
4 ^d	2a	3e : R ¹ = <i>t</i> -Bu	4i : R ¹ = <i>t</i> -Bu	74%
5 ^e	2a	3f : R ¹ = (<i>E</i>)-CH=CHMe	4j : R ¹ = (<i>E</i>)-CH=CHMe	41%
6 ^f	2a	3g : R ¹ = CH=CHMe ₂	4k : R ³ = CH=CHMe ₂	76%
7 ^f				64%
8 ^f	2a	3i : <i>n</i> = 1	4m : <i>n</i> = 1	22% ^g
9 ^f	2a	3j : <i>n</i> = 2	4n : <i>n</i> = 2	86%
10	2a	3k : X = O	4o : X = O	90%
11	2a	3l : X = NBoc	4p : X = NBoc	92%

^a Conditions: **1** (1 equiv), **2a** (3 equiv), **3** (3 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), 0 °C, 15 min. ^b Product was obtained as a single isomer. ^c 10 equiv of **3b** was used. ^d Conditions: **1** (1 equiv), **2a** (2 equiv), **3** (5 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), rt, 60 min. ^e 2 equiv of **2a** was used. ^f Conditions: **1** (1 equiv), **2a** (2 equiv), **3** (5 equiv), Et₃B (3 equiv), CH₂Cl₂ (0.1 M), 0 °C, 15 min. ^g Compound **7** was obtained in 63% yield.

transition state **F** would then control the stereocenters of the α - and β -positions of the ketone, resulting in formation of the adduct **4** in a stereoselective fashion.

The broad scope of the present coupling was further demonstrated using **3b–l** as the third component (Table 2).

The reactions of enolizable acetaldehyde **3b** and *n*-heptanal **3c** with O,Te-acetal **1**, cyclopentenone **2a** and Et₃B/O₂ resulted in efficient formation of **4f** and **4g**, respectively, as the sole stereoisomers (entries 1 and 2). The sterically more cumbersome aliphatic aldehydes **3d** and **3e** also served as electrophiles without decreasing the yields to afford **4h** and **4i**, respectively (entries 3 and 4). Even the α,β -unsaturated aldehydes **3f** and **3g** were found to be applicable to the three-component coupling reactions (entries 5 and 6), although these compounds can function as both radical and carbanion acceptors. The bridgehead radical **B** was preferentially added to α,β -unsaturated ketone **2a** in the presence of **3f** and **3g**, and the resultant boron enolate selectively reacted with the aldehyde, leading to **4j** and **4k**, respectively. Significantly, the adducts **4f–k** were obtained as single isomers, further supporting the stereochemical rationale in Figure 1.

When acetone **3h** was mixed with **1** and **2a** in the presence of Et₃B/O₂, the in situ aldol reaction constructed a new tetrasubstituted carbon center to afford **4l** (entry 7). Furthermore, the three cyclic structures were connected in a single step upon using the cyclic ketones (entries 8–11). Cyclopentanone **3i** and cyclohexanone **3j** were coupled with **1** and **2a** to deliver the tricycyclic structures **4m** and **4n**, respectively (entries 8 and 9). The tetrahydropyran and piperidine structures were effectively incorporated into the coupling adducts **4o** and **4p** by using **3k** and **3l**, respectively, as the third components (entries 10 and 11). The smooth formation of **4l–p** under mild conditions reflected the generality of the present method for connecting the sterically hindered bonds between the three components.

In conclusion, we have developed a new three-component coupling reaction to construct **4** from O,Te-acetal **1**, α,β -unsaturated ketone **2** and aldehyde/ketone **3** by the action of Et₃B and O₂. The single-step reaction simultaneously constructed the bridgehead carbon center in a stereospecific fashion, and the contiguous three stereocenters in a stereoselective fashion. The tin-free procedure, the mild conditions, the broad substrate scope, the high stereoselectivity, and the efficient construction of the two hindered linkages are particularly useful features for the target-oriented synthesis. Because of these advantages, this three-component coupling introduces a novel and valuable strategy for streamlined synthesis of multiply functionalized terpenoids with complex architectures.

Acknowledgment. This research was financially supported by the Funding Program for Next Generation World-Leading Researchers (JSPS) to M.I. and a Grant-in-Aid for Young Scientists (B) (JSPS) to D.U. We thank S. Yoshioka, Dr. Y. Inokuma, and Prof. M. Fujita (all at The University of Tokyo) and Dr. K. Yoza (Bruker AXS) for X-ray crystallographic analyses.

Supporting Information Available. Experimental procedures and spectroscopic and analytical data for relevant compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.