

Multicomponent Prins Cyclization from Allylsilyl Alcohols Leading to Dioxaspirodecane

Asunción Barbero,* Alberto Diez-Varga, and Francisco J. Pulido

Departamento de Química Orgánica, Facultad de Ciencias, Campus Miguel Delibes,
47011 Valladolid, Spain

barbero@qo.uva.es

Received August 22, 2013

ABSTRACT



A multicomponent reaction for the preparation of dioxaspirodecanes starting from allylsilyl alcohols was achieved. The one-pot sequence involves the sequential acid-catalyzed reaction of an allylsilyl alcohol with an aldehyde to afford an alkenediol. The subsequent Prins cyclization of the homoallylic alcohol moiety generates a tetrahydropyranyl carbocation which is intramolecularly trapped by the second hydroxyl group. The chemoselectivity of the process shows dependence on the nature of the aldehyde and the concentration of the catalyst.

The stereoselective synthesis of substituted tetrahydropyran and oxepane scaffolds has attracted considerable attention, due to the presence of these units in a wide range of natural products with important biological activities,¹ such as the marine tetracyclic ether Hemibrevetoxin B,² isolated from dinoflagellate *Gymnodinium breve*, or the bromotriterpene polyether Armatol F,³ isolated from the red alga *Chondria armata*.

Among the many useful methods for the construction of these cyclic ethers, Prins cyclization has shown to be one of the most effective.⁴ Typically the process involves the acid-catalyzed reaction of an unsaturated alcohol with an aldehyde. The silyl-Prins modification has emerged as a very versatile approach for the synthesis of oxygenated

heterocycles. Keck⁵ and Wender⁶ and others⁷ have applied this protocol to the synthesis of macrocycles containing pyran units. As shown in Scheme 1, the acid-catalyzed reaction of an allylsilyl alcohol with an aldehyde proceeds in high yield and with total stereocontrol to provide the methylene-tetrahydropyran derivative.

An even more powerful strategy consists of generating the homoallylic alcohol in situ. Minehan⁸ has used this methodology in the synthesis of 4-methylene-tetrahydropyrans, via an initial Barbier allylation to generate a silylated homoallylic alcohol intermediate, which then undergo indium-mediated Prins cyclization. Chan⁹ and Saikia¹⁰ have described a multicomponent reaction involving a Sakurai allylation followed by an intramolecular silyl-Prins cyclization (Scheme 2).

(1) (a) Yasumoto, T.; Murata, M. *Chem. Rev.* **1993**, *93*, 1897–1909. (b) Fernandez, J. J.; Souto, M. L.; Norte, M. *Nat. Prod. Rep.* **2000**, *17*, 235–246.

(2) Prasad, A. V. K.; Shimizu, Y. *J. Am. Chem. Soc.* **1989**, *111*, 6476–6477.

(3) Ciavatta, M. L.; Wahidulla, S.; D'Souza, L.; Scognamiglio, G.; Cimino, G. *Tetrahedron* **2001**, *57*, 617–623.

(4) (a) Olier, C.; Kaafarani, M.; Gastaldi, S.; Bertrand, M. P. *Tetrahedron* **2010**, *66*, 413–445. (b) Pastor, I.; Yus, M. *Curr. Org. Chem.* **2007**, *11*, 925–957. (c) Pastor, I.; Yus, M. *Curr. Org. Chem.* **2012**, *16*, 1277–1312. (d) Han, X.; Peh, G.; Floreancig, P. E. *Eur. J. Org. Chem.* **2013**, 1193–1208.

(5) (a) Keck, G. E.; Kraft, M. B.; Truong, A. P.; Li, W.; Sanchez, C. C.; Kedei, N.; Lewin, N. E.; Blumberg, P. M. *J. Am. Chem. Soc.* **2008**, *130*, 6660–6661. (b) Keck, G. E.; Truong, A. P. *Org. Lett.* **2005**, *7*, 2149–2152. (c) Keck, G. E.; Truong, A. P. *Org. Lett.* **2005**, *7*, 2153–2156.

(6) (a) Wender, P. A.; DeChristopher, B. A.; Schrier, A. J. *J. Am. Chem. Soc.* **2008**, *130*, 6658–6659. (b) Wender, P. A.; Verma, V. A. *Org. Lett.* **2008**, *10*, 3331–3334.


(7) Within others, see: (a) Aubele, D. L.; Lee, C. A.; Floreancig, P. A. *Org. Lett.* **2003**, *5*, 4521–4523. (b) Dobbs, A. P.; Dunn, J. *Tetrahedron Lett.* **2012**, *53*, 2392–2395. (c) Furman, B.; Dziedzic, M.; Justyniak, I. *Tetrahedron* **2008**, *64*, 3103–3110.

(8) Pham, M.; Allatabakhsh, A.; Minehan, T. G. *J. Org. Chem.* **2008**, *73*, 741–744.


(9) (a) Wei, Z. Y.; Li, J. S.; Wang, D.; Chan, T. H. *Tetrahedron Lett.* **1987**, *28* (3), 441–444. (b) Wei, Z. Y.; Wang, D.; Li, J. S.; Chan, T. H. *J. Org. Chem.* **1989**, *54*, 5768–5774. (c) Chan, K.-P.; Loh, T.-P. *Tetrahedron Lett.* **2004**, *45*, 8387–8390.

(10) Reddy, U. C.; Rama Raju, B.; Pramod Kumar, E. K.; Saikia, A. K. *J. Org. Chem.* **2008**, *73*, 1628–1630.

Scheme 1. Silyl-Prins Cyclizations



Scheme 2. Multicomponent Sakurai–Prins Cyclization



Yet, there are rather few examples of Prins cyclization involving an internal trap of the pyranil carbocation.¹¹ These strategies usually involve the intramolecular Prins cyclization of an alkene which contains a homoallylic alcohol or amide bearing another nucleophilic atom (either oxygen or nitrogen) to give fused diheterobicyclic compounds.¹²

Following our interest in the development of new approaches to the synthesis of different sized carbo- and heterocycles, promoted by silicon-mediated cyclizations,¹³ we now present our recent results on the extension of this methodology to the synthesis of oxacycles using the Prins cyclization.

Here, we describe a new multicomponent Prins cyclization in which an alkenyl monoalcohol undergoes a tandem allylation–Prins cyclization followed by internal trapping of the tetrahydropyranil cation to give dioxaspirodecanes.

The requisite allylsilyl alcohols for this study were prepared by silylcupration of allene and capture of the intermediate cuprate with α,β -unsaturated aldehydes or ketones. Initially, we attempted the standard Prins cyclization of allylsilyl alcohol **1a** (1.0 equiv) with benzaldehyde (1.0 equiv) in the presence of TMSOTf (1.2 equiv) at -78°C . Surprisingly, we did not obtain the expected oxepane **3a**, from a direct Prins cyclization. Instead, the reaction provided a product which turned out to be the

shown dioxaspirodecane **2a**, along with unreacted hydroxyallylsilane **1a**. We then performed the same reaction using 2.0 equiv of benzaldehyde. The reaction then proceeded in high yield and with very high stereocontrol ($>95:5$) (a single stereoisomer could be detected in the reaction mixture).

Other Lewis acids, which are conventional in this type of cyclizations, were screened for this reaction. As shown in Table 1, AlCl_3 gave dioxaspirodecane **2a** in low yield (entry 3), whereas TiCl_4 gave a complex reaction mixture (entry 5) and Et_2AlCl gave also a complex mixture from which oxepane **3a** could be isolated in very low yield (entry 4). TMSOTf, at -78°C , was the most effective catalyst. This catalyst and these conditions were used for the subsequent reactions.

Table 1. Prins Cyclization from Allylsilyl Alcohols

entry	Lewis acid	temp ($^\circ\text{C}$)	ratio 2a/3a	yield ^a (%)
1	TMSOTf	-78	$>95:5$	85
2	TMSOTf	-40	$>95:5$	56
3	AlCl_3	-78	$>95:5$	64
4	Et_2AlCl	-78	$<5:95$	35
5	TiCl_4	-78	—	—

^a Conditions: **1a** (1.0 mmol), benzaldehyde (2.2 mmol), Lewis acid (1.2 mmol).

We next evaluated the scope of this process by employing a variety of alkyl, aryl, and α,β -unsaturated aldehydes and two different allylsilyl alcohols. The results are shown in Table 2.

As shown in Table 2, the product outcome is dependent on both the (arylic, vinylic, or alkylic) nature of the aldehyde and the electronic character of the substituents on it. Thus, there is a major predominance for the formation of the dioxaspirodecane derivatives **2** and the best diastereoselectivities ($>95:5$) are found for unsubstituted vinylic and aryl derivatives (Table 2, entries 1, 5, 9, and 13).


Yet, two different behaviors are found for aryl derivatives depending on the electronic character of the aromatic substituent. Thus, electron-rich aromatic aldehydes (Table 2, entries 2, 3, 10, 11) provide the corresponding dioxaspirodecanes **2b–c** and **2j–k** in high yield and with good to excellent diastereoselectivities.

However the reaction with electron-poor aromatic aldehydes led to a moderate selectivity toward **2** or even to a slight predominance of the oxepane derivative when nitrobenzaldehyde is used (Table 2, entries 4 and 12). Moreover, substituted vinylic aldehydes provided with good selectivities dioxaspirodecanes **2f–g** and **2n–o**

(11) Elsworth, J. D.; Willis, C. L. *Chem. Commun.* **2008**, 1587–1589.

(12) (a) Reddy, B. V. S.; Borkar, P.; Yadav, J. S.; Reddy, P. P.; Kunwar, A. C.; Sridhar, B.; Gree, R. *Org. Biomol. Chem.* **2012**, *10*, 1349–1358. (b) Reddy, B. V. S.; Jalal, S.; Borkar, P.; Yadav, J. S.; Reddy, P. P.; Kunwar, A. C.; Sridhar, B. *Org. Biomol. Chem.* **2012**, *10*, 6562–6568. (c) Reddy, B. V. S.; Narasimulu, G.; Reddy, Y. V.; Chakravarthy, P. P.; Yadav, J. S.; Sridhar, B. *Tetrahedron Lett.* **2012**, *53*, 3100–3103. (d) Reddy, B. V. S.; Borkar, P.; Chakravarthy, P. P.; Yadav, J. S.; Gree, R. *Tetrahedron Lett.* **2010**, *51*, 3412–3416. (e) Yadav, J. S.; Borkar, P.; Chakravarthy, P. P.; Reddy, B. V. S.; Sarma, A. V. S.; Basha, S. J.; Sridhar, B.; Gree, R. *J. Org. Chem.* **2010**, *75*, 2081–2084.

(13) (a) Barbero, A.; Pulido, F. J. *Acc. Chem. Res.* **2004**, *37*, 817–825. (b) Barbero, A.; Castreño, P.; Pulido, F. J. *J. Am. Chem. Soc.* **2005**, *127*, 8022–8023. (c) Barbero, A.; Castreño, P.; Pulido, F. J. *Org. Lett.* **2003**, *5*, 4045–4048. (d) Pulido, F. J.; Barbero, A.; Castreño, P. *J. Org. Chem.* **2011**, *76*, 5850–5855.

Table 2. Scope of the Multicomponent Prins Cyclization


entry	1	R ¹	R ²	ratio 2/3 ^a	product ^b (yield, %)
1	1a	Me	C ₆ H ₅	>95:5	2a (85)
2	1a	Me	4-MeOC ₆ H ₄	92:8	2b+3b (89)
3	1a	Me	4-MeC ₆ H ₄	>95:5	2c (85)
4	1a	Me	4-ClC ₆ H ₄	74:26	2d+3d (75)
5	1a	Me	CH ₂ =CH	>95:5	2e (70)
6	1a	Me	(<i>E</i>)- PhCH=CH	94:6	2f+3f (88)
7	1a	Me	(<i>E</i>)- MeCH=CH	83:17	2g+3g (87)
8	1a	Me	<i>c</i> -C ₆ H ₁₁	55:45	2h+3h (63)
9	1b	H	C ₆ H ₅	>95:5	2i (86)
10	1b	H	4-MeOC ₆ H ₄	89:11	2j+3j (80)
11	1b	H	4-MeC ₆ H ₄	87:13	2k+3k (87)
12	1b	H	4-NO ₂ C ₆ H ₄	40:60	2l+3l ^c (68)
13	1b	H	CH ₂ =CH	>95:5	2m (70)
14	1b	H	(<i>E</i>)- PhCH=CH	86:14	2n+3n (82)
15	1b	H	(<i>E</i>)- MeCH=CH	72:28	2o+3o (88)
16	1b	H	<i>c</i> -C ₆ H ₁₁	40:60	2p+3p (76)

^aThe ratio of products was determined by ¹H NMR (400 MHz).

^b Conditions: **1** (1.0 mmol), aldehyde (2.2 mmol), TMSOTf (1.2 mmol), at -78 °C. ^c The oxepane derivative was isolated as the regioisomeric 4-methyl-2-*p*-nitrophenyl-2,3,6,7-tetrahydro-oxepine.

(Table 2, entries 6, 7, 14, and 15). Interestingly, aliphatic aldehydes (Table 2, entries 8 and 16) furnished an almost equimolar mixture of both products **2** and **3**.



It has to be noted that the results with allylsilyl alcohols **1a** and **1b** are similar, although **1b** shows slightly lower selectivity toward the formation of the dioxaspirodecane derivatives.

The structures of **2b** and **2j** were unambiguously confirmed by NMR spectroscopy, HRMS analysis, and X-ray crystal structure analysis (Figure 1).

A mechanistical proposal for the formation of dioxaspirodecane **2** involves the acid-catalyzed Sakurai reaction between the hydroxyallylsilane and benzaldehyde to provide an alkoxy-alkenol **I** which then will react with another molecule of aldehyde to form an oxocarbenium (Scheme 3). Final cyclization will provide a pyranil carbocation which in turn will be intramolecularly trapped by the neighboring hydroxy group.

Yet, formation of methyleneoxepanes **3** would proceed through initial formation of an (*E*)-oxocarbenium ion **II**, which would be trapped by the nucleophilic allylsilane.

Presumably, whether the initial formation of intermediate **I** or **II** is the quickest process determines the final

**Figure 1.** X-ray crystal structures of **2b** and **2j**.**Scheme 3.** Mechanistical Proposal

outcome of the reaction. In this sense, the results shown in Table 2 seem to indicate that the reaction outcome depends on both steric and electronic factors. However, more investigation is needed to determine the real influence of these factors.

Yet, it has been described that the Hosomi–Sakurai reaction of allylsilanes and aldehydes catalyzed by TMSOTf is not favored, although in the presence of a silyl ether as a coreactant the reaction is readily promoted.¹⁴ A possible explanation for the success of our initial Sakurai reaction leading to **I** is that the active species is the catalytic amount of triflic acid present in TMSOTf solution.¹⁵

Moreover, in every case an excellent degree of diastereoselectivity was observed in the formation of both the 2,7-*cis*-oxepane and the 7,9-*cis*-dioxaspirodecane derivatives.

(15) Similar findings have been described by Hinkle in a ISMS reaction: Lambert, R. F.; Hinkle, R. J.; Ammann, S. E.; Lian, Y.; Liu, J.; Lewis, S. E.; Pike, R. D. *J. Org. Chem.* **2011**, *76*, 9269–9277.

(16) Alder, R. W.; Harvey, J. N.; Oakley, M. T. *J. Am. Chem. Soc.* **2002**, *124*, 4960–4961.

(14) (a) Mekhalfia, A.; Markó, I. E. *Tetrahedron Lett.* **1991**, *32*, 4779–4782.

The stereochemical outcome of Prins cyclization leading to tetrahydropyran derivatives has been rationalized by Alder,¹⁶ using density functional calculations. This model suggests that Prins cyclization proceeds through a chairlike transition state in which the substituents on C-2 and C-6 are equatorial for minimal repulsions. Moreover, calculations show that equatorial nucleophilic attack to the tetrahydropyranyl cation will be favored for secondary carbocations, while axial attack is predicted for tertiary pyranyl carbocations. It has to be noted that Alder's calculations have been done for intermolecular nucleophilic attack.

However, dioxaspirodecane **2** shows high selectivity for the equatorial trapping of the tertiary tetrahydropyranyl cation. The rationale for this unexpected selectivity could be found in the steric factors. Thus, the intramolecular trapping of the cation takes place on the less hindered face of the cyclic cation, since axial trapping will have an unfavorable 1,3-diaxial interaction.

We next decided to study the influence of the amount of Lewis acid in the outcome of the reaction. The results are shown in Table 3.


As shown in Table 3, the degree of selectivity toward the dioxaspirodecane derivatives is enhanced when the amount of Lewis acid increases. In this sense, our hypothesis that TfOH is the real initiator of the Hosomi reaction is in accordance with the rate effect on this process when the concentration of acid increases.

Interestingly, the reaction of allylsilyl alcohol **1b** with (*E*)-cinnamaldehyde gives an almost equimolar mixture of **2** and **3** when 0.8 equiv of TMSOTf is used, but a 15:1 ratio when the amount of Lewis acid is increased to 2.4 equiv. As far as we know, this chemoselective dependence on the concentration of Lewis acid has never been previously reported in Prins cyclizations.

In conclusion, we have reported an easy access to substituted dioxaspirodecanes in a very stereoselective manner, through a multicomponent Sakurai–Prins cyclization. To the best of our knowledge, this is the first described three-component Prins cyclization starting from an alkenyl alcohol. Moreover the chemoselectivity of this reaction toward the dioxaspirodecane (vs the oxepane derivative) is dependent on both stereoelectronic factors and the concentration of the Lewis acid.

This multicomponent coupling allows the synthesis of complex dioxaspirodecanes in a sequence where three

Table 3. Influence of the Amount of Catalyst in Prins Cyclization



entry	1 R ¹	R ² -CHO R ²	TMSOTf equiv	ratio 2/3 ^a	yield ^b (%)
1	Me	(<i>E</i>)- MeCH=CH	0.8	78:22	89
2	Me	(<i>E</i>)- MeCH=CH	1.2	83:17	87
3	Me	(<i>E</i>)- MeCH=CH	2.4	87:13	85
4	Me	(<i>E</i>)- MeCH=CH	4.8	90:10	86
5	H	(<i>E</i>)- PhCH=CH	0.8	57:43	80
6	H	(<i>E</i>)- PhCH=CH	1.2	86:14	82
7	H	(<i>E</i>)- PhCH=CH	2.4	94:6	81
8	H	4-MeC ₆ H ₄	0.8	79:21	89
9	H	4-MeC ₆ H ₄	1.2	87:13	87
10	H	4-MeC ₆ H ₄	2.4	91:9	86

^aThe ratio of products was determined by ¹H NMR (400 MHz).
^b Conditions: **1** (1.0 mmol), aldehyde (2.2 mmol), –78 °C.

new stereogenic centers are created with excellent stereoselectivity.

Acknowledgment. We thank the Ministry of Science and Technology of Spain (CTQ2009-09302) and the “Junta de Castilla y León” (GR170) for financial support. We are very grateful to L.T.I. (UVA) and to Dr. D. Miguel (Department of Inorganic Chemistry, University of Valladolid) for X-ray determination.

Supporting Information Available. Experimental details and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.