


Iron-Catalyzed Reductive Cyclization of 1,6-Enynes

Aijun Lin, Zhi-Wei Zhang, and Jiong Yang*

Department of Chemistry, Texas A&M University, College Station, Texas 77843-3255, United States


S Supporting Information

ABSTRACT: A precatalyst of FeCl₂ and iminopyridine was activated in situ by a combination of diethylzinc and magnesium bromide etherate; it catalyzed the reductive cyclization of 1,6-enynes to give pyrrolidine and tetrahydrofuran derivatives from *N*- and *O*-tethered 1,6-enynes. The scope of the transformation was explored.



Iron catalysis for organic synthesis has experienced significant growth in recent decades.¹ Compared with other late transition metals, iron stands out because it is one of the most abundant elements in the Earth's crust, and most of iron salts are nontoxic. Thus, development of iron-based catalysis is desirable because of its relevance to "green" chemical processes and to a more sustainable chemical industry. Further, iron catalysis frequently shows reactivities and selectivities complementary to other transition metal catalysis: facets that can be exploited synthetically. The reductive cyclization and cycloisomerization of 1,6-enynes are powerful approaches for the synthesis of five-membered carbo- and heterocycles from acyclic substrates (Scheme 1).² Whereas a number of transition

Scheme 1. Reductive Cyclization and Cycloisomerization of 1,6-Enynes



metal catalysts have been developed for these transformations, very few examples of iron catalysis exist.³ Specifically, Fürstner described the Alder-ene cycloisomerization of 1,6-enynes using a well-defined low-valent organoiron complex [Li(TMEDA)]-[(η^5 -C₅H₅)Fe(C₂H₄)₂] as the catalyst,⁴ and the reductive cyclization of 1,6-enynes with the bis(imino)pyridine iron bis(dinitrogen) complex [(*i*-PrPDI)Fe(N₂)₂] was reported by Chirik.^{5,6} Both of these transformations make use of air-sensitive organoiron complexes that have to be separately prepared, and the latter also has to be carried out under an atmosphere of H₂ under pressure. As part of our efforts to develop practical and sustainable iron-catalyzed transformations, herein we report an operationally simple approach for iron-catalyzed reductive cyclization of unactivated 1,6-enynes to give pyrrolidine and tetrahydrofuran derivatives. This trans-


formation was enabled by the discovery of Et₂Zn and MgBr₂·Et₂O as a unique combination for activating a precatalyst consisting of FeCl₂ and a bidentate iminopyridine ligand.

At the outset of our research, we envisioned that a catalytically competent low-valent iron species might be formed by the reaction of FeCl₂ and the Grignard reagents.⁷ Thus, we commenced with the reductive enyne cyclization of the *N*-tethered 1,6-enyne **1a** by treatment with FeCl₂ and cyclopentylmagnesium bromide in THF (Table 1, entry 1). The initial results were promising as the desired cyclization product **2a** was obtained in 20% yield as a single (*Z*)-isomer. Similar yields were obtained when ethylmagnesium bromide (entry 2) or isopropylmagnesium bromide (entry 3) were used. In an effort to improve reaction efficiency, we also tested diethylzinc as a reducing agent for generating the catalytic iron species. Whereas diethylzinc alone gave **2a** in 26% yield at room temperature over 12 h or in 43% yield at 60 °C over 6 h (entries 4 and 5), addition of MgBr₂·Et₂O and ligand **L1** led to a dramatic improvement of the reaction,⁸ which was complete in 6 h at room temperature to give **2a** in 78% yield (entry 8). The same product was obtained with a comparable yield when FeCl₂ of 99.99% purity was employed (entry 9), indicating that the transformation was indeed under iron catalysis.⁹ Interestingly, low conversion of the reaction was observed when **L1** was used as the sole additive (entry 6), and rapid degradation of the starting material occurred when MgBr₂·Et₂O was used by itself (entry 7). Reduction of the amount of either Et₂Zn or FeCl₂ led to decreased yield (entries 10 and 11). Further screening of the reaction conditions, which included testing various ligands (**L2**–**L7**, entries 10–15), iron precatalysts (Table S1, Supporting Information), and solvents (Table S1, Supporting Information), led us to choose the conditions shown in entry 8 for all subsequent reactions.¹⁰

The scope of the reaction was evaluated using various substrates, and the results are shown in Figure 1. *N*-Tethered 1,6-enynes with a methoxy group at either the *para*- or *meta*-position of the aryl group were found to be compatible with the

Received: November 11, 2013

Published: December 26, 2013

Table 1. Screening of Reaction Conditions^a


entry	L	RM	additive	time (h)	yield ^b 2a (%)
1 ^c	—	C ₅ H ₅ MgBr	—	6	20
2 ^c	—	EtMgBr	—	6	17
3 ^c	—	<i>i</i> -PrMgBr	—	6	22
4	—	Et ₂ Zn	—	12	26
5 ^d	—	Et ₂ Zn	—	6	43
6	L1	Et ₂ Zn	—	16	<10
7	—	Et ₂ Zn	MgBr ₂ ·Et ₂ O	16	trace
8	L1	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	78
9 ^e	L1	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	75
10 ^f	L1	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	56
11 ^g	L1	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	43
12	L2	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	41
13	L3	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	29
14	L4	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	33
15	L5	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	18
16	L6	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	30
17	L7	Et ₂ Zn	MgBr ₂ ·Et ₂ O	6	22

^aReactions typically carried out with 0.2 mmol of **1a**, 20 mol % of FeCl₂, 20 mol % of the ligand, 3.0 equiv of Et₂Zn, and 1.0 equiv of MgBr₂·Et₂O in 3.0 mL of THF at room temperature. ^bIsolated yield. ^c1.5 equiv of RMgBr was used. ^dThe reaction was carried out at 60 °C. ^eFeCl₂ of 99.99% purity was used. ^f2.0 equiv of Et₂Zn was used. ^g10 mol % of FeCl₂ was used.

reaction, and the reductive cyclization products were obtained in good yields (72% yield for **2b** and 81% yield for **2c**). However, the same substituent at the *ortho*-position of the aryl led to formation of **2d** with significantly reduced efficiency (29%). This was attributed to unfavorable interactions of the lone pair of the methoxy group and the catalytic center. Indeed, the corresponding *ortho*-methyl substrate gave the product (**2e**) in 69% yield.¹¹ Electron-withdrawing chloro- or trifluoromethyl groups at the *para*-position of the aryl led to products in 68% (for **2f**) and 64% (for **2g**) yield, respectively, and the chloride substituent remained intact under the reaction conditions. Disubstitution at the 3'- and 5'-positions of the aryl with two methyl groups caused a somewhat reduced yield (**2h**, 55%). The reaction also proceeded smoothly with a thiophene-containing 1,6-enyne substrate to give **2i** in 52% yield. An alkyl-substituted enyne substrate was found to give the desired product **2j** in 41% yield. The *N*-tosyl of the *N*-tethered enyne substrate could be replaced with the *N*-Cbz, and the reductive cyclization product **2k** was obtained in 57% yield. We further tested this transformation using *O*-tethered 1,6-enynes. Despite





Figure 1. Substrate scope. Reactions were carried out with 20 mol % of FeCl₂, 20 mol % of L1, 3.0 equiv of Et₂Zn, and 1.0 equiv of MgBr₂·Et₂O in THF at room temperature.


the potential competing reductive elimination pathway with these substrates, methylene tetrahydrofurans **2l–2n** were obtained in good to moderate yields under the reaction conditions. The Thorpe–Ingold effect likely contributed to the higher efficiency for formation of **2o**.¹² Our experiments also revealed a limitation of the reaction. Despite our efforts, the enyne substrate with an internal alkene failed to deliver the desired product **2p** under the current reaction conditions. A diethyl malonate-tether substrate also failed to deliver the expected product **2q**.

Although the exact nature of the iron catalyst is unknown, we speculate that the reductive cyclization of **1a** proceeds as shown in Scheme 2. It starts with formation of metallacycle **A** by oxidative cyclization of **1a** with a low-valent iron species, which was generated from FeCl₂ and MgBr₂·Et₂O with Et₂Zn in the presence of L1. Transmetalation of **A** with Et₂Zn cleaves the metallacycle and gives intermediates **B** and **C**, which can convert to **D** and **E** through rapid β -hydride elimination followed by reductive elimination of the iron hydride intermediates. The catalytic iron species is also regenerated. Protonolysis of **D** and **E** gave **2a**.¹³ This proposed mechanistic pathway is consistent with the following experimental observation: The monodeuterated products **3** and **4** and the undeuterated **2a** were obtained in 49:31:19 ratio from **1a** upon quenching the reaction with D₂O (Scheme 3). Whereas deuteration of **D** and **E** led to **3** and **4**, formation of a small

Scheme 2. Proposed Reaction Mechanism



Scheme 3. Deuteration Study



amount of the undeuterated **2a** likely proceeded by transmetalation of **D** and **E** with an iron species to give **F** and **G**. Compound **2a** was formed upon β -hydride and reductive eliminations of **F** and **G**. For reasons yet to understand, our attempts of trapping the organometallic intermediates with common electrophiles have been unsuccessful.¹⁴

In summary, we have developed an operationally simple iron-catalyzed reductive cyclization of 1,6-enynes. This transformation was enabled by the unique combination of diethylzinc and $\text{MgBr}_2 \cdot \text{Et}_2\text{O}$ for converting the precatalyst FeCl_2 and the α -iminopyridine ligand into the catalytically competent species in situ. Functionalized pyrrolidine and tetrahydrofuran derivatives were obtained from *N*- and *O*-tethered 1,6-enynes in good-to-moderate yields. We elucidated the scope of the reaction and revealed its limitations. Further exploration of the utility of this iron-based catalytic system and elucidation of its exact mechanism of action will be reported in due course.

■ ASSOCIATED CONTENT

Supporting Information

Experimental details and NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: yang@mail.chem.tamu.edu.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the National Science Foundation (CHE-1150606) and the Robert A. Welch Foundation (A-1700) for financial support.

■ REFERENCES

- (1) For some reviews: (a) Bolm, C.; Legros, J.; Le Pailh, J.; Zani, L. *Chem. Rev.* **2004**, *104*, 6217–6254. (b) *Iron Catalysis in Organic Chemistry*; Plietker, B., Ed.; Wiley-VCH: Weinheim, Germany, 2008. (c) Enthaler, S.; Junge, K.; Beller, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 3317–3321. (d) Sherry, B. D.; Fürstner, A. *Acc. Chem. Res.* **2008**, *41*, 1500–1511. (e) Czaplik, W. M.; Mayer, M.; Cvengros, J.; von Wangelin, J. A. *ChemSusChem* **2009**, *2*, 396–417. (f) Nakamura, E.; Yoshikai, N. *J. Org. Chem.* **2010**, *75*, 6061–6067. (g) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. *Chem. Rev.* **2011**, *111*, 1293–1314.
- (2) For some recent reviews: (a) Michelet, V.; Toullec, P. Y.; Genêt, J.-P. *Angew. Chem., Int. Ed.* **2008**, *47*, 4268–4315. (b) Watson, I. D. G.; Toste, F. D. *Chem. Sci.* **2012**, *3*, 2899–2919. (c) Marinetti, A.; Jullien, H.; Voituriez, A. *Chem. Soc. Rev.* **2012**, *41*, 4884–4908. (d) Aubert, C.; Fensterbank, L.; Garcia, P.; Malacria, M.; Simmonneau, A. *Chem. Rev.* **2011**, *111*, 1954–1993. (e) Belmont, P.; Parker, E. *Eur. J. Org. Chem.* **2009**, 6075–6089. (f) Shen, H. C. *Tetrahedron* **2008**, *64*, 7847–7870. (g) Jiménez-Núñez, E.; Echavarren, A. M. *Chem. Rev.* **2008**, *108*, 3326–3350. For some examples: (h) Park, J. H.; Kim, S. M.; Chung, Y. K. *Chem.—Eur. J.* **2011**, *17*, 10852–10856. (i) Fan, B.-M.; Xie, J.-H.; Li, S.; Wang, L.-X.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2007**, *46*, 1275–1277. (j) Jang, H.-Y.; Hughes, F. W.; Gong, H.; Zhang, J.; Brodbelt, J. S.; Krische, M. J. *J. Am. Chem. Soc.* **2005**, *127*, 6174–6175. (k) Jang, H.-Y.; Krische, M. J. *J. Am. Chem. Soc.* **2004**, *126*, 7875–7880. (l) Montchamp, J.-L.; Negishi, E.-i. *J. Am. Chem. Soc.* **1998**, *120*, 5345–5346.
- (3) (a) Nieto-Oberhuber, C.; Muñoz, M. P.; López, S.; Jiménez-Núñez, E.; Nevado, C.; Herrero-Gómez, E.; Raducan, M.; Echavarren, A. M. *Chem.—Eur. J.* **2006**, *12*, 1677–1693. (b) Takacs, J. M.; Newsome, P. W.; Kuehn, C.; Takusagawa, F. *Tetrahedron* **1990**, *46*, 5507–5522. (c) Takacs, J. M.; Anderson, L. G. *J. Am. Chem. Soc.* **1987**, *109*, 2200–2202. (d) Takacs, J. M.; Weidner, J. J.; Takacs, B. E. *Tetrahedron Lett.* **1993**, *34*, 6219–6222. (e) Takacs, J. M.; Weidner, J. J.; Newsome, P. W.; Takacs, B. E.; Chidambaram, R.; Shoemaker, R. J. *Org. Chem.* **1995**, *60*, 3473–3476. (f) Takacs, J. M.; Myoung, Y.-C.; Anderson, L. G. *J. Org. Chem.* **1994**, *59*, 6928–6942. (g) Takacs, J. M.; Anderson, L. G.; Newsome, P. W. *J. Am. Chem. Soc.* **1987**, *109*, 2542–2544. (h) Pearson, A. J.; Dubbert, R. A. *Organometallics* **1994**, *13*, 1656–1661.
- (4) (a) Fürstner, A.; Majima, K.; Martín, R.; Krause, H.; Kattnig, E.; Goddard, R.; Lehmann, C. W. *J. Am. Chem. Soc.* **2008**, *130*, 1992–2004. (b) Fürstner, A.; Martín, R.; Majima, K. *J. Am. Chem. Soc.* **2005**, *127*, 12236–12237.
- (5) (a) Sylvester, K. T.; Chirik, P. J. *J. Am. Chem. Soc.* **2009**, *131*, 8772–8774. (b) Hoyt, J. M.; Sylvester, K. T.; Semproni, S. P.; Chirik, P. J. *J. Am. Chem. Soc.* **2013**, *135*, 4862–4877.
- (6) For related reports of iron-catalyzed reductive cyclization of activated 1,6-enynes: (a) Hata, T.; Hirone, N.; Sujaku, S.; Nakano, K.; Urabe, H. *Org. Lett.* **2008**, *10*, 5031–5033. (b) Hata, T.; Sujaku, S.; Hirone, N.; Nakano, K.; Imoto, J.; Imade, H.; Urabe, H. *Chem.—Eur. J.* **2011**, *17*, 14593–14602.
- (7) For studies of the so-called “inorganic Grignard reagent” by reaction of FeX_2 and RMgX , see: (a) Bogdanovic, B.; Schwickardi, M. *Angew. Chem., Int. Ed.* **2000**, *39*, 4610–4612. (b) Aleandri, L. E.; Bogdanovic, B.; Bons, P.; Dürr, C.; Gaidies, A.; Hartwig, T.; Hockett, S. C.; Lagarden, M.; Wilczok, U.; Brand, R. A. *Chem. Mater.* **1995**, *7*,

1153–1170. (c) Siedlaczek, G.; Schwickardi, M.; Kolb, U.; Bogdanović, B.; Blackmond, D. G. *Catal. Lett.* **1998**, *55*, 67–72. (d) Fürstner, A.; Martin, R.; Krause, H.; Seidel, G.; Goddard, R.; Lehmann, C. W. *J. Am. Chem. Soc.* **2008**, *130*, 8773–8787.

(8) For FeCl₂-iminopyridine complexes, see: (a) Gibson, V. C.; O'Reilly, R. K.; Wass, D. F.; White, A. J. P.; Williams, D. J. *Dalton Trans.* **2003**, 2824–2830. (b) Wu, J. Y.; Moreau, B.; Ritter, T. *J. Am. Chem. Soc.* **2009**, *131*, 12915–12917.

(9) Buchwald, S. L.; Bolm, C. *Angew. Chem., Int. Ed.* **2009**, *48*, 5586–5587.

(10) Formation of material of high polarity, which we tentatively attributed to polymerization of the substrate, was also observed under the reaction conditions. For iron-catalyzed polymerization of olefins, see: (a) Small, B. L.; Brookhart, M.; Bennett, A. M. A. *J. Am. Chem. Soc.* **1998**, *120*, 4049–4050. (b) Britovsek, G. J. P.; Gibson, V. C.; McTavish, S. J.; Solan, G. A.; White, A. J. P.; Williams, D. J.; Kimberley, B. S.; Maddox, P. J. *Chem. Commun.* **1998**, 849–850. (c) Raynaud, J.; Wu, J. Y.; Ritter, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 11805–11808.

(11) We thank an anonymous reviewer for suggesting the experiment with **2e**.

(12) (a) Beesley, R. M.; Ingold, C. K.; Thorpe, J. F. *J. Chem. Soc.* **1915**, *107*, 1080–1106. (b) Ingold, C. K. *J. Chem. Soc.* **1921**, *119*, 305–329. (c) Ingold, C. K.; Sako, S.; Thorpe, J. F. *J. Chem. Soc.* **1922**, 1177–1198.

(13) For a similar mechanistic hypothesis of a nickel-catalyzed reductive cyclization of 1,6-enynes, see: Chen, M.; Weng, Y.; Guo, M.; Zhang, H.; Lei, A. *Angew. Chem., Int. Ed.* **2008**, *47*, 2279–2282.

(14) The following aldehydes (3 equiv) were used as the electrophile: 4-trifluoromethylbenzaldehyde, 4-methoxybenzaldehyde, and hydrocinnamaldehyde.