

Cross-Coupling Hydrogen Evolution Reaction in Homogeneous Solution without Noble Metals

Jian-Ji Zhong, Qing-Yuan Meng, Bin Liu, Xu-Bing Li, Xue-Wang Gao, Tao Lei, Cheng-Juan Wu, Zhi-Jun Li, Chen-Ho Tung, and Li-Zhu Wu*

Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry & University of Chinese Academy of Sciences, the Chinese Academy of Sciences, Beijing, 100190, P. R. China

S Supporting Information

ABSTRACT: A highly efficient noble-metal-free homogeneous system for a cross-coupling hydrogen evolution (CCHE) reaction is developed. With cheap, earth-abundant eosin Y and molecular catalyst $\text{Co}(\text{dmgH})_2\text{Cl}_2$, good to excellent yields for coupling reactions with a variety of isoquinolines and indole substrates and H_2 have been achieved without any sacrificial oxidants. Mechanistic insights provide rich information on the effective, clean, and economic CCHE reaction.



Developing efficient and atom-economic methods for C–C bond formation is always an important concern in synthetic chemistry.^{1,2} The cross-dehydrogenative coupling reaction³ is a desirable synthetic approach toward C–C bond formation, as it does not require additional functionalization of subcomponents. Reactions of this type often involve the use of stoichiometric sacrificial oxidants, which leads to low atom economy and possible generation of toxic wastes. A new type of reaction, cross-coupling hydrogen evolution (CCHE) reaction,⁴ has recently appeared, which avoids the use of any sacrificial oxidants and yields hydrogen gas (H_2) as the sole byproduct. By combining eosin Y and a graphene-supported RuO_2 nanocomposite (G-RuO_2)⁵ as a photosensitizer and a catalyst, the CCHE reaction proceeds smoothly to afford the desired cross-coupling products and H_2 in quantitative yields under visible light irradiation. This novel reaction offers a potential solution for highly efficient, clean, and economical C–C bond formation.

In an effort to expand our knowledge and to develop a new system of the CCHE reaction, we replaced the noble metal heterogeneous catalyst G-RuO_2 with inexpensive molecular catalysts. In this contribution, we report an elegant new catalytic system for the CCHE reaction in a homogeneous solution. Here, an earth-abundant $\text{Co}(\text{dmgH})_2\text{Cl}_2$ (dmgH = dimethylglyoximate) complex is used as a catalyst to capture the electrons and protons eliminated from the C–H bonds of the substrates (Scheme 1). For a systematic comparison of the G-RuO_2 system, eosin Y is employed as a photosensitizer to initiate the cross-coupling of *N*-phenyl-tetrahydroisoquinolines and indoles, a typical cross-coupling reaction under visible-light catalysis.^{6,7}

Our initial study focused on the cross-coupling of *N*-phenyl-tetrahydroisoquinolines and indoles for their prevalent skeleton in natural products.⁸ After 1 mol % of eosin Y and 5 mol % of $\text{Co}(\text{dmgH})_2\text{Cl}_2$ were added into a mixture of CH_3CN and H_2O containing *N*-phenyl-tetrahydroisoquinoline **1a** and indole **2a**,

Scheme 1. Cross-Coupling Hydrogen Evolution in Homogeneous Solution without Noble Metals



the reaction mixture was deaerated by bubbling nitrogen and then irradiated by green LEDs ($\lambda = 525 \pm 10$ nm) at room temperature. Significantly, 55% and 63% yields of the desired cross-coupling product and H_2 , respectively, could be obtained with no use of any sacrificial oxidants (Table 1, entry 1). And the cross-coupling product and H_2 are the only products detected. Increasing the amount of $\text{Co}(\text{dmgH})_2\text{Cl}_2$ and eosin Y resulted in a large improvement in both conversion and yield of the reaction (Table 1, entries 2–5). When 3 mol % of eosin Y and 8 mol % of $\text{Co}(\text{dmgH})_2\text{Cl}_2$ were used, the conversion approached 100% and the yields of the cross-coupling product and H_2 were achieved in yields of 94% and 98%, respectively. When the CCHE reaction proceeded in H_2O , however, only a 12% yield of the cross-coupling product and a 16% yield of H_2 were obtained (Table 1, entry 6). Addition of an organic solvent greatly enhanced the performance of the reaction (Table 1, entries 5, 7, 8). Among various organic solvents, CH_3CN was the best for the CCHE reaction. H_2O is also of significance in the reaction. When the CCHE reaction was carried out in CH_3CN , a negligible amount of desired cross-

Received: February 19, 2014

Published: March 14, 2014

Table 1. Optimization of the Reaction Conditions^a

entry	EY	cat.	solvent ^b	convn ^c	H ₂ (%) ^d	3a (%) ^e
1	1%	5%	H ₂ O/CH ₃ CN	65	63	55
2	1%	8%	H ₂ O/CH ₃ CN	80	76	69
3	1%	10%	H ₂ O/CH ₃ CN	82	79	71
4	3%	8%	H ₂ O/CH ₃ CN	100	98	94
5	5%	8%	H ₂ O/CH ₃ CN	100	97	91
6	3%	8%	H ₂ O	18	16	12
7	3%	8%	H ₂ O/DMF	45	43	39
8	3%	8%	H ₂ O/CH ₃ OH	55	54	50
9 ^f	3%	8%	CH ₃ CN	6	3	5

^aReaction conditions: 0.1 mmol of *N*-phenyl-tetrahydroisoquinoline **1a**, 0.2 mmol of indole **2a**, corresponding eosin Y, and cat. (Co(dmgH)₂Cl₂) were added, and the solution was strictly deaerated and irradiated for 16 h by 525 nm green LEDs, with methane as the internal standard. ^bH₂O/CH₃CN 4:1. ^cCorresponding to **1a**. ^dYields detected by gas chromatography using pure methane as an internal standard based on **1a**. ^eYields detected by NMR using 4-nitroacetophenone as an internal standard based on **1a**. ^fCo(dmgH)₂pyCl instead of Co(dmgH)₂Cl₂.

coupling product and H₂ were detected (Table 1, entry 9). When D₂O instead of H₂O was used as the solvent, D₂ was obtained as the only byproduct with no alteration of the reaction efficiency (Scheme S1), possibly a result of a quick exchange of H₂O with the released proton from the substrate. And the best ratio of H₂O to CH₃CN was found to be 4:1 (Table S1). Further experiments indicated that the conversion of **1a** was negligible when eosin Y or Co(dmgH)₂Cl₂ was absent from the reaction mixture, and no conversion of **1a** could be detected if the reaction was conducted in darkness.

Given that noble-metal G-RuO₂⁴ was replaced by the inexpensive Co(dmgH)₂Cl₂ molecular catalyst, we compared the two systems and found that the amount of eosin Y was decreased from 20 to 3 mol % for the CCHE reaction in the homogeneous solution, and simultaneously **1a** could be completely converted to the desired cross-coupling product and H₂ in a shorter reaction time but with higher yields (Table 1, entry 4). For the indole with an ester group that showed the lowest reactivity in the G-RuO₂ heterogeneous system,⁴ the reactivity of this noble-metal-free homogeneous reaction system enhanced four times (Scheme 3). Moreover, the homogeneous reaction can be carried out in a mixture of organic solvent and water. Therefore, it is possible to expand the scope of water-insoluble substrates for the CCHE reaction.

The catalyst changed from G-RuO₂ to Co(dmgH)₂Cl₂, which resulted in greatly improved catalytic performance, prompting us to shed light on the mechanism of the CCHE reaction. For consistency regarding reaction conditions, all subsequent mechanistic studies were performed under the optimal conditions. Although no obvious change could be detected in the UV-vis absorption spectrum of each component, eosin Y, **1a**, Co(dmgH)₂Cl₂, and indoles as well as their mixtures, these spectra suggested that only eosin Y in the reaction mixture could be excited by green LED light (SI, Figure S1). In this situation, two reaction pathways are possible. One is that the photoexcited eosin Y might be oxidatively

quenched by Co(dmgH)₂Cl₂ to generate [eosin Y]^{•+} for the oxidation of **1a**. The other is that the photoexcited eosin Y could be reductively quenched by **1a** to produce [eosin Y]^{•-}, which delivers an electron to the Co(dmgH)₂Cl₂ catalyst for proton reduction to H₂. To determine the dominant reaction pathway in the system, a flash photolysis investigation was undertaken in terms of the rich spectroscopic property of eosin Y.⁹ As shown in Figure 1a, immediately after a laser excitation




Figure 1. (a) Transient absorption spectra observed after laser excitation ($\lambda_{\text{ex}} = 532$ nm) of system containing (1) eosin Y (1.0×10^{-5} M); (2) eosin Y (1.0×10^{-5} M), **1a** (3.3×10^{-4} M); (3) eosin Y (1.0×10^{-5} M), **1a** (3.3×10^{-4} M), Co(dmgH)₂Cl₂ (2.7×10^{-5} M). The concentration ratio of the substrate is identical to that in the reaction. (b) Spectroelectrochemical absorption spectrum of **1a**^{•+} in CH₃CN/H₂O by oxidation of **1a** (1.0×10^{-5} M) at 0.90 V voltage relative to NHE. (c and d) UV-vis absorption spectra of system containing **1a** (4.0×10^{-4} M), **2a** (8.0×10^{-4} M), eosin Y (1.2×10^{-5} M), Co(dmgH)₂Cl₂ (3.2×10^{-5} M) in H₂O/CH₃CN 4:1 before and after irradiation by green LEDs (525 ± 10 nm).


by 532 nm light, a strong negative bleach of the ground state absorption of eosin Y at approximately 520 nm and the characteristic absorption assigned to the triplet excited state (³[eosin Y]^{*}) at 560 nm were observed. When *N*-phenyl-tetrahydroisoquinoline **1a** was introduced into the solution of eosin Y, a new absorption band, which is in accordance with the eosin Y radical anion ([eosin Y]^{•-}), appeared at 410 nm at the expense of the ³[eosin Y]^{*} absorption at 560 nm. The result suggested that the electron transfer from **1a** to ³[eosin Y]^{*} took place. In contrast, introduction of Co(dmgH)₂Cl₂ into the eosin Y solution with 532 nm laser irradiation resulted in no change of the ³[eosin Y]^{*} absorption (SI, Figure S2c). Clearly, the reductive quenching pathway of the photoexcited ³[eosin Y]^{*} dominates the homogeneous catalytic system.

Further addition of Co(dmgH)₂Cl₂ into a mixture of eosin Y and **1a** in a 4:1 H₂O/CH₃CN solution led to a distinct quenching of [eosin Y]^{•-} upon laser excitation at 532 nm. The lifetime of [eosin Y]^{•-} changed from 39.8 to 4.3 μs indicating the electron transfer from [eosin Y]^{•-} to Co(dmgH)₂Cl₂. Simultaneously, a new absorption with a maximum appeared at 370 nm in the transient absorption spectra. The emerged absorption band is similar to that of **1a** under oxidation potential at 0.90 V versus NHE in CH₃CN/H₂O (Figure 1b). The result indicated that the **1a**^{•+} at 370 nm is generated by electron transfer from **1a** to the excited eosin Y. This is, to the best of knowledge, the first time the *N*-phenyl-tetrahydroisoquinoline radical cation was observed directly in the cross-coupling reactions. Importantly, monitoring the absorption spectral change before and after irradiation provided evidence of the formation of the reduced intermediates of Co-

(dmgH)₂Cl₂. Before irradiation, the absorption spectrum was simply the sum of the individual components and eosin Y showed the absorption in the visible light region. Upon irradiation by green LEDs for 2 min, a characteristic absorption appeared at 450 nm consistent with the formation of Co^{II} species as reported in the literature (Figure 1c). Prolonged irradiation within 4 min led to new lower energy absorptions at 550–650 nm, in addition to the one at 450 nm (Figure 1d). The new lower energy bands are identical to Co^I species reported in the literature, which is a crucial intermediate for H₂ evolution.¹⁰

On the basis of the above observations, we proposed a plausible mechanism for the CCHE reaction (Scheme 2). Upon


Scheme 2. Proposed Mechanism



visible light irradiation, eosin Y is immediately excited to its singlet excited state ¹[eosin Y]* followed by a quick intersystem crossing (ISC) to its triplet excited state ³[eosin Y]* with a lifetime of 98.8 μs (SI, Figure S3a). Subsequent electron transfer from **1a** to ³[eosin Y]* results in the formation of the eosin Y radical anion, [eosin Y]^{•-}, and amine radical cation **1a**^{•+}; the electron transfer rate constant (*k*₁) from **1a** to the excited ³[eosin Y]* was estimated to be 3.76 × 10⁹ M⁻¹ s⁻¹ according to the lifetime of ³[eosin Y]* with (0.8 μs) and without (98.8 μs) **1a** (SI, Figure S3b). The generated amine radical cation **1a**^{•+} releases a proton and is further oxidized to generate an iminium intermediate, which can effectively react with nucleophiles to afford the desired cross-coupling product. The formed eosin Y radical anion ([eosin Y]^{•-}), on the other hand, delivers an electron to Co(dmgh)₂Cl₂ to regenerate eosin Y to its ground state. Based on the lifetime alternation of [eosin Y]^{•-} from 39.8 to 4.3 μs (SI, Figure S3c, d), the rate of electron transfer (*k*₂) from [eosin Y]^{•-} to Co(dmgh)₂Cl₂ was determined to be 7.78 × 10⁹ M⁻¹ s⁻¹. The cascade electron transfer from substrate **1a** to the excited ³[eosin Y]* to produce [eosin Y]^{•-}, and then from the reduced radical anion [eosin Y]^{•-} to Co(dmgh)₂Cl₂, is believed to be responsible for the efficient CCHE reaction in the homogeneous solution. It is suggested that the formation of H₂ is protonation of Co^I to form a Co^{III}-H hydrid.¹⁰ H₂ is generated by either protonation of the Co^{III}-H hydrid, H₂ elimination and reduction of Co^{III} to Co^{II}, or reduction of the Co^{III}-H hydrid to the Co^{II}-H hydrid followed by protonation to give H₂ and Co^{II}. The alternative involves a reaction between two Co^{III}-H hydrids to form H₂ and 2 molecules of Co^{II}.

Under the optimized conditions, various indole derivatives were examined to show the generality of the CCHE reaction in a homogeneous solution (Scheme 3). When the electron-withdrawing or -donating groups were introduced at different positions of indoles, the desired cross-coupling product with concomitant emission of H₂ could also be obtained in good to excellent yields. In contrast to the reported heterogeneous


Scheme 3. Scope of Indoles^a



^aReaction conditions: 0.1 mmol of **1a**, 0.2 mmol of **2**, 4 mL of H₂O, 1 mL of CH₃CN, 2.04 mg of eosin Y, 2.9 mg of Co(dmgh)₂Cl₂; the solution is strictly deaerated, with pure methane as the internal standard, and irradiated for 16–22 h by 525 nm green LEDs. ^bYields detected by ¹H NMR using 4-nitroacetophenone as an internal standard based on **1a**; isolated yields are given in parentheses. ^cCorresponding to **1a**.

system, the yield of the cross-coupling product and H₂ is greatly enhanced in the homogeneous solution when the substituted group is Br or ester in the indole skeleton. When *N*-phenyl-tetrahydroisoquinolines with various substituted groups were examined, good to excellent yields were also obtained (Scheme 4). As tabulated in Scheme 4, the steric hindrance seems not to have a big impact on the final results, while the yield of the cross-coupling product and H₂ decreased when R₆ was F, Cl, or Br, respectively.

In summary, we have developed a noble-metal-free homogeneous strategy for the CCHE reaction with no use of any sacrificial oxidants under visible light irradiation. Using low-cost, earth-abundant eosin Y and Co(dmgh)₂Cl₂ as a photosensitizer and a catalyst, good to excellent yields of cross-coupling products and virtual equivalents of H₂ evolution have been achieved without any sacrificial oxidants. The homogeneous CCHE reaction is attractive because it can be performed in a mixture of organic solvent and water, which greatly expands the scope of substrates not soluble in water. Mechanistic insights revealed that the efficient CCHE reaction relies on the cascade electron transfer from substrate **1a** to the excited ³[eosin Y]* to generate the [eosin Y]^{•-} radical anion, which delivers an electron to Co(dmgh)₂Cl₂ for proton reduction to H₂. Importantly, direct evidence of the formation of cation radical **1a**^{•+} was obtained for the first time in the cross-coupling reaction by visible light catalysis and reduced Co^{II} and Co^I species as intermediates, which greatly enriches our knowledge for the development of cleaner, safer, and more atom-economic CCHE reactions.

Scheme 4. Scope of Tetrahydroisoquinolines^a

^aReaction conditions: 0.1 mmol of **1a**, 0.2 mmol of **2**, 4 mL of H₂O, 1 mL of CH₃CN, 2.04 mg of eosinY, 2.9 mg of Co(dmgH)₂Cl₂. The solution is strictly deaerated, with pure methane as the internal standard, and irradiated for 16–22 h by 525 nm green LEDs. ^bYields detected by ¹H NMR using 4-nitroacetophenone as an internal standard based on **1a**; isolated yields are given in parentheses. ^cCorresponding to **1a**.

■ ASSOCIATED CONTENT

Supporting Information

Experimental details and characterization of all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: lwzhu@mail.ipc.ac.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support for this research from the Ministry of Science and Technology of China (2013CB834804, 2014CB239402, and 2013CB834505), the National Natural Science Foundation of China (21390404, 21090343, and 91027041), and the Chinese Academy of Sciences is gratefully acknowledged.

■ REFERENCES

- (1) (a) Young, A. J.; White, M. C. *J. Am. Chem. Soc.* **2008**, *130*, 14090. (b) Chen, K.; Baran, P. S. *Nature* **2009**, *459*, 824. (c) Wang, D.-H.; Engle, K.-M.; Shi, B.-F.; Yu, J.-Q. *Science* **2010**, *327*, 315. (d) Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147. (e) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. *Chem. Rev.* **2011**, *111*, 1293. (f) Liu, C.; Zhang, H.; Shi, W.; Lei, A. *Chem. Rev.* **2011**, *111*, 1780.
- (2) (a) Trost, B. *Science* **1991**, *254*, 1471. (b) Sheldon, R. A. *Chem. Soc. Rev.* **2012**, *41*, 1437. (c) Li, C.-J.; Trost, B. M. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 13197. (d) Anastas, P. T.; Warner, J. C. *Green Chemistry: Theory and Practice*; Oxford University Press: New York, 1998. (e) Newhouse, T.; Baran, P. S.; Hoffmann, R. W. *Chem. Soc. Rev.* **2009**, *38*, 3010.
- (3) (a) Jia, C.; Kitamura, T.; Fujiwara, Y. *Acc. Chem. Res.* **2001**, *34*, 633. (b) Komiya, N.; Terai, H.; Nakae, T.; Murahashi, S.-I. *J. Am. Chem. Soc.* **2003**, *125*, 15312. (c) Stuart, D. R.; Fagnou, K. *Science* **2007**, *316*, 1172. (d) Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2007**, *129*, 11904. (e) Neel, A. J.; Hehn, J. P.; Triplet, P. F.; Toste, F.

D. *J. Am. Chem. Soc.* **2013**, *135*, 14044. (f) Li, C.-J. *Acc. Chem. Res.* **2008**, *42*, 335. (g) Yeung, C. S.; Dong, V. M. *Chem. Rev.* **2011**, *111*, 1215. (h) Girard, S. A.; Knauber, T.; Li, C.-J. *Angew. Chem., Int. Ed.* **2014**, *53*, 74.

(4) Meng, Q.-Y.; Zhong, J.-J.; Liu, Q.; Gao, X.-W.; Zhang, H.-H.; Lei, T.; Li, Z.-J.; Feng, K.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *J. Am. Chem. Soc.* **2013**, *135*, 19052.

(5) Meng, Q.-Y.; Liu, Q.; Zhong, J.-J.; Zhang, H.-H.; Li, Z.-J.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *Org. Lett.* **2012**, *14*, 5992.

(6) (a) Zeitler, K. *Angew. Chem., Int. Ed.* **2009**, *48*, 9785. (b) Yoon, T. P.; Ischay, M. A.; Du, J. *Nat. Chem.* **2010**, *2*, 527. (c) Narayanam, J. M. R.; Stephenson, C. R. J. *Chem. Soc. Rev.* **2011**, *40*, 102. (d) Xuan, J.; Xiao, W.-J. *Angew. Chem., Int. Ed.* **2012**, *51*, 6828. (e) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322.

(7) (a) Condie, A. G.; González-Gómez, J. C.; Stephenson, C. R. J. *J. Am. Chem. Soc.* **2010**, *132*, 1464. (b) Hari, D. P.; König, B. *Org. Lett.* **2011**, *13*, 3852. (c) Rueping, M.; Vila, C.; Koenigs, R. M.; Poschorny, K.; Fabry, D. C. *Chem. Commun.* **2011**, *47*, 2360. (d) Su, F.; Mathew, S. C.; Möhlmann, L.; Antonietti, M.; Wang, X.; Blechert, S. *Angew. Chem., Int. Ed.* **2011**, *50*, 657. (e) To, W.-P.; Tong, G. S.-M.; Lu, W.; Ma, C.; Liu, J.; Chow, A. L.-F.; Che, C.-M. *Angew. Chem., Int. Ed.* **2012**, *51*, 2654. (f) Liu, Q.; Li, Y.-N.; Zhang, H.-H.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *Chem.—Eur. J.* **2012**, *18*, 620. (g) Zhong, J.-J.; Meng, Q.-Y.; Wang, G.-X.; Liu, Q.; Chen, B.; Feng, K.; Tung, C.-H.; Wu, L.-Z. *Chem.—Eur. J.* **2013**, *19*, 6443. (h) Gao, X.-W.; Meng, Q.-Y.; Xiang, M.; Chen, B.; Feng, K.; Tung, C.-H.; Wu, L.-Z. *Adv. Synth. Catal.* **2013**, *355*, 2158. (i) Kreis, L. M.; Krautwald, S.; Pfeiffer, N.; Martin, R. E.; Carreira, E. M. *Org. Lett.* **2013**, *15*, 1634. (j) Weiss, M. E.; Kreis, L. M.; Lauber, A.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2011**, *50*, 11125.

(8) (a) Li, Z. P.; Li, C.-J. *J. Am. Chem. Soc.* **2005**, *127*, 6968. (b) Liu, P.; Zhou, C.-Y.; Xiang, S.; Che, C.-M. *Chem. Commun.* **2010**, *46*, 2739. (c) Klusmann, M.; Boess, E.; Schmitz, C. J. *Am. Chem. Soc.* **2012**, *134*, 5317.

(9) (a) Joselevich, E.; Willner, I. *J. Phys. Chem.* **1995**, *99*, 6903. (b) Islam, S. D. M.; Konishi, T.; Fujitsuka, M.; Ito, O.; Nakamura, Y.; Usui, Y. *Photochem. Photobiol.* **2000**, *71*, 675.

(10) (a) Dempsey, J. L.; Brunschwig, B. S.; Winkler, J. R.; Gray, H. B. *Acc. Chem. Res.* **2009**, *42*, 1995. (b) Artero, V.; Chavarot-Kerlidou, M.; Fontecave, M. *Angew. Chem., Int. Ed.* **2011**, *50*, 7238. (c) Eisenberg, R.; Lazarides, T.; McCormick, T.; Du, P.; Luo, G.; Lindley, B. J. *Am. Chem. Soc.* **2009**, *131*, 9192.