

# Shagenes A and B, New Tricyclic Sesquiterpenes Produced by an Undescribed Antarctic Octocoral


Jacqueline L. von Salm,<sup>†</sup> Nerida G. Wilson,<sup>‡</sup> Brian A. Vesely,<sup>§</sup> Dennis E. Kyle,<sup>§</sup> Jason Cuce,<sup>†</sup> and Bill J. Baker<sup>\*,†</sup>

<sup>†</sup>Department of Chemistry and Center for Drug Discovery and Innovation and <sup>§</sup>Department of Global Health, University of South Florida, Tampa, Florida 33620, United States

<sup>‡</sup>Western Australia Museum, Perth, Western Australia, Australia

## S Supporting Information


**ABSTRACT:** The isolation and characterization of two new tricyclic sesquiterpenoids, shagenes A (**1**) and B (**2**) are presented. These compounds were isolated from an undescribed soft coral collected from the Scotia Arc in the Southern Ocean. One- and two-dimensional NMR spectroscopy and mass spectrometry provided the data necessary to characterize the compounds and their relative stereochemical configurations. Exploration of the bioactivity of shagenes A and B found **1** active against the visceral leishmaniasis causing parasite, *Leishmania donovani*, with no cytotoxicity against the mammalian host.



Sessile marine organisms such as sponges and corals are well-known for their chemically diverse natural products. It is common for these organisms to lack physical or behavioral defenses, therefore creating a necessity for the production of chemical defenses. From 1985 to 2008, cnidarians were second only to porifera in new marine natural products reported each year.<sup>1</sup> The largest class of cnidarians, Anthozoa, are almost solely responsible for producing the natural products isolated from this phylum with greater than 80% of the compounds identified coming from the subclass Octocorallia.<sup>2</sup> Many of these secondary metabolites have been investigated for their ecological roles in feeding deterrence, allelopathy, and antifouling, providing evidence of widespread bioactivity.<sup>3</sup> This has led many researchers to study their potential in drug discovery; however, the majority of published octocoral chemistry originates in tropical shallow waters with only a few dozen exceptions for deeper cold-water corals (>100 m).<sup>4,5</sup>

Important metabolites for drug discovery that have been identified from soft corals include the potent anticancer diterpene glycosides eleutherobin<sup>6</sup> and sarcodictyin A<sup>7</sup> and the anti-inflammatory pseudopterosins.<sup>8</sup> Terpenoids such as these comprise approximately 92% of all reported octocoral secondary metabolites.<sup>2</sup> Because terpenes are common in plant and algal extracts, it was once thought that the origin of octocoral terpenes was symbiotic microalgae called zooxanthellae.<sup>9</sup> However, more recent results favor soft corals producing their own terpenoid metabolites;<sup>10,11</sup> consider, for example, octocorals from depths of more than 100 m, where light has limited penetration through the water column and photosynthetic microalgal symbionts are absent, which are rich sources of terpenoids.<sup>12</sup> This paper presents the new terpenoid

skeleton of shagene A and B (**1** and **2**)<sup>13</sup> from a previously undescribed genus of octocoral collected via trawling near the South Georgia Islands.<sup>14</sup> Shagenes are reminiscent of tamariscene, from the liverwort *Frullania tamarisci*, which bears the same fused 3/6/5 tricyclic ring system but displays a different methylation pattern on that ring system.<sup>15</sup>



Other cold-water sesquiterpenoids isolated from corals include the paesslerins,<sup>16</sup> ainigmaptilonones,<sup>17</sup> and alcyopterensins,<sup>18</sup> all of which have shown moderate cytotoxicity toward either human cancer cell lines or microbial pathogens. Shagene A (**1**) demonstrates 5  $\mu$ M half maximal inhibitory concentration (IC<sub>50</sub>) of the infected macrophage stage of the visceral leishmaniasis causing parasite,<sup>19</sup> *Leishmania donovani*, showing a 10-fold increase in activity over the axenic amastigote form (54  $\mu$ M) with no cytotoxicity up to 100  $\mu$ g/mL (345  $\mu$ M) on

Received: March 17, 2014

Published: April 29, 2014

Table 1.  $^1\text{H}^a$  and  $^{13}\text{C}^b$  NMR Data for Shagenes A (1) and B (2) in  $\text{CDCl}_3$ 

position	shagene A (1)			shagene B (2)		
	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (m, J, Hz)	HMBC	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (m, J, Hz)	HMBC
1	33.2	1.19 (1H, m)	2, 8, 9, 14	32.2	1.42 (1H, m)	2, 8, 9, 14
2	35.7	1.25 (1H, q, 11.8)	1, 3, 4, 9, 14	36.4	1.35 (1H, q, 11.8)	1, 3, 4, 9, 14
		1.65 (1H, m)	1, 3, 4, 9, 14		1.68 (1H, m)	1, 3, 4, 9, 14
3	72.2	5.34 (1H, dt, 11.7, 6.4)	2, 4, 10, 16	71.4	5.33 (1H, dt, 11.7, 6.4)	2, 4, 10, 16
4	22.6	1.65 (1H, m)	2, 5, 6, 10, 11	25.5	2.03 (1H, d, 6.1)	2, 3, 5, 6, 9, 10, 11
5	41.9			40.2		
6	147.3			178.0		
7	123.2	5.60 (1H, m)	5, 6, 8, 9, 15	129.7	5.94 (1H, m)	5, 6, 8, 9, 15
8	88.6	4.09 (1H, d, 0.6)	1, 5, 6, 7, 18	208.5		
9	48.4	1.48 (1H, dd, 1.5, 9)	1, 2, 5, 8, 10, 14	49.0	1.89 (1H, d, 10.9)	1, 2, 5, 6, 8, 10, 14
10	31.5	1.78 (1H, br s)	3, 4, 5, 6, 9	33.1	2.06 (1H, br s)	4, 5, 6, 9, 11, 12
11	140.6			139.4		
12	115.9	5.05 (1H, q, 1.6)	10, 13	116.9	5.11 (1H, q, 1.6)	10, 13
		5.20 (1H, br s)	10, 11, 13		5.20 (1H, br s)	10, 11, 13
13	24.1	1.77 (3H, s)	10, 11, 12	24.1	1.76 (3H, s)	10, 11, 12
14	19.5	1.01 (3H, d, 6.4)	1, 2, 3, 9	19.3	1.23 (3H, d, 6.1)	1, 2, 3, 9
15	11.6	1.49 (3H, t, 1.2)	5, 6, 7	13.2	1.84 (3H, d, 1.2)	5, 6, 7
16	170.9			170.8		
17	21.6	2.09 (3H, s)	3, 16	21.5	2.11 (3H, s)	3, 16
18	55.6	3.34 (3H, s)	8			

$^a$  $^1\text{H}$  NMR experiments at 500 MHz, recorded in ppm (integration, multiplicity,  $J$ -coupling in Hz);  $^b$  $^{13}\text{C}$  NMR experiments at 125 MHz, recorded in ppm.

J774.A1 macrophages.<sup>20</sup> The selectivity index (SI) of shagene A (1) is therefore greater than 70 as compared to the lower SI of the control drug miltefosine. Miltefosine shows an  $\text{IC}_{50}$  of 3  $\mu\text{M}$  against the axenic amastigote form, 1.4  $\mu\text{M}$  for the infected macrophage stage, but greater than 50  $\mu\text{M}$  cytotoxicity against mammalian cells. Shagene B (2) showed no activity (>20  $\mu\text{g}/\text{mL}$ ) alluding to the importance of having the methoxy substituent at C-8.

The bright yellow and highly branched octocoral was collected during austral spring 2011 at depths of approximately 180 m via trawling. The location sampled was the sesquiterpenes namesake, Shag Rocks, which are a series of small islands along the Scotia Arc (S53° 26' 16.10", W42° 02' 15.52"). Specimens were frozen immediately due to noticeable oxidation of the yellow pigment. Once freeze-dried, the coral was processed with a Soxhlet extractor with refluxing  $\text{CH}_2\text{Cl}_2$ . The crude extract was mounted on silica gel and subsequently fractionated by normal-phase MPLC. Further purification of the nonpolar portion eluting in EtOAc-*n*-hexane (1:1) was accomplished by HPLC equipped with UV and ELSD detection. The more lipophilic colorless oil shagene A (1) (27 mg, 0.009% dry wt) was collected as the major component of the mixture with shagene B (2) (2 mg colorless oil, 0.0007% dry wt) considerably less abundant.

The  $^1\text{H}$  NMR spectra of shagenes A (1),  $[\alpha]_{\text{D}}^{26} -0.9$ , and B (2),  $[\alpha]_{\text{D}}^{26} +0.2$ , displayed characteristic peaks indicative of smaller cyclic terpenes with 16 proton signals largely composed of single proton integrations, multiple downfield olefinic protons, and methyl substituents. The  $^{13}\text{C}$  NMR spectrum of 1 showed 18 carbon signals including five downfield signals of four olefinic carbons and the carbonyl carbon of the acetyl group ( $\delta_{\text{C}}$  170.9). As represented in Table 1, the spectra display functionality indicative of one acetate methyl ( $\delta_{\text{H}}$  2.09), one methoxy group ( $\delta_{\text{H}}$  3.34), one olefinic methylene ( $\delta_{\text{H}}$  5.05, 5.20), one trisubstituted olefin ( $\delta_{\text{H}}$  5.60), three vinyl or aliphatic methyl groups, six  $\text{sp}^3$  methines, one  $\text{sp}^3$  methylene,

and two olefinic quaternary carbons. HREIMS of 1 provided the molecular formula  $\text{C}_{18}\text{H}_{26}\text{O}_3$  ( $m/z$  calcd 290.1882, found  $m/z$  290.1878) demonstrating six degrees of unsaturation and providing validation for the hypothesized 15 carbon skeleton of a tricyclic sesquiterpene bearing a quaternary carbon at the junction of all three rings. Further verification of this scaffold and relative stereochemistry were accomplished via 2D NMR experiments (COSY, gHMQC, gHMBCAD, and ROESY).

One-bond correlations from the HMQC experiment for 1 confirmed the olefinic methylene protons at C-12 ( $\delta_{\text{C}}$  115.9) and the quaternary nature of carbons C-5 and C-11 ( $\delta_{\text{C}}$  41.9, 140.6). Shared HMBC correlations (Figure 1) of the two




Figure 1. Key COSY and HMBC correlations of 1.

protons at C-12 and  $\text{H}_3$ -13 ( $\delta_{\text{H}}$  5.05, 5.20, and 1.77) with C-10 ( $\delta_{\text{C}}$  31.5) and C-11 provided proof of the isopropenyl attachment to the cyclopropane at C-10. Placement of C-5 as a bridgehead carbon for this cyclopropane, the cyclohexane and cyclopentene fused rings, is verified by HMBC correlations with H-4, H-7, H-8, H-9, H-10, and  $\text{H}_3$ -15. Chemical shifts for C-4 ( $\delta_{\text{C}}$  22.6), C-5, and C-10 of shagene A are similar to those reported for the cyclopropane ring of the tricyclic skeleton comprising tamariscene ( $\delta_{\text{C}}$  30.1,  $\delta_{\text{C}}$  34.2,  $\delta_{\text{C}}$  33.6)<sup>15</sup> with the

added functionality of shagene A accounting for shift variance. Connectivity of the cyclopentene moiety can be completed on the basis of COSY coupling of the olefinic proton H-7 ( $\delta_{\text{H}}$  5.60) with vinyl methyl protons H<sub>3</sub>-15 ( $\delta_{\text{H}}$  1.49) and the oxygen-bearing methine H-8 ( $\delta_{\text{H}}$  4.09). HMBC coupling of the methoxy protons with the oxygen bearing methine at C-8 verifies their connectivity. Further analysis of the 2D COSY spectrum shows H-1 ( $\delta_{\text{H}}$  1.19) to be vicinal to H-9 ( $\delta_{\text{H}}$  1.48), the methyl protons at H<sub>3</sub>-14 ( $\delta_{\text{H}}$  1.01), and the diastereotopic methylene protons at H-2 ( $\delta_{\text{H}}$  1.25,  $\delta_{\text{H}}$  1.65). The acetate methyl protons provided HMBC correlations to the carbonyl C-16 ( $\delta_{\text{C}}$  170.9) and C-3 ( $\delta_{\text{C}}$  72.2) of the cyclohexane ring. COSY coupling of H-3 ( $\delta_{\text{H}}$  5.34) with neighboring methylene protons H-2 and methine H-4 ( $\delta_{\text{H}}$  1.65) confirms a large portion of the cyclohexane ring. HMBC correlations of H-4 with C-2, C-5, C-6, C-10, and C-11 provide the evidence of a third bridgehead carbon, this time between the cyclohexane and cyclopropane rings to complete the connectivity of the overall scaffold.

Relative stereochemistry of shagene A (**1**) was deducible upon analysis of strong, unanticipated ROESY correlations (Figure 2) between the exocyclic methylene protons on C-12




Figure 2. Key ROESY correlations for shagene A (**1**).


with the cyclopentane and cyclohexane bridgehead H-9, a methylene proton at C-2, and the methyl protons of the acetate. This places the isopropenyl group in the periphery of the acetate and methyl substituents of the cyclohexane ring essentially blocking that face of the molecule. Spatial placement of the methyl substituent at C-1 is a result of correlations between H<sub>3</sub>-14 with H-9, and ROESY correlations between H-1 and H-8 place the methoxy group on the same face as H-9. Additional support for the  $\beta$ -methoxy is provided by conformational analysis of the H-9 coupling constants. Energy-minimized conformation of the  $\alpha$ -methoxy configuration requires a large  $^3J_{\text{HH}}$  coupling between H-8 and H-9 due to a nearly eclipsed conformation. On the other hand, the  $\beta$ -methoxy displays a roughly  $118^\circ$  dihedral angle resulting in a small  $J_{\text{H8,H9}}$ . The observed  $J_{\text{H8,H9}}$  of 1.5 Hz indicates a  $\beta$ -methoxy substituent at C-8. H<sub>3</sub>-18 and H<sub>3</sub>-13 show weak ROESY correlation.

The obvious difference in the  $^1\text{H}$  NMR spectra between compounds **1** and **2** was the absence of the proton signals associated with the methoxy functionality at  $\delta_{\text{H}}$  3.34 ppm and its oxymethine at  $\delta_{\text{H}}$  4.09 ppm as well as the downfield shift of multiple proton signals surrounding C-8. A carbon shift in **2**, but absent in **1** was observed at  $\delta_{\text{C}}$  208.5, establishing the oxidized  $\alpha,\beta$ -unsaturated ketone moiety of **2**. The presence of a carbonyl at C-8 was further verified by the absence of H-8 COSY correlations with H-9 and H-7. Evidence of a conjugated ketone was indicated by absorbance at a wavelength of 257 nm in the UV spectrum and the extra carbonyl stretch on IR ( $1696\text{ cm}^{-1}$ ). HREIMS of **2** revealed an expected mass similar to **1**

save one carbon and four protons ( $m/z$  calcd 274.1569, found 274.1569) providing the molecular formula  $\text{C}_{17}\text{H}_{22}\text{O}_3$ . Validation of **2** having the same connectivity and stereochemistry as **1** is shown in the 2D NMR experiments (COSY, HSQC, HMBC, and ROESY).

The proposed biosynthetic pathway to **1** and **2** is shown in Scheme 1. Structural similarities to biosynthetic intermediates

#### Scheme 1. Proposed Biosynthetic Pathway for Sesquiterpenes **1** and **2**



displayed in the biosynthesis of valerenic acid makes these molecules likely to undergo cyclization through a caryophyllene intermediate (**3**) via copalyl diphosphate synthase (CPS).<sup>21</sup> Alternative pathways have been suggested with a concerted three-ring cyclization step; however, this would produce an undesirable stereochemical outcome.

### ■ ASSOCIATED CONTENT

#### Supporting Information

$^1\text{H}$ ,  $^{13}\text{C}$ , COSY, HSQC, HMQC, HMBC, and ROESY NMR spectra for structure elucidation of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

### ■ AUTHOR INFORMATION

#### Corresponding Author

\*E-mail: [bjbaker@usf.edu](mailto:bjbaker@usf.edu).

#### Notes

The authors declare no competing financial interest.

### ■ ACKNOWLEDGMENTS

We thank the crew and research scientists on board the Nathaniel B. Palmer during the above-mentioned field season that kept us afloat and were always willing to lend a helping hand. The project was supported by National Science Foundation awards ANT-1043749 (N.G.W.) and ANT-0838776 (B.J.B.) from the Antarctic Organisms and Ecosystems program, and National Institutes of Health grant AI103673 (B.J.B. and D.E.K.) from the National Institute of Allergy and Infectious Diseases. Greg W. Rouse, Scripps Institution of Oceanography, was instrumental in the logistical and conceptual implementation of this research. We extend special thanks to family and friends of The Baker Lab for their support, hard work, and dedication.

## ■ REFERENCES

- (1) Hu, G. P.; Yuan, J.; Sun, L.; She, Z. G.; Wu, J. H.; Lan, X. J.; Zhu, X.; Lin, Y. C.; Chen, S. P. *Mar. Drugs* **2011**, *9*, 514–525.
- (2) Harper, M. K.; Bugni, T. S.; Copp, B. R.; James, R. D.; Lindsay, B. S.; Richardson, A. D.; Schnabel, P. C.; Tasdemir, D.; VanWagoner, R. M.; Verbitski, S. M.; Ireland, C. M. Introduction to the Chemical Ecology of Marine Natural Products. In *Marine Chemical Ecology*; McClintock, J. B., Baker, B. J., Eds.; CRC Press: Boca Raton, FL, 2001; pp 3–70.
- (3) Coll, J. C. *Chem. Rev.* **1992**, *92*, 613–631.
- (4) Skropeta, D. *Nat. Prod. Rep.* **2008**, *25*, 1131–1166.
- (5) Lebar, M. D.; Heimbegner, J. L.; Baker, B. J. *Nat. Prod. Rep.* **2007**, *24*, 774–797.
- (6) Lindel, T.; Jensen, P. R.; Fenical, W.; Long, B. H.; Casazza, A. M.; Carboni, J.; Fairchild, C. R. *J. Am. Chem. Soc.* **1997**, *119*, 8744–8745.
- (7) Dambrosio, M.; Guerriero, A.; Pietra, F. *Helv. Chim. Acta* **1987**, *70*, 2019–2027.
- (8) Look, S. A.; Fenical, W.; Jacobs, R. S.; Clardy, J. *Proc. Natl. Acad. Sci. U.S.A.* **1986**, *83*, 6238–6240.
- (9) Coll, J. C.; Bowden, B. F.; Tapiolas, D. M.; Willis, R. H.; Djura, P.; Streamer, M.; Trott, L. *Tetrahedron* **1983**, *41*, 1085–1092.
- (10) Bandurraga, M. M.; McKittrick, B.; Fenical, W.; Arnold, E.; Clardy, J. *Tetrahedron* **1982**, *38*, 305–310.
- (11) Michalek-Wagner, K.; Bourne, D. J.; Bowden, B. F. *Mar. Biol.* **2001**, *138*, 753–760.
- (12) Scheuer, P. J. *Science* **1990**, *248*, 173–177.
- (13) Shagene A (1): UV ( $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  ( $\epsilon$ ) 240 (5945) nm; IR (thin film) 2959, 3050, 2933, 2877, 1734, 1655, 1458, 1369, 1246, 1089, 1026  $\text{cm}^{-1}$ ;  $^1\text{H}$  and  $^{13}\text{C}$  NMR data, see Table 1; 70 eV HREIMS  $m/z$  290.1878 ( $[\text{M}^+]$ ) ( $\text{C}_{18}\text{H}_{26}\text{O}_3$  calcd, 290.1882). Shagene B (2): UV ( $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  ( $\epsilon$ ) 257 (5425), 240 (6357) nm; IR (thin film) 3080, 2960, 2929, 2858, 1734, 1696, 1607, 1462, 1380, 1242, 1186, 1030  $\text{cm}^{-1}$ ;  $^1\text{H}$  and  $^{13}\text{C}$  NMR data, see Table 1; 70 eV HREIMS  $m/z$  274.1569 ( $[\text{M}^+]$ ) ( $\text{C}_{17}\text{H}_{22}\text{O}_3$  calcd, 274.1569).
- (14) This coral was assumed to be *Paragorgia* sp. upon collection; however, the bright yellow coloration and DNA sequence data from the mismatch repair (mtMutS) gene suggests this species is not related to any known species, genera, or families of Octocorallia. Further genetic analysis of the specimen is necessary for further resolution. A sample voucher has been accessioned in the Benthic Invertebrate Collection, Scripps Institution of Oceanography (S3528).
- (15) Paul, C.; Konig, W. A.; Muhle, H. *Phytochemistry* **2001**, *57*, 307–313.
- (16) Rodriguez Brasco, M. F.; Seldes, A. M.; Palermo, J. A. *Org. Lett.* **2001**, *3*, 1415–1417.
- (17) Iken, K. B.; Baker, B. J. *J. Nat. Prod.* **2003**, *66*, 888–890.
- (18) Palermo, J. A.; Rodriguez Brasco, M. F.; Spagnuolo, C.; Seldes, A. M. *J. Org. Chem.* **2000**, *65*, 4482–4486.
- (19) Siqueira-Neto, J. L.; Moon, S.; Jang, J.; Yang, G.; Lee, C.; Moon, H. K.; Chatelain, E.; Genovesio, A.; Cecchetto, J.; Freitas-Junior, L. H. *PLoS Negl. Trop. Dis* **2012**, *6*, e1671.
- (20) The axenic amastigote bioassay for leishmaniasis was an MTS [3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulphophenyl)-2H-tetrazolium] 72 h assay using 66000 amastigotes per well.  $\text{IC}_{50}$  values were calculated from sigmoidal inhibition curves using SigmaPlot 11.0 software. The procedure for the J447.A-1 cytotoxicity assay is the same as the axenic amastigote bioassay with one exception, 50000 amastigotes are used per well.
- (21) Yeo, Y. S.; Nybo, S. E.; Chittiboyina, A. G.; Weerasooriya, A. D.; Wang, Y. H.; Gongora-Castillo, E.; Vaillancourt, B.; Buell, C. R.; DellaPenna, D.; Celiz, M. D.; Jones, A. D.; Wurtele, E. S.; Ransom, N.; Dudareva, N.; Shaaban, K. A.; Tibrewal, N.; Chandra, S.; Smillie, T.; Khan, I. A.; Coates, R. M.; Watt, D. S.; Chappell, J. *J. Biol. Chem.* **2013**, *288*, 3163–3173.