


Stereocontrolled Construction of Six Vicinal Stereogenic Centers on Spiropyrazolones via Organocascade Michael/Michael/1,2-Addition Reactions

Pankaj Chauhan, Suruchi Mahajan, Charles C. J. Loh, Gerhard Raabe, and Dieter Enders*

Institute of Organic Chemistry, RWTH Aachen University Landoltweg 1, 52074 Aachen, Germany

S Supporting Information

ABSTRACT: A highly stereoselective one-pot procedure for the synthesis of spiropyrazolone derivatives bearing six contiguous stereogenic centers including two tetrasubstituted carbons has been developed. Under sequential catalysis by two organocatalysts, a cinchona-derived aminosquaramide and DBU, a series of diversely functionalized spiropyrazolones are obtained in good yields (47–62%) and excellent stereoselectivities (up to >25:1 dr and 98–99% ee). The opposite enantiomers of the spiropyrazolones are also accessible by employing a pseudoenantiomeric aminosquaramide catalyst.



The synthesis of pyrazolone derivatives has attracted considerable attention in recent years due to their wide spectrum of applications in dye, analytical, and pharmaceutical chemistry.^{1,2} Edaravone (**A**), a neuroprotective agent, and metamizole (**B**), an effective analgesic and antipyretic agent, are two important pharmaceutically valuable pyrazolones (Figure 1).^{1c–f} In




Figure 1. Biologically active pyrazolone derivatives.

addition to these, the pyrazolone derivatives such as **C** exhibit HIV inhibition activity.^{1g} Recently, the pyrazolones were found to be inhibitors of the CD80 protein and also possess potent activity in inhibiting protease-resistant prion protein accumulation, cytokines, and p38 kinases.^{1h–k} Pyrazol-3-one derivatives have also been studied as multidrug resistance modulators.^{1l}

On the other hand, spirocyclic frameworks frequently found in many synthetic bioactive compounds and natural products fascinated many researchers to develop synthetic strategies for their construction. Because of their complex three-dimensional structure, the stereoselective synthesis of spirocyclic molecules from simple precursors in an atom-economic fashion is considered as a tough challenge.³ Recently, the organocatalytic cascade/domino reactions emerged as an efficient strategy to provide such complex structures in an operationally simple one-pot procedure,⁴ and we have witnessed tremendous growth in

the asymmetric synthesis of spirocyclic oxindole derivatives especially employing organocascade sequences.⁵ Despite the many important applications of the pyrazolone moiety, the catalytic asymmetric synthesis of spiropyrazolones is less explored.

The Rios group reported the asymmetric synthesis of spiropyrazolones bearing three contiguous stereogenic centers via a secondary amine catalyzed domino reaction (Scheme 1).⁶ Rios and co-workers were also able to construct four stereocenters on a spiropyrazolones in a three-component domino reaction.⁷ The cinchona-derived primary amines were found to catalyze the domino Michael/Michael reaction of the enones with the pyrazolones⁸ and the unsaturated pyrazolones⁹ to afford the spirocyclohexanonepyrazolones with three consecutive stereogenic centers. Recently, spiropyrazolones bearing an *N*-heterocyclic ring with three adjacent stereogenic centers were synthesized by aminothiureas¹⁰ and quinine¹¹ catalyzed domino Michael/cyclization reactions.


To the best of our knowledge, the asymmetric synthesis of spirocyclohexanonepyrazolones bearing more than four stereocenters are not known. We took up this challenge and developed a one-pot protocol for the asymmetric synthesis of the spiropyrazolones bearing as many as six stereogenic centers including two adjacent tetrasubstituted ones. It was envisaged that an organocascade sequence involving a stereoselective Michael/Michael/1–2-addition reaction between β -dicarbonyl compounds **1**, nitroalkenes **2**, and unsaturated pyrazolones **3** mediated by sequential organocatalysis¹² using a chiral bifunctional organocatalyst¹³ and an achiral base could afford such highly functionalized spiropyrazolones (Scheme 1).

To attain our objective at the onset, we performed a one-pot reaction that involved a quinine-derived squaramide (1 mol %)

Received: April 15, 2014

Published: May 19, 2014

Scheme 1. Catalytic Enantioselective Strategies for the Synthesis of Spiropyrzalone Derivatives



catalyzed Michael addition of ethyl acetoacetate (**1a**) to (*E*)- β -nitrostyrene (**2a**) in dichloromethane. After 24 h, the unsaturated pyrazolone **3a** and a guanidine base TBD (20 mol %) in dichloromethane were added sequentially. To our delight, the desired spiropyrzalone bearing four tertiary and two tetrasubstituted stereogenic centers was obtained in a good yield of 41% and excellent enantio- (99% ee) and diastereoselectivity (>25:1 dr) (Table 1, entry 1). The screening of other achiral bases (entries 2–6) showed that the DBU provides the best yield as well as excellent stereoselectivity (entry 2). The screening of solvents such as chloroform, toluene, and THF did not result in any improvement in the product yield (entries 7–9). Further efforts for optimizing the yield of **4a** by increasing the catalyst loading of DBU revealed that with 50 mol % DBU a better yield was observed; however, a further increase in the amount of DBU did not show any improvement in the product yield. Using 2 equiv of **3** the spiropyrzalone **4** was isolated in maximum yield of 60% with excellent stereoselectivity (entries 13 and 14).

Armed with optimized conditions, we further evaluated the substrate scope on a 0.5 mmol scale (Table 2). Various pyrazolone-derived olefins bearing electron-withdrawing and electron-releasing substituents at the different aromatic position reacted efficiently under the standard reaction conditions to afford the desired spiropyrzalones **4b–i** in good yields and excellent enantio- and diastereoselectivities (entries 2–9). The heteroaromatic pyrazolone derivative was also tolerated under this one-pot organocascade procedure, thus resulting in the desired spiropyrzalone **4j** in 55% yield with excellent ee (entry 10). The unsaturated pyrazolone bearing an *o*-chlorophenyl group gives 47% yield of product **4k** with 99% ee (entry 11).

Different aromatic nitroalkenes bearing electron-withdrawing as well as electron-donating substituents also allowed an efficient access to the spiropyrzalones **4l–o** in good yield and excellent ee of 98–99% (entries 12–15). The nitroalkenes

Table 1. Optimization of the Reaction Conditions for the Asymmetric Synthesis of the Spiropyrzalone **4a**^a

entry	base (x mol %)	solvent	yield ^b (%)	ee ^c (%)
1	TBD (20)	CH ₂ Cl ₂	41	99
2	DBU (20)	CH ₂ Cl ₂	44	99
3	DBN (20)	CH ₂ Cl ₂	43	99
4	DABCO (20)	CH ₂ Cl ₂	12	99
5 ^d	piperidine (20)	CH ₂ Cl ₂	14	98
6 ^d	pyrrolidine (20)	CH ₂ Cl ₂	12	98
7	DBU (20)	CHCl ₃	35	99
8	DBU (20)	toluene	34	99
9	DBU (20)	THF	35	99
10	DBU (30)	CH ₂ Cl ₂	46	99
11	DBU (50)	CH ₂ Cl ₂	52	99
12 ^e	DBU (100)	CH ₂ Cl ₂	36	99
13 ^f	DBU (30)	CH ₂ Cl ₂	57	99
14 ^f	DBU (50)	CH ₂ Cl ₂	60	99

^aReaction conditions: 0.2 mmol of **1a**, 0.2 mmol of **2a**, 1 mol % of **I**, 0.24 mmol of **3a**, and x mol % of base (0.1 M in solvent). ^bYield of isolated **4a** after flash column chromatography. ^cEnantiomeric excess of the major diastereomer (>25:1 dr) determined by HPLC analysis on a chiral stationary phase. ^dThe reaction was run for 96 h in the second step. ^eThe reaction was run for 0.5 h in the second step. ^f0.40 mmol of **3a** was used.


bearing heteroaromatic groups also provide the desired adducts **4p** and **4q** in good yields and excellent stereoselectivities (entries 16 and 17). We also tried other dicarbonyl compounds such as methyl acetoacetate and acetylacetone, which efficiently reacted under this one-pot protocol to afford the adducts **4r** and **4s** in 57% and 53% yield and 99% ee (entries 18 and 19).

After generating two tetrasubstituted adjacent carbon atoms on the spiropyrzalone, we tried to create three consecutive tetrasubstituted carbons by employing a trisubstituted β -ketoester. A one-pot organocatalytic sequence promoted by **I** and DBU between the β -ketoester **5**, (*E*)- β -nitrostyrene (**2a**), and pyrazolone **3a** provided the spiropyrzalone **6** bearing three contiguous tertiary and three tetrasubstituted stereogenic centers in excellent stereoselectivity (>25:1 dr and 95% ee), albeit in a low yield of 16% (Scheme 2).

The absolute configuration of the spiropyrzalones **4a–s** could be assigned as (1*R*), (2*S*), (3*S*), (4*S*), (5*S*), and (6*R*) based on the X-ray structure of **4c**.¹⁴

We have also tested the catalytic potential of amino-squaramide catalyst **II** derived from quinidine in place of catalyst **I** to generate the opposite enantiomer of the corresponding product. With catalyst **II**, various spiropyrzalone derivatives were accessible in good yields and excellent enantio- and diastereoselectivities (Table 3).


In order to demonstrate the practical utility of this one-pot protocol, we have performed a gram-scale cascade

Table 2. Substrate Scope of the Asymmetric Synthesis of Spiropyrzazolones **4** with Catalyst **I**^a


entry	R ¹	R ²	R ³	R ⁴	4	yield ^b (%)	ee ^c (%)
1	OEt	Ph	Ph	Ph	4a	62	99
2	OEt	Ph	4-ClC ₆ H ₄	Ph	4b	58	99
3	OEt	Ph	4-BrC ₆ H ₄	Ph	4c	62	99
4	OEt	Ph	4-CF ₃ C ₆ H ₄	Ph	4d	61	99
5	OEt	Ph	3-ClC ₆ H ₄	Ph	4e	60	99
6	OEt	Ph	4-MeC ₆ H ₄	Ph	4f	52	99
7	OEt	Ph	4-MeOC ₆ H ₄	Ph	4g	57	99
8	OEt	Ph	2-MeC ₆ H ₄	Ph	4h	55	99
9	OEt	Ph	3-MeOC ₆ H ₄	Ph	4i	57	99
10	OEt	Ph	2-thienyl	Ph	4j	55	99
11	OEt	Ph	Ph	2-ClC ₆ H ₄	4k	47	99
12	OEt	4-FC ₆ H ₄	Ph	Ph	4l	57	99
13	OEt	4-ClC ₆ H ₄	Ph	Ph	4m	58	99
14	OEt	4-MeC ₆ H ₄	Ph	Ph	4n	56	98
15	OEt	4-MeOC ₆ H ₄	Ph	Ph	4o	56	99
16 ^d	OEt	2-furanyl	Ph	Ph	4p	55	99
17	OEt	2-thienyl	Ph	Ph	4q	59	99
18	OMe	Ph	Ph	Ph	4r	57	99
19	Me	Ph	Ph	Ph	4s	53	99


^aReaction conditions: 0.5 mmol of **1**, 0.5 mmol of **2**, 1 mol % of **I**, 1.0 mmol of **3**, and 50 mol % of DBU (0.1 M in CH₂Cl₂). ^bYield of the isolated product after flash column chromatography. ^cEnantiomeric excess of the major diastereomer (>25:1 dr). ^d12:1 dr.

Scheme 2. Asymmetric Synthesis of Spiropyrzalone with Three Contiguous Tetrasubstituted Stereocenters



Michael/Michael/1,2-addition reaction between **1a**, **2a**, and **3a** using a lower loading (0.5 mol %) of catalyst **I** (Scheme 3). The desired spiropyrazolone **4a** was obtained in 59% yield without deteriorating the stereochemical outcome of the reaction.


In conclusion, we have disclosed a one-pot procedure for the synthesis of a new series of potentially biologically important spiropyrazolone derivatives. A variety of spirocyclohexanepyrzalone derivatives bearing six stereocenters including two vicinal tetrasubstituted carbons were obtained in good yields and excellent stereoselectivities via sequential organocatalytic Michael/Michael/1,2-addition reactions facilitated by a low loading of a cinchona-derived amino squaramide and a readily available achiral base. This one-pot cascade sequence can be scaled up without losing the reaction efficiency in terms of product yield and stereoselectivity. The opposite enantiomer of the spiropyrazolones can be also synthesized in good yield and excellent stereoselectivity by employing a pseudoenantiomeric catalyst.

Table 3. Substrate Scope of the Asymmetric Synthesis of Spiropyrzazolones *ent*-**4** with Catalyst **II**^a

entry	R ¹	R ²	R ³	<i>ent</i> - 4	yield ^b (%)	ee ^c (%)
1	OEt	Ph	Ph	<i>ent</i> - 4a	61	99
2	OEt	Ph	4-ClC ₆ H ₄	<i>ent</i> - 4b	57	97
3	OEt	Ph	3-ClC ₆ H ₄	<i>ent</i> - 4e	60	98
4	OEt	Ph	4-MeC ₆ H ₄	<i>ent</i> - 4f	50	98
5	OEt	Ph	3-MeOC ₆ H ₄	<i>ent</i> - 4i	57	98
6	OEt	4-FC ₆ H ₄	Ph	<i>ent</i> - 4l	60	97
7	OEt	4-ClC ₆ H ₄	Ph	<i>ent</i> - 4m	57	98
8	OEt	4-MeC ₆ H ₄	Ph	<i>ent</i> - 4n	56	97
9 ^d	OEt	2-thienyl	Ph	<i>ent</i> - 4q	61	98
10	OMe	Ph	Ph	<i>ent</i> - 4r	59	98

^aReaction conditions: 0.5 mmol of **1**, 0.5 mmol of **2**, 1 mol % of **II**, 1.0 mmol of **3**, and 50 mol % of DBU (0.1 M in CH₂Cl₂). ^bYield of the isolated product after flash column chromatography. ^cEnantiomeric excess of the major diastereomer (>25:1 dr). ^d10:1 dr.

Scheme 3. Gram-Scale One-Pot Stereoselective Synthesis of Spiropyrazolone 4a



■ ASSOCIATED CONTENT

Supporting Information

Experimental details and full spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: enders@rwth-aachen.de. Tel: +49 241 809 4676. Fax: +49 241 809 2127.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Support from the European Research Council (ERC Advanced Grant "DOMINOCAT") is gratefully acknowledged.

■ REFERENCES

- (1) For a book on pyrazolone chemistry, see: (a) Varvounis, G. *Pyrazol-3-ones. Part IV: Synthesis and Applications*. In *Advances in Heterocyclic Chemistry*; Katritzky, A. R., Ed.; Academic Press: New York, 2009; Vol. 98, p 143. For reviews, see: (b) Horton, D. A.; Bourne, G. T.; Smythe, M. L. *Chem. Rev.* **2003**, *103*, 893. (c) Watanabe, T.; Tanaka, K.; Watanabe, K.; Takamatsu, Y.; Tobe, A. *Yakugaku Zasshi* **2004**, *124*, 99. (d) Yoshida, H.; Yanai, H.; Namiki, Y.; Fukatsu-Sasaki, K.; Furutani, N.; Tada, N. *CNS Drug Rev.* **2006**, *12*, 9. For selected examples, see: (e) Hinz, B.; Cheremina, O.; Bachmakov, J.; Renner, B.; Zolk, O.; Fromm, M. F.; Brune, K. *FASEB J.* **2007**, *21*, 2343. (f) Wong, A.; Sibbald, A.; Ferrero, F.; Plager, M.; Santolaya, M. E.; Escobar, A. M.; Campos, S.; Barragan, S.; Gonzalez, M. D. L.; Kesselring, G. L. F. *Clin. Pediatr.* **2001**, *40*, 313. (g) Hadi, V.; Koh, Y.-H.; Sanchez, T. W.; Barrios, D.; Neamati, N.; Jung, K. W. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6854. (h) Uvebrant, K.; Thrige, D. D. G.; Rosén, A.; Åkesson, M.; Berg, H.; Walse, B.; Björk, P. J. *Biomol. Screen.* **2007**, *12*, 464. (i) Kimata, A.; Nakagawa, H.; Ohyama, R.; Fukuuchi, T.; Ohta, S.; Suzuki, T.; Miyata, N. *J. Med. Chem.* **2007**, *50*, 5053. (j) Golebiowski, A.; Townes, J. A.; Laufersweiler, M. J.; Brugel, T. A.; Clark, M. P.; Clark, C. M.; Jane, F. D.; Laughlin, S. K.; Sabat, M. P.; Bookland, R. G.; VanRens, J. C.; De, B.; Hsieh, L. C.; Janusz, M. J.; Walter, R. L.; Webster, M. E.; Mekel, M. J. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2285. (k) Clark, M. P.; Laughlin, S. K.; Laufersweiler, M. J.; Bookland, R. G.; Brugel, T. A.; Golebiowski, A.; Sabat, M. P.; Townes, J. A.; VanRens, J. C.; Djung, J. F.; Natchus, M. G.; De, B.; Hsieh, L. C.; Xu, S. C.; Walter, R. L.; Mekel, M. J.; Heitmeyer, S. A.; Brown, K. K.; Juergens, K.; Taiwo, Y. O.; Janusz, M. J. *J. Med. Chem.* **2004**, *47*, 2724. (l) Wang, X. H.; Wang, X. K.; Liang, Y. J.; Shi, Z.; Zhang, J. Y.; Chen, L. M.; Fu, L. W. *Chin. J. Cancer* **2010**, *29*, 980.
- (2) For recent examples of the construction of pyrazolone derivatives, see: (a) Gogoi, S.; Zhao, C.-G. *Tetrahedron Lett.* **2009**, *50*, 2252. (b) Gogoi, S.; Zhao, C.-G.; Ding, D. *Org. Lett.* **2009**, *11*, 2249. (c) Mariappan, G.; Saha, B. P.; Bhuyan, N. R.; Bharti, P. R.; Kumar, D. J. *Adv. Pharm. Technol. Res.* **2010**, *1*, 260. (d) Ma, R.; Zhu, J.; Liu, J.; Chen, L.; Shen, X.; Jiang, H.; Li, J. *Molecules* **2010**, *15*, 3593.
- (e) Liao, Y.-H.; Chen, W.-B.; Wu, Z.-J.; Du, X.-L.; Cun, L.-F.; Zhang, X.-M.; Yuan, W.-C. *Adv. Synth. Catal.* **2010**, *352*, 827. (f) Yang, Z.; Wang, Z.; Bai, S.; Liu, X.; Lin, L.; Feng, X. *Org. Lett.* **2011**, *13*, 596. (g) Mazzanti, A.; Calbet, T.; Font-Bardia, M.; Moyano, A.; Rios, R. *Org. Biomol. Chem.* **2012**, *10*, 1645. (h) Šimek, M.; Remeš, M.; Veselý, J.; Rios, R. *Asian J. Org. Chem.* **2013**, *2*, 64.
- (3) For recent reviews on the stereoselective synthesis of spirocyclic compounds, see: (a) Trost, B. M.; Brennan, M. K. *Synthesis* **2009**, 3003. (b) Rios, R. *Chem. Soc. Rev.* **2012**, *41*, 1060. (c) Nakazakia, A.; Kobayashi, S. *Synlett* **2012**, 1427. (d) Franz, A. K.; Hanhan, N. V.; Ball-Jones, N. R. *ACS Catal.* **2013**, *3*, 540.
- (4) For selected reviews on organocatalytic domino/cascade reactions, see: (a) Enders, D.; Grondal, C.; Hüttl, M. R. M. *Angew. Chem., Int. Ed.* **2007**, *46*, 1570. (b) Yu, X.; Wang, W. *Org. Biomol. Chem.* **2008**, *6*, 2037. (c) Grondal, C.; Jeanty, M.; Enders, D. *Nat. Chem.* **2010**, *2*, 167. (d) Albrecht, L.; Jiang, H.; Jørgensen, K. A. *Angew. Chem., Int. Ed.* **2011**, *50*, 8492. (e) Moyano, A.; Rios, R. *Chem. Rev.* **2011**, *111*, 4703. (f) Grossmann, A.; Enders, D. *Angew. Chem., Int. Ed.* **2012**, *51*, 314. (g) Pellissier, H. *Adv. Synth. Catal.* **2012**, *354*, 237. (h) Loh, C. C. J.; Enders, D. *Chem.—Eur. J.* **2012**, *18*, 10212. (i) Lu, L.-Q.; Chen, J.-R.; Xiao, W.-J. *Acc. Chem. Res.* **2012**, *45*, 1278. (j) Volla, C. M. R.; Atodiresse, I.; Rueping, M. *Chem. Rev.* **2014**, *114*, 2390. For recent highlights, see: (k) Chauhan, P.; Enders, D. *Angew. Chem., Int. Ed.* **2014**, *53*, 1485.
- (5) For recent reviews on the organocatalytic asymmetric synthesis of spirooxindoles, see: (a) Zhou, F.; Liu, Y.-L.; Zhou, J. *Adv. Synth. Catal.* **2010**, *352*, 1381. (b) Badillo, J. J.; Hanhan, N. V.; Franz, A. K. *Curr. Opin. Drug Discovery Dev.* **2010**, *13*, 758. (c) Dalpozzo, R.; Bartoli, G.; Bencivenni, G. *Chem. Soc. Rev.* **2012**, *41*, 7247. (d) Ball-Jones, N. R.; Badillo, J. J.; Franz, A. K. *Org. Biomol. Chem.* **2012**, *10*, S165. (e) Hong, L.; Wang, R. *Adv. Synth. Catal.* **2013**, *355*, 1023. (f) Liu, Y.; Wang, H.; Wan, J. *Asian J. Org. Chem.* **2013**, *2*, 374. (g) Chauhan, P.; Chimni, S. S. *Tetrahedron: Asymmetry* **2013**, *24*, 343. (h) Cheng, D.; Ishihara, Y.; Tan, B.; Barbas III, C. F. *ACS Catal.* **2014**, *4*, 743.
- (6) (a) Companyó, X.; Zea, A.; Alba, A.-N. R.; Mazzanti, A.; Moyano, A.; Rios, R. *Chem. Commun.* **2010**, 46, 6953. (b) Alba, A.-N. R.; Zea, A.; Valero, G.; Calbet, T.; Font-Bardia, M.; Mazzanti, A.; Moyano, A.; Rios, R. *Eur. J. Org. Chem.* **2011**, *7*, 1318.
- (7) Zea, A.; Alba, A.-N. R.; Mazzanti, A.; Moyano, A.; Rios, R. *Org. Biomol. Chem.* **2011**, *9*, 6519.
- (8) Wu, B.; Chen, J.; Li, M.-Q.; Zhang, J.-X.; Xu, X.-P.; Ji, S. J.; Wang, X.-W. *Eur. J. Org. Chem.* **2012**, 1318.
- (9) (a) Liang, J.; Chen, Q.; Liu, L.; Jiang, X.; Wang, R. *Org. Biomol. Chem.* **2013**, *11*, 1441. (b) Zhang, J.-X.; Li, N. K.; Liu, Z.-M.; Huang, X. F.; Geng, Z.-C.; Wang, X.-W. *Adv. Synth. Catal.* **2013**, *355*, 797.
- (10) Liu, L.; Zhong, Y.; Zhang, P.; Jiang, X.; Wang, R. *J. Org. Chem.* **2012**, *77*, 10228. (b) Chen, Q.; Liang, J.; Wang, S.; Wang, D.; Wang, R. *Chem. Commun.* **2013**, 49, 1657.
- (11) Cui, B. D.; Li, S.-W.; Zuo, J.; Wu, Z. J.; Zhang, X.-M.; Yuan, W.-C. *Tetrahedron* **2014**, *70*, 1895.
- (12) For recent examples on organocatalytic sequential one-pot reactions, see: (a) Jiang, K.; Jia, Z.-J.; Chen, S.; Wu, L.; Chen, Y.-C. *Chem.—Eur. J.* **2010**, *16*, 2852. (b) Enders, D.; Urbanietz, G.; Cassens-Sasse, E.; Keeß, S.; Raabe, G. *Adv. Synth. Catal.* **2012**, *354*, 1481. (c) Hong, B.-C.; Dange, N. S.; Ding, C.-F.; Liao, J.-H. *Org. Lett.* **2012**, *14*, 448. (d) Dange, N. S.; Hong, B.-C.; Lee, C.-C.; Lee, G.-H. *Org. Lett.* **2013**, *15*, 3914. (e) Li, X.; Yang, L.; Peng, C.; Xie, X.; Leng, H.-J.; Wang, B.; Tang, Z.-W.; He, G.; Ouyang, L.; Huang, W.; Han, B. *Chem. Commun.* **2013**, 8692.
- (13) For selected reviews on the bifunctional hydrogen bonding organocatalysis, see: (a) Doyle, A. G.; Jacobsen, E. N. *Chem. Rev.* **2007**, *107*, 5713. (b) Bhadury, P. S.; Song, B.-A.; Yang, S.; Hu, D.-Y.; Xue, W. *Curr. Org. Synth.* **2009**, *6*, 380. (c) Siau, W. Y.; Wang, J. *Catal. Sci. Technol.* **2011**, *1*, 1298. (d) Alemán, J.; Parra, A.; Jiang, H.; Jørgensen, K. A. *Chem.—Eur. J.* **2011**, *17*, 6890. (e) Chauhan, P.; Chimni, S. S. *RSC Adv.* **2012**, *2*, 737.
- (14) CCDC 997149 (for 4c) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif. For X-ray structure, see Supporting Information.