

Origins of Diastereoselectivity in Lewis Acid Promoted Ketene–Alkene [2 + 2] Cycloadditions


Christopher M. Rasik,[†] Young J. Hong,[‡] Dean J. Tantillo,^{*,‡} and M. Kevin Brown^{*,†}

[†]Indiana University, Department of Chemistry, 800 East Kirkwood Avenue, Bloomington, Indiana 47405, United States

[‡]University of California, Davis, Department of Chemistry, 1 Shields Avenue, Davis, California 95616, United States

S Supporting Information


ABSTRACT: A detailed analysis of a Lewis acid promoted ketene–alkene [2 + 2] cycloaddition is reported. The studies have led to a rationalization for an observed inversion of diastereoselectivity between thermally induced and Lewis acid promoted ketene–alkene [2 + 2] cycloadditions. The model is supported with both experimental and computational results.



We recently reported a method for Lewis acid promoted ketene–alkene [2 + 2] cycloadditions.¹ This method offers a number of advantages relative to traditional thermally induced variants.² The advantages include increased reactivity, increased diastereoselectivity, increased regioselectivity, and, for certain cases, inverse diastereoselectivity.¹ The latter point raised interesting questions concerning issues of stereocontrol. Herein, we describe a mechanistic analysis of how the ketene substitution pattern affects the diastereoselectivity of cycloaddition. This analysis of experimental results is also supported with related computational data.

The standard conditions for cycloaddition are illustrated in Scheme 1.¹ Several points are important to note: (1) A

Scheme 1. Lewis Acid Promoted Ketene–Alkene [2 + 2] Cycloadditions




stoichiometric Lewis acid is necessary due to severe product inhibition. (2) For reactions with disubstituted ketenes, Al(III) Lewis acids are uniquely effective. (3) The ketene can be either generated and used as a solution in dichloromethane without removal of the Et₃NHCl byproduct or isolated and purified with little change in yield or diastereoselectivity. In the former case ~2.5 equiv of EtAlCl₂ are necessary because 1 equiv of the Lewis acid reacts with Et₃NHCl. With the latter, if isolated and purified ketenes are used, 1.5 equiv of EtAlCl₂ is sufficient. (4) For many cases, the reactions proceed at –78 °C in <1 h.

During the course of our studies we made an interesting observation regarding the diastereoselectivity of the Lewis acid

promoted [2 + 2] cycloaddition compared to traditional thermal cycloadditions (Scheme 2).¹ For reactions with

Scheme 2. Comparison of Lewis Acid Promoted and Thermally Induced [2 + 2] Cycloadditions



activated alkenes and aryl/alkyl ketenes, different major diastereomers were observed depending on the reaction conditions. Under traditional thermal cycloaddition conditions the *endo*-Ph (7b, 9b) adducts were obtained as the major products, while, under Lewis acid promoted conditions, the *exo*-Ph (7a, 9a) product was preferentially generated. For example, the reaction of ketene 5 with cyclopentadiene at 22 °C for 12 h provides the *endo*-Ph product 7b in 6:1 dr while conditions employing EtAlCl₂ lead to formation of the *exo*-Ph 7a product in 7:1 dr. Similar observations were noted for cycloadditions with indene (Scheme 2, 1 + 8 → 9a/9b). It


Received: August 26, 2014

Published: September 17, 2014

should be noted that control experiments confirm that the inversion in selectivity is not due to a solvent effect.¹

To rationalize the observed differences in diastereoselectivity, we proposed a mechanistic model (Scheme 3; $R_1 = \text{Et}$).¹ In

Scheme 3. Proposed Stereochemical Models¹



accord with previous mechanistic data, under thermal conditions, the alkene likely approaches the ketene *syn* to the smaller Et-group (relative to the conjugated Ph-group). Under Lewis acid promoted conditions, the alkene must approach *syn*-to-Ph. A necessary component of this model is that the Ph-group must rotate out of plane to allow for the approach of the incoming alkene. Neither of these models assumes a concerted or stepwise cycloaddition. However, if the process is stepwise, the putative dipolar intermediate must be short-lived to limit bond rotations that may lead to changes in diastereoselectivity.

To test the proposed models,¹ we carried out cycloadditions with modified ketenes. Since our original hypothesis for Lewis acid promoted reactions assumed that the alkene approaches *syn*-to-Ph, increasing the size of the alkyl group should lead to increased diastereoselectivities. Indeed, increasing the size of the alkyl group (R_1 in Table 1) from Me, Et, *i*-Bu, to *i*-Pr led to

Table 1. Effect of Substituent Size on Diastereoselectivity^a

Reaction scheme showing the Diels-Alder reaction of ketene **1** (Ph-C(=O)-R₁) with cyclopentadiene **8** under conditions A or B to form bicyclic adducts **9** and **10**.

conditions A

toluene 120 °C, 24 h

conditions B

EtAlCl₂, CH₂Cl₂, -78 °C, 2 h


entry	ketene (R ₁)	yield ^b (%)	dr ^c	yield ^b (%)	dr ^c
1 ^d	Me (10)	82	1:7 (13a:13b)	23	3:1 (13a:13b)
2	Et (1)	>95	1:2 (9a:9b)	86	16:1 (9a:9b)
3	<i>i</i> -Bu (11)	>95	1:1 (14a:14b)	56	>20:1 (14a:14b)
4	<i>i</i> -Pr (12)	42	>20:1 (15a:15b)	80	>20:1 (15a:15b)

^aSee the Supporting Information (SI) for details. ^bYields determined by ¹H NMR analysis with an internal standard. ^cDetermined by ¹H NMR analysis of the unpurified reaction mixture. ^dKetene was generated *in situ*. See the SI.

a significant increase in diastereoselectivity (compare entries 1–4, Table 1, conditions B). Thermal cycloaddition with Ph/Me or Ph/Et substituted ketenes predominantly generated the *endo*-Ph adducts (9b, 13b) likely from the approach of the alkene *syn*-to-alkyl. However, when the steric bulk of the alkyl substituent was increased to *i*-Pr, the *exo*-Ph isomer was generated in >20:1 dr. It seems that the steric bulk imposed by the *i*-Pr group is too significant to allow for the approach of the alkene.

Reactions with ketene 16 were also investigated because it bears an aryl ring that is incapable of rotation out of plane. Thus, approach of the alkene can only occur from the less hindered face according to the model illustrated in Scheme 4.

Scheme 4. Controlling Facial Selectivity



In accordance with this hypothesis, cycloadditions carried out under both thermal and Lewis acid promoted conditions provided the *endo*-Ar adduct 17 in excellent diastereoselectivity.

Based on the data described above it seems that the selectivity of the cycloaddition is dictated by the approach of the alkene to different faces of the ketene. A question that remained is why does the Lewis acid affect the approach of the alkene from different faces of the ketene? To assist in our understanding, we turned toward computational evaluation³ of the potential energy surfaces of both the thermal and Lewis acid promoted reactions.⁴ A recent theoretical study by Wei, Tang and co-workers described many features of the potential energy surface for the reaction of phenylbenzylketene with cyclopentadiene (5 + 6 in Scheme 2).⁵ Herein, we describe results for 1 + 8, comparing and contrasting our results with theirs and highlighting additional important features contributing to stereocontrol.

Initial computational investigations revealed that coordination of MeAlCl₂ to a ketene reduces the barrier to rotation of the aryl ring (Figure 1). This observation is consistent with the notion of aryl rotation for the Lewis acid promoted




Figure 1. Barrier to rotation of Ph/Me ketene and Ph/Me ketene Lewis acid complex.

cycloadditions described above. Note that, in the Lewis acid complexed structure, the C=C bond of the ketene shortens slightly (from 1.33 to 1.31 Å) and the C–C_{ipso} bond lengthens slightly (from 1.48 to 1.49 Å).

MeAlCl₂-promoted cycloaddition of **1** + **8** was found to involve reaction coordinates in the borderlands between stepwise and concerted asynchronous processes.⁶ Pathways to the major and minor products were found, depending on the level of theory used, to involve either dipolar intermediates with very small barriers for cyclobutane ring closure (<2.5 kcal/mol) or no intermediates but very asynchronous formation of the two new C–C bonds (free energies, –78 °C, in CH₂Cl₂ or the gas phase). In either case, the lifetime of a dipolar species is expected to be very short and formation of the first new C–C bond occurs in the rate-determining transition state structures (Figure 2, top). Note that the central ketene carbons in these




Figure 2. Transition state structures for formation of diastereomers **9a/b** in the presence of MeAlCl₂. Selected distances shown in Å (M06-2X/6-31G(d)).³

(very early) transition state structures sit over the center of the indene π -bond; only later along the reaction coordinates is this symmetrical bridging disrupted. In addition, the aryl groups in both transition state structures are indeed twisted out of conjugation: in the major transition state structure so as to avoid the indene (and possibly engage in favorable C–H $\cdots\pi$ interactions) and in the minor transition state structure so as to avoid the Lewis acid. The predicted free energy barriers (from MeAlCl₂ complexed **1** + free **8**) are 6.67 and 6.85 kcal/mol (M06-2X/6-31G(d)),³ corresponding to a smaller product ratio than that observed experimentally (a 20:1 product ratio

corresponds to a preference of 1.3 kcal/mol). A third transition state structure was located with a different orientation of the Lewis acid (Figure 2, bottom). This transition state structure is 2.00 kcal/mol lower in free energy than the lower energy transition state structure from the top of Figure 2. If this transition state structure leads to the major product, then a larger product ratio is expected. The pathway forward from this transition state structure leads to a flat portion of the energy surface, initially forming a dipolar intermediate with the appropriate C–C bond formed. However, an exit channel from this region of the energy surface leads to a regioisomer of **9a**.

The uncatalyzed thermal reaction was also predicted to be concerted with highly asynchronous bond forming events. Barriers for thermal cycloaddition were predicted to be much larger (37–38 kcal/mol), consistent with experiment. Transition state structures leading to **9a** and **9b** were predicted to be nearly equienergetic (low selectivity is observed experimentally, Scheme 2), although the transition state structure leading to **9a** was favored slightly ($\Delta\Delta G = 0.65$ kcal/mol with M06-2X/6-31G(d)). In the transition state structure leading to **9a**, the phenyl ring twists out of conjugation, but in the transition state structure leading to **9b**, the phenyl ring does not twist out of conjugation (Figure 3), consistent with the twist in the




Figure 3. Transition state structures for formation of diastereomers **9a/b** in the absence of a Lewis acid. Selected distances shown in Å (M06-2X/6-31G(d)).³

transition state structure leading to Lewis acid complexed **9b** being dependent on the presence of the Lewis acid, likely the result of a steric effect.

Unusual energy surfaces have been described previously by Singleton and co-workers for other ketene–alkene and ketene–diene cycloadditions, and dynamics calculations have played an important role in understanding the origins of selectivity for these processes.⁷ Dynamics calculations on the complex situation discussed above will be described in a future report, but our tentative diastereoselectivity model is summarized in Scheme 3. In short, selectivity for **9a** is increased in the Lewis acid promoted case due to steric clashes between the Lewis acid and phenyl groups that disfavor the **9b**-forming transition state structure.⁸ These clashes induce a conformational change, which does not occur in the Lewis acid free system (Figure 3), that causes a reduction in conjugation between the phenyl ring and the C=C bond of the ketene.

In summary, we have provided both experimental and computational data that support the proposed pathway for diastereoselectivity. Subsequent studies will be directed toward utilizing this knowledge for future iterations of reaction development.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental data, computational data, and analytical data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: brownmkb@indiana.edu.

*E-mail: djtantillo@ucdavis.edu.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We gratefully acknowledge support from UC Davis and the National Science Foundation (CHE-030089 for computer time via the XSEDE program). Indiana University is also acknowledged for generous financial support.

■ REFERENCES

- (1) (a) Rasik, C. M.; Brown, M. K. *J. Am. Chem. Soc.* **2013**, *135*, 1673. For a review, see: (b) Rasik, C.; Brown, M. *Synlett* **2014**, 25, 760.
- (2) For excellent reviews, see: (a) Snider, B. B. *Chem. Rev.* **1988**, *88*, 793–811. (b) Hyatt, J. A.; Reynolds, P. W. *Organic Reactions*; Wiley: New York, NY, 1994. (c) Tidwell, T. T. *Ketenes*; Wiley: New York, NY, 1995. (d) Tidwell, T. T. *Ketenes II*; Wiley-Interscience: Hoboken, NJ, 2006. (e) Tidwell, T. T. *Eur. J. Org. Chem.* **2006**, 563–576. (f) Danheiser, R. L. *Science of Synthesis: Compounds with Four and Three Carbon Heteroatom Bonds*, Vol. 23; Georg Thieme Verlag KG: Germany, 2006. (g) Allen, A. D.; Tidwell, T. T. *Chem. Rev.* **2013**, *113*, 7287.
- (3) Calculations were performed with GAUSSIAN09 (Frisch, M. J. et al.; full reference in Supporting Information). Geometries were optimized using the B3LYP and M06-2X methods with the 6-31G(d) basis set, and all stationary points were characterized as minima or transition state structures using frequency calculations at the same level: (a) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648–5652. (b) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 1372–1377. (c) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785–789. (d) Stephens, P. J.; Devlin, F. J.; Chabalowski, C. F.; Frisch, M. J. *J. Phys. Chem.* **1994**, *98*, 11623–11627. (e) Zhao, Y.; Truhlar, D. *Theor. Chem. Acc.* **2008**, *120*, 215–241 (this method provides a better treatment of dispersion). Intrinsic reaction coordinate (IRC) calculations were used for further characterization of transition state structures: (f) Gonzalez, C.; Schlegel, H. B. *J. Phys. Chem.* **1990**, *94*, 5523–5527. (g) Fukui, K. *Acc. Chem. Res.* **1981**, *14*, 363–368. Free energies in solution (toluene (120 °C) and CH₂Cl₂ (–78 °C) in the absence and presence of the Lewis acid, respectively) are computed using the integral equation formalism variant (IEFPCM) with the SMD solvation method: (h) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. *J. Phys. Chem. B* **2009**, *113*, 6378–6396. Reported free energies are relative to that of the separate reactants (alkene + ketene for thermal condition and alkene + Lewis acid–ketene complex for the Lewis acid promoted reaction). Images of computed structures are created with *Ball & Stick*: (i) Müller, N.; Falk, A.; Gsaller, G. *Ball & Stick V.4.0a12, molecular graphics application for MacOS computers*; Johannes Kepler University: Linz, 2004. MeAlCl₂ was used as the model Lewis acid. M06-2X results are described in the text; see Supporting Information for B3LYP results.
- (4) For earlier computational evaluation of ketene–alkene [2 + 2] cycloadditions, see: (a) Wang, X.; Houk, K. N. *J. Am. Chem. Soc.* **1990**, *112*, 1754. (b) Bernardi, F.; Bottoni, A.; Robb, M. A.; Venturini, A. *J. Am. Chem. Soc.* **1990**, *112*, 2106. (c) Valenti, E.; Pericas, M. A.; Moyano, A. *J. Org. Chem.* **1990**, *55*, 3582. (d) Salzner, U.; Bachrach, S. M. *J. Org. Chem.* **1996**, *61*, 237.
- (5) Wang, Y.; Wei, D.; Li, Z.; Zhu, Y.; Tang, M. *J. Phys. Chem. A* **2014**, *118*, 4288.
- (6) (a) Tantillo, D. J. *J. Phys. Org. Chem.* **2008**, *21*, 561–570. (b) Williams, A. *Concerted Organic and Bio-Organic Mechanisms*; CRC Press: Boca Raton, FL, 2000.
- (7) (a) Ussing, B. R.; Hang, C.; Singleton, D. A. *J. Am. Chem. Soc.* **2006**, *128*, 7594–7607. (b) Gonzalez-James, O. M.; Kwan, E. E.; Singleton, D. A. *J. Am. Chem. Soc.* **2012**, *134*, 1914–1917. (c) Pemberton, R. P.; Hong, Y. J.; Tantillo, D. J. *Pure Appl. Chem.* **2013**, *85*, 1949–1957.
- (8) Wei, Tang and co-workers also invoke steric effects in rationalizing the diastereoselectivity for the 5 + 6 reaction (see ref 5).