


120 °C (Table 1, entry 1). The 2,4-diphenylpyridine product should come from the cyclization of two molecules of

Table 1. Optimization of the Reaction Conditions^a



entry	[Fe]	oxidant	solvent	temp (°C)	yield (%)
1	FeCl ₂		toluene	120	18
2	FeCl ₂	O ₂	toluene	120	22
3	FeCl ₂	TBHP ^b	toluene	120	16
4	FeCl ₂	H ₂ O ₂ ^c	toluene	120	0
5	FeCl ₂	DTBP	toluene	120	52
6	FeCl ₃	DTBP	toluene	120	60
7	Fe(OTf) ₂	DTBP	toluene	120	26
8	Fe(OTf) ₃	DTBP	toluene	120	20
9		DTBP	toluene	120	0
10	FeCl ₃	DTBP	1,4-dioxane	120	16
11	FeCl ₃	DTBP	CH ₃ CN	120	74
12	FeCl ₃	DTBP	DCE	120	86(68) ^d
13	FeCl ₃		DCE	120	37
14	FeCl ₃	DTBP	DCE	100	62
15	FeCl ₃	DTBP	DCE	140	80

^aReaction conditions: **1a** (0.4 mmol), **2a** (0.4 mmol), [Fe] (10 mol %), oxidant (0.6 mmol), solvent (3 mL), 5 h, in air; isolated yields.

^bTBHP (5–6 M in decane). ^c30 wt % solution in water. ^dReaction was performed under argon.

acetophenone oxime acetate **1a** and a methyl group on *N,N*-dimethylaniline **2a**. The reaction indicated that N–O bond cleavage of the ketoxime acetates occurred in the presence of Fe catalyst and tertiary anilines. This interesting primary result prompted us to optimize the reaction conditions to develop an Fe-catalyzed reaction of ketoxime acetates and tertiary anilines for the synthesis of substituted pyridines.


Various oxidants such as O₂, *tert*-butyl hydroperoxide (TBHP), H₂O₂, and di-*tert*-butyl peroxide (DTBP) were then screened to optimize the reaction (Table 1, entries 2–5). DTBP was found to be the most effective oxidant, providing **3aa** in 52% yield. Further experiments showed that FeCl₃ was superior to FeCl₂ (Table 1, entry 6). Other iron catalyst precursors such as Fe(OTf)₂ and Fe(OTf)₃ were less reactive in the reaction (Table 1, entries 7 and 8). Notably, no reaction occurred in the absence of the iron catalyst or in the presence of copper catalyst,¹⁹ indicating that the iron catalyst is essential for the reaction (Table 1, entry 9). Next, various solvents such as 1,4-dioxane, CH₃CN, and DCE were screened (Table 1, entries 10–12). DCE was found to be the optimal solvent for the reaction, affording pyridine **3aa** in 86% yield (Table 1, entry 12). Notably, the 2,4-diphenylpyridine **3aa** was obtained in 68% yield when the reaction was performed under argon, and the 2,4-diphenylpyridine **3aa** was observed in 37% yield in absence of the DTBP under air (Table 1, entry 13). These results indicate that oxygen combined with DTBP acted as the oxidant in the reaction. Moreover, the reaction temperature was also varied. The optimum temperature for the reaction was found to be 120 °C (Table 1, entries 14 and 15).

To evaluate the potential of amines to function as a source of one-carbon synthon, various tertiary and secondary amines were investigated under the standard conditions (see Scheme S1). The reaction of acetophenone oxime acetate **1a** with 4-methyl-*N,N*-

dimethylaniline **2b** affords the desired pyridine **3aa** in 74% yield, whereas electron-withdrawing (–Br, –CN) groups on aniline decreased the yield of **3aa**. These results clearly indicate that the choice of amines is crucial for the reaction. Subsequently, different acetophenone oxime carboxylates were investigated to test their reactivity. These results indicate that the carboxylate group at the ketoximes has little effect on the cyclization reaction (see Scheme S2).

With the optimized reaction conditions in hand, the scope of the reaction was investigated (Scheme 2). This novel Fe-

Scheme 2. Fe-Catalyzed Cyclization of Ketoxime Acetates and *N,N*-Dimethylaniline^a




^aReaction conditions: **1** (0.4 mmol), **2a** (0.4 mmol), FeCl₃ (10 mol %), (*t*-BuO)₂ (0.6 mmol, 1.5 equiv), DCE (3 mL), in air; isolated yields. ^bCH₃CN was used as solvent.

catalyzed cyclization reaction exhibited good functional-group tolerance and proved to be a general method for the facile construction of 2,4-disubstituted pyridines. Ketoxime acetates with electron-neutral or electron-donating groups on aryl rings such as methyl, alkyl, and methoxyl afforded the corresponding 2,4-diarylpyridines **3ba–da,ga–ha** in good yields (61–80%).²⁰ Ketoxime acetates with electron-withdrawing substituents, such as fluoro, chloro, and bromo, were well tolerated and afforded the corresponding 2,4-diarylpyridines **3ia–ja,la** in 70–74% yields. Moreover, trifluoromethyl-substituted ketoxime acetate proceeded smoothly to afford the desired 2,4-di(4-(trifluoromethyl)phenyl)pyridine **3ma** in 68% yield in CH₃CN.²¹ These results indicated that the electronic nature of

the ketoxime acetates has little effect on the reaction. However, the steric effect plays a role in the reaction. *o*-Methyl- or chloro-substituted ketoxime acetates afforded the desired 2,4-diarylpyridines **3a–fa,ka** in moderate yields. In addition, β -naphthyl methyl ketoxime acetate **1n**, furanyl methyl ketoxime acetate **1o**, and thiophene-2-yl methyl ketoxime acetate **1p** also reacted well to give the desired 2,4-disubstituted pyridines **3na–pa** in 60–66% yields, respectively. However, no reaction occurred when an aliphatic ketoxime acetate such as 3,3-dimethylbutan-2-one oxime acetate was employed as the substrate.

On the basis of the aforementioned results, it was expected that 2,4,6-trisubstituted pyridines could be synthesized by using alternative *N,N*-dialkylanilines. Expectedly, 6-methyl-2,4-diphenylpyridine **3ai** was obtained in 62% yield in the presence of *N,N*-diethylaniline **2i** (Scheme 3, eq 1). 2,4,6-Triphenylpyridine **3aj**

Scheme 3. Construction of 2,4,6-Trisubstituted Pyridine by Using Alternative *N,N*-Dialkylanilines




was also obtained in 71% yield when *N,N*-dibenzylaniline **2j** was used as the substrate (Scheme 3, eq 2). These results also reveal that the carbon at the 6-position of pyridine product **3** definitely comes from the methylene carbon on the *N,N*-dialkylaniline **2**.

To study the byproduct in the dealkylation of *N,N*-dialkylanilines, the cyclization of acetophenone oxime acetate **1a** and *N*-phenyl-1,2,3,4-tetrahydroisoquinoline **2k** was performed under the standard reaction conditions (Scheme 3, eq 3). In this reaction, *N*-(2-(4,6-diphenylpyridin-2-yl)phenethyl)-aniline **3ak** was obtained in 45% yield, clearly indicating that the dealkylation of *N,N*-dialkylanilines produced *N*-alkylaniline species in the reaction.


On the basis of the above results,²² a tentative mechanism for the Fe-catalyzed cyclization is proposed, as shown in Scheme 4. The first step of the reaction is the reduction of Fe^{3+} by *N,N*-dimethylaniline **2a** to afford Fe^{2+} species and iminium ion intermediate **2a'**.²³ Then, reductive cleavage of the N–O bond of ketoxime acetate **1** by Fe^{2+} species via a two-step single electron-transfer process forms the imine anion intermediate **A** and Fe^{3+} species. Rapid coordination of intermediate **A** and Fe^{3+} species gives the imino- Fe^{3+} intermediate **B**. Next, nucleophilic addition of **B** to iminium ion intermediate **2a'** produces the enamine intermediate **C**. Addition of enamine intermediate **C** to a second molecular of ketoxime acetate **1** gives the intermediate **D** followed by the release of NH_2OAc to afford the intermediate **E**.¹⁴ Tautomerization of **E** gives intermediate **F**, and intramolecular deamination cyclization of the intermediate **F** assisted by FeCl_3 forms the intermediate **G**. Finally, oxidative aromatization of intermediate **G** produces the 2,4-diarylpyridine **3**.

Scheme 4. Tentative Mechanisms for Fe-Catalyzed Cyclization of Ketoxime Acetates and Tertiary Anilines



In the above reaction, the 2,4-diarylpyridines with identical aryls were synthesized. To extend the flexibility of the reaction, Fe-catalyzed cycloaddition of ketoxime acetate **1a** and *N,N*-dicinnamylaniline **2l** was conducted under the above conditions. Delightfully, the desired 2,4-diphenylpyridine **3aa** was obtained in 66% yield. Inspired by this result, we synthesized a series of 2,4-diarylpyridines with different aryls by this method using different ketoxime acetates (Scheme 5). Notably, the procedure

Scheme 5. Fe-Catalyzed Cyclization of Ketoxime Acetates and Dicinnamylaniline^a




^aReaction conditions: **1** (0.2 mmol), **2l** (0.24 mmol), FeCl_3 (10 mol %), $(t\text{-BuO})_2$ (0.6 mmol), DCE (3 mL), in air; isolated yields.

shows a good method for the synthesis of 2,4-diarylpyridines with different aryls. Ketoxime acetates with methyl, alkyl, fluoro, chloro, and bromo on the aromatic ring gave the corresponding 2,4-diarylpyridines **3bl–ll** in 60–72% yields, indicating that the reaction was insensitive to the electronic nature of the ketoxime acetates. β -Naphthyl-substituted ketoxime acetate **1n** proceeded smoothly as well to give the corresponding 2-(naphthalen-2-yl)-4-phenylpyridine **3nl** in 56% yield. In addition, a heterocyclic ketoxime acetate such as **1p** was also tolerated and afforded the desired 4-phenyl-2-(thiophene-2-yl)pyridine **3pl** in 69% yield.

The mechanism proposed for the reaction is illustrated in Scheme 6. Nucleophilic addition of intermediate **B** to the

Scheme 6. Proposed Mechanism for the Synthesis of Pyridines with Different Aryls



iminium ion **2I'** affords intermediate **H**. The intramolecular cyclization/deamination of intermediate **H** gives the intermediate **I**. Finally, oxidative aromatization of **I** produces the 2,4-diarylpyridine **3**.

In conclusion, we have developed an unprecedented Fe-catalyzed cyclization of ketoxime carboxylates and *N,N*-dialkylanilines for the synthesis of 2,4-disubstituted and 2,4,6-trisubstituted pyridines. It is a novel reaction for Fe-catalyzed N–O bond cleavage of the ketoxime carboxylates in the presence of tertiary anilines. The methylene carbon on *N,N*-dialkylanilines functioned as a source of one-carbon synthon in the reaction. The catalytic system was allowed to extend to cyclization of ketoxime carboxylates and *N,N*-dicinnamylaniline for the synthesis of 2,4-diarylpyridines with different aryls. These methods show good procedures for rapid elaboration of the readily available ketoxime carboxylates into diverse valuable substituted pyridines in high yields under mild reaction conditions. Further study of the scope and mechanism is underway.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b00326.

Detailed experimental procedures, characterization data, and copies of NMR spectra for all products (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: guanzzhh@nwnu.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by generous grants from the National Natural Science Foundation of China (21272183 and 21472147) and the Fund of Raising Stars of Shanxi Province (2012KJXX-26).

■ REFERENCES

- (1) (a) Desimoni, G.; Faita, G.; Quadrelli, P. *Chem. Rev.* **2014**, *114*, 6081. (b) Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. *J. Med. Chem.* **2014**, *57*, 5845. (c) Byrne, J. P.; Kitchen, J. A.; Gunnlaugsson, T. *Chem. Soc. Rev.* **2014**, *43*, 5302. (d) Chelucci, G. *Chem. Soc. Rev.* **2006**, *35*, 1230.
- (2) For reviews, see: (a) Allais, C.; Grassot, J.-M.; Rodriguez, J.; Constantieux, T. *Chem. Rev.* **2014**, *114*, 10829. (b) Gulevich, A. V.; Dudnik, A.; Chernyak, N.; Gevorgyan, V. *Chem. Rev.* **2013**, *113*, 3084. (c) Hill, M. D. *Chem. - Eur. J.* **2010**, *16*, 12052. (d) Henry, G. D. *Tetrahedron* **2004**, *60*, 6043.
- (3) (a) Yoshida, M.; Mizuguchi, T.; Namba, K. *Angew. Chem., Int. Ed.* **2014**, *53*, 14550. (b) Lei, C.-H.; Wang, D.-X.; Zhao, L.; Zhu, J.; Wang,

M.-X. *J. Am. Chem. Soc.* **2013**, *135*, 4708. (c) Jiang, Y.; Park, C.-M.; Loh, T.-P. *Org. Lett.* **2014**, *16*, 3432. (d) Chen, M. Z.; Micalizio, G. C. *J. Am. Chem. Soc.* **2012**, *134*, 1352. (e) Gati, W.; Rammah, M. M.; Rammah, M. B.; Couty, F.; Evano, G. *J. Am. Chem. Soc.* **2012**, *134*, 9078.

(4) (a) Loy, N. S. Y.; Singh, A.; Xu, X.; Park, C.-M. *Angew. Chem., Int. Ed.* **2013**, *52*, 2212. (b) Wang, Y.-F.; Toh, K. K.; Ng, E. P. J.; Chiba, S. *J. Am. Chem. Soc.* **2011**, *133*, 6411.

(5) (a) Xi, L.-Y.; Zhang, R.-Y.; Liang, S.; Chen, S.-Y.; Yu, X.-Q. *Org. Lett.* **2014**, *16*, 5269. (b) Shi, Z.; Loh, T.-P. *Angew. Chem., Int. Ed.* **2013**, *52*, 8584. (c) Michlik, S.; Kempe, R. *Angew. Chem., Int. Ed.* **2013**, *52*, 6326.

(6) (a) Thiel, I.; Jiao, H.; Spannenberg, A.; Hapke, M. *Chem. - Eur. J.* **2013**, *19*, 2548. (b) Domínguez, G.; Pérez-Castells, J. *Chem. Soc. Rev.* **2011**, *40*, 3430.

(7) (a) Prechter, A.; Henrion, G.; Faudot dit Bel, P.; Gagosz, F. *Angew. Chem., Int. Ed.* **2014**, *53*, 4959. (b) Movassaghi, M.; Hill, M. D. *J. Am. Chem. Soc.* **2006**, *128*, 4592.

(8) Sukhorukov, A. Y.; Ioffe, S. L. *Chem. Rev.* **2011**, *111*, 5004.

(9) (a) Narasaka, K.; Kitamura, M. *Eur. J. Org. Chem.* **2005**, *2005*, 4505.

(b) Narasaka, K. *Pure Appl. Chem.* **2002**, *74*, 143.

(10) Zaman, S.; Mitsuru, K.; Abell, A. D. *Org. Lett.* **2005**, *7*, 609.

(11) Gerfaud, T.; Neuville, L.; Zhu, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 572.

(12) Tan, Y.; Hartwig, J. F. *J. Am. Chem. Soc.* **2010**, *132*, 3676.

(13) Faulkner, A.; Bower, J. F. *Angew. Chem., Int. Ed.* **2012**, *51*, 1675.

(14) (a) Ren, Z.-H.; Zhang, Z.-Y.; Yang, B.-Q.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2011**, *13*, 5394. (b) Guan, Z.-H.; Zhang, Z.-Y.; Ren, Z.-H.; Wang, Y.-Y.; Zhang, X. *J. Org. Chem.* **2011**, *76*, 339. (c) Liang, H.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Chem. - Eur. J.* **2013**, *19*, 9789. (d) Ran, L.; Liang, H.; Guan, Z.-H. *Youji Huaxue* **2013**, *33*, 66. (e) Ran, L.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Green Chem.* **2014**, *16*, 112. (f) Du, W.; Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Chem. Commun.* **2014**, *50*, 7437. (g) Zhao, M.-N.; Hui, R.-R.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2014**, *16*, 3082.

(15) (a) Huang, H.; Ji, X.; Wu, W.; Jiang, H. *Chem. Soc. Rev.* **2015**, *44*, 1155. (b) Tang, X.; Huang, L.; Xu, Y.; Yang, J.; Wu, W.; Jiang, H. *Angew. Chem., Int. Ed.* **2014**, *53*, 4205. (c) Ke, J.; Tang, Y.; Yi, H.; Li, Y.; Cheng, Y.; Liu, C.; Lei, A. *Angew. Chem., Int. Ed.* **2015**, *54*, 6604. (d) Liu, S.; Liebeskind, L. S. *J. Am. Chem. Soc.* **2008**, *130*, 6918. (e) Jiang, H.; Yang, J.; Tang, X.; Li, J.; Wu, W. *J. Org. Chem.* **2015**, *80*, 8763. (f) Mahernia, S.; Mahdavi, M.; Adib, M. *Synlett* **2014**, *25*, 1299. (g) Cai, Z.; Lu, X.; Wang, S.; Ji, S. *Acta Chim. Sinica* **2014**, *72*, 914.

(16) Wei, Y.; Yoshikai, N. *J. Am. Chem. Soc.* **2013**, *135*, 3756.

(17) (a) Deb, I.; Yoshikai, N. *Org. Lett.* **2013**, *15*, 4254. (b) Majireck, M. M.; Witek, J. A.; Weinreb, S. M. *Tetrahedron Lett.* **2010**, *51*, 3555. (c) Nakanishi, S.; Hirano, T.; Otsuji, Y.; Itoh, K. *Chem. Lett.* **1987**, *16*, 2167.

(18) (a) Ouyang, K.; Hao, W.; Zhang, W.-X.; Xi, Z. *Chem. Rev.* **2015**, *115*, 12045. (b) Li, H.; He, Z.; Guo, X.; Li, W.; Zhao, X.; Li, Z. *Org. Lett.* **2009**, *11*, 4176. (c) Liu, W.; Liu, J.; Ogawa, D.; Nishihara, Y.; Guo, X.; Li, Z. *Org. Lett.* **2011**, *13*, 6272. (d) Wu, W.; Su, W. *J. Am. Chem. Soc.* **2011**, *133*, 11924.

(19) No reaction occurred when a variety of copper catalysts such as CuI, CuBr, CuCl, CuBr₂, CuCl₂, and Cu(OAc)₂ were screened in the reaction.

(20) The byproduct of the reaction was the corresponding ketone.

(21) All of the reactions listed in Scheme 2 were performed both in DCE and CH₃CN. DCE gave better yields in most cases, but CH₃CN gave better result for **3ea** and **3ma**.

(22) Fe-catalyzed three-component cyclization of ketoxime acetate, ketone (or N-Ts-protected ketoxime), and *N,N*-dimethylaniline under the standard conditions was not successful; see the Supporting Information. This result indicated that the ketone is not an intermediate in the reaction.

(23) (a) Ratnikov, M. O.; Xu, X.; Doyle, M. P. *J. Am. Chem. Soc.* **2013**, *135*, 9475. (b) Jia, F.; Li, Z. *Org. Chem. Front.* **2014**, *1*, 194. (c) Takasu, N.; Oisaki, K.; Kanai, M. *Org. Lett.* **2013**, *15*, 1918.