


Cp*Co^{III}-Catalyzed Dehydrative C–H Allylation of 6-Arylpurines and Aromatic Amides Using Allyl Alcohols in Fluorinated AlcoholsYouka Bunno,[†] Nanami Murakami,[†] Yudai Suzuki,[‡] Motomu Kanai,[‡] Tatsuhiko Yoshino,^{*,†,§} and Shigeki Matsunaga^{*,†,§}[†]Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan[‡]Graduate School of Pharmaceutical Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan[§]ACT-C, Japan Science and Technology Agency, Sapporo 060-0812, Japan

S Supporting Information

ABSTRACT: Cp*Co^{III}-catalyzed C–H allylation of various aromatic C–H bonds using allyl alcohols as allylating reagents is described. Improved reaction conditions using fluorinated alcohol solvents afforded efficient directed C–H allylation of 6-arylpurines, benzamides, and a synthetically useful Weinreb amide with good functional group compatibility.



Allylation of aromatic compounds is very important in organic synthesis because the allyl moiety can be further converted into a wide variety of functional groups. In terms of atom economy,¹ step economy,² and functional group compatibility, direct allylation of inert C–H bonds under transition metal catalysis³ is more attractive than classical allylation methods, such as cross-coupling reactions of functionalized arenes.^{4,5} Therefore, aromatic and olefinic C–H allylation reactions using allyl acetates,⁶ carbonates,⁷ phosphates,⁸ halides,⁹ and ethers¹⁰ as allylating reagents were developed over the past decade using various transition metals. Among them, Cp*Rh^{III}-catalyzed¹¹ C–H allylation reactions using preactivated allyl alcohols^{6c,7b–d,8d,9d,10b} are especially noteworthy due to their mild reaction conditions and broad substrate scope.¹² Those reactions are thought to proceed via C–H metalation, olefin insertion, and subsequent β -elimination of oxygen-containing functional groups.


We¹³ and other groups^{14,15} have investigated Cp*Co^{III} catalysts for C–H bond functionalization as an inexpensive alternative¹⁶ to Cp*Rh^{III} catalysts after our first report in 2013.^{13a} Glorius¹⁷ and Ackermann¹⁸ independently reported Cp*Co^{III}-catalyzed C–H allylation of indoles, pyrroles, and 2-phenylpyridines using preactivated allyl alcohol derivatives under mild conditions. The scope of this reaction was further expanded to benzamide derivatives.¹⁹ On the other hand, we recently reported that unactivated allyl alcohols can be used directly for aromatic C–H allylation under Cp*Co^{III} catalysis, while Cp*Rh^{III} exhibited inferior catalytic activity.²⁰ Key to this reaction is likely the more favored β -hydroxy elimination over the potentially competing β -hydride elimination after insertion of a C–C double bond.²¹ The substrate scope of our previous protocol, however, was limited to indoles and 1-phenylpyrazole.

Although it was reported that *N*-methylbenzamide can be allylated using an allyl alcohol, the yield was low, and its generality was not demonstrated.¹⁹ During the preparation of our manuscript, Kapur reported Ru^{II}-catalyzed C–H allylation using allyl alcohols, but the scope was still limited to indoles, an indoline, and 2-phenylpyridine.²² Here we report an improved protocol for Cp*Co^{III}-catalyzed dehydrative C–H allylation of aromatic compounds using allyl alcohols. The use of fluorinated alcohol solvent significantly enhanced the reactivity, enabling allylation of 6-arylpurines, benzamides, and an aromatic Weinreb amide in moderate to excellent yield.

We first selected 6-phenylpurine **1a** as a model substrate (Table 1). Transition-metal-catalyzed C–H bond functionalization of 6-arylpurines has recently attracted much attention²³ due to their biological activities.²⁴ Previously reported conditions for allylation of indoles²⁰ using [Cp*Co(CO)I₂] (5 mol %), AgOTf (10 mol %), and AgOAc (10 mol %) in DCE at 60 °C selectively afforded monoallylated product **3a**, but in only 13% yield (entry 1). Screening of the solvent led to a significantly improved yield of **3a** (entries 2–6). Although most of the solvents were ineffective, 2,2,2-trifluoroethanol (TFE) and 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) exhibited high reactivity,²⁵ and **3a** was obtained in good yields (entries 5, 6). Changing AgOTf to other Ag salts did not affect the reactivity (entries 7–10), whereas the yield was decreased when no cationic Ag salts were used (entry 11). Other acetate bases were less effective than AgOAc (entries 12–14). Almost no product was detected in the absence of acetate bases (entry 15), indicating that a carboxylate-assisted concerted

Received: March 23, 2016

Published: April 27, 2016

Table 1. Optimization of Reaction Conditions^a



| entry | solvent | Ag salt | base | yield ^b (%) |
|-------|---------|--------------------|-------|------------------------|
| 1 | DCE | AgOTf | AgOAc | 13 |
| 2 | THF | AgOTf | AgOAc | trace |
| 3 | toluene | AgOTf | AgOAc | 6 |
| 4 | MeOH | AgOTf | AgOAc | 3 |
| 5 | TFE | AgOTf | AgOAc | 81 (78 ^c) |
| 6 | HFIP | AgOTf | AgOAc | 72 |
| 7 | TFE | AgSbF ₆ | AgOAc | 78 |
| 8 | TFE | AgNTf ₂ | AgOAc | 79 |
| 9 | TFE | AgBF ₄ | AgOAc | 80 |
| 10 | TFE | AgPF ₆ | AgOAc | 73 |
| 11 | TFE | none | AgOAc | 42 |
| 12 | TFE | AgOTf | NaOAc | 8 |
| 13 | TFE | AgOTf | KOAc | 29 |
| 14 | TFE | AgOTf | CsOAc | 12 |
| 15 | TFE | AgOTf | none | trace |

^aThe reactions were run using **1a** (0.10 mmol) and **2a** (0.15 mmol) in indicated solvent (0.1 M) at 60 °C. ^bDetermined by ¹H NMR analysis of the crude mixture using dibenzyl ether as an internal standard. ^cIsolated yield after silica gel column chromatography at 0.30 mmol scale.


metalation–deprotonation mechanism works in this reaction.²⁶ Under the best reaction conditions in entry 5, the allylated product **3a** was isolated in 78% yield after chromatographic purification.

After optimizing the conditions, the scope of 6-aryl-purines was investigated (Scheme 1). In addition to *N*9-alkylated purine derivatives **1a**, **1b**, and *N*9-tosylated purine **1c**, fully acetylated 6-phenylpurine riboside **1d** was allylated without any difficulty to give **3d** in 88% yield. Both electron-donating and -withdrawing groups were compatible on the phenyl moiety, and products **3e–g** were obtained in 70%–93% yield. Heteroaryl-purine **1h** also successfully afforded allylated product **3h** in 59% yield. The reaction using tertiary allyl alcohol **2b** proceeded γ -selectively to afford prenylated product **3i**. The scalability of the new protocol was also confirmed by a gram-scale experiment in which **3b** was obtained in 95% yield.

Next we turned our attention to the allylation of other aromatic compounds. After minor modification of the reaction conditions,²⁷ benzamide derivatives **4** were also allylated using 20 mol % of AgNTf₂ and HFIP solvent, as shown in Scheme 2. Unsubstituted benzamide **4a** and *p*-Me-benzamide **4b** afforded the corresponding allylated products in moderate yields along with the bis-allylated byproducts.²⁸ Allylation of *o*-F-benzamide **4c** proceeded in 71% yield. Various *m*-substituted benzamide derivatives, including *I*-substituted substrate **4h**, afforded the monoallylated products in acceptable yields with >10/1 regioselectivity (**5d–5h**). *N*-Methylbenzamide **4i** and *N,N*-dimethylbenzamide **4j** also afforded the allylated products **5i** and **5j** albeit in diminished yields. To confirm the importance of the fluorinated alcohol, a negative control experiment using previously

Scheme 1. Scope of Dehydrative Allylation of 6-Arylpurines^a


^aThe reactions were run using **1** (0.30 mmol), **2**, [Cp*Co(CO)I₂] (5 mol %), AgOTf (10 mol %), AgOAc (10 mol %) in TFE (0.1 M) at 60 °C for 8 h unless otherwise noted. The indicated yields are isolated yields after silica gel column chromatography. ^b1.0 g (3.5 mmol) of **1b** was used, and 1.08 g of **3b** was isolated. ^cThe reaction time was 24 h. ^dThe reaction was run at 70 °C.

Scheme 2. Dehydrative Allylation of Benzamides^a

^aThe reactions were run using **4** (0.30 mmol), **2** (0.90 mmol), [Cp*Co(CO)I₂] (5 mol %), AgNTf₂ (20 mol %), AgOAc (10 mol %) in HFIP (1.0 M) at 80 °C for 24 h unless otherwise noted. The indicated yields are isolated yields after silica gel column chromatography. ^bUsing DCE instead of HFIP as solvent.

used DCE as a solvent was performed for **4d**. As expected, a lower yield was obtained compared with that in HFIP (34% in DCE vs 67% in HFIP).

Synthetically useful Weinreb amide²⁹ **6** was also allylated in moderate yield (eq 1). It is noteworthy that the Weinreb



amide-directed C–H bond functionalization reaction is rather limited possibly due to its weak coordinating ability,³⁰ and this is the first example of Weinreb amide-directed C–H bond functionalization under high-valent cobalt catalysis.

Figure 1 shows a plausible catalytic cycle for the allylation of 6-arylpyrines **1** based on previous Cp*Rh^{III}-^{7b} and




Figure 1. Plausible catalytic cycle for allylation of **1**.

Cp*Co^{III}-catalyzed^{17–20} allylation reactions, and C–H functionalization reactions of **1**.²³ The active catalyst would be coordinatively unsaturated cationic complex **I** generated from [Cp*Co(CO)I₂], AgOTf, and AgOAc. Coordination of **1** (**II**) followed by acetate-assisted C–H metalation would afford metallacycle intermediate **III**. After insertion of allyl alcohol **2** (**IV**), β -hydroxy elimination would release product **3** and hydroxide complex **V**,³¹ which would undergo protonation by AcOH to regenerate **I**. We observed trace amounts of aldehyde **8** as a byproduct in the reaction of **1b**. This byproduct supports the intermediate **IV**, from which β -hydride elimination to afford **8** is unfavorable but possible.²¹ The reason for the observed notable solvent effect is unclear, but the high polarity, moderate acidity, and weak coordinating ability of TFE and HFIP may facilitate the dissociation of acetate or other ligands and promote the generation of coordinatively unsaturated active species such as **I** and **III**. Coordination of acetate or another substrate **1** to these species could interfere with the desired reaction pathway.

In summary, efficient and general reaction conditions for Cp*Co^{III}-catalyzed aromatic C–H allylation of 6-arylpyrines, and benzamide derivatives using allyl alcohols as allylating reagents were developed. Use of fluorinated alcohol solvents was crucial for improving the scope of the dehydrative C–H allylation. We also describe our preliminary result of allylation of an aromatic Weinreb amide to further demonstrate the possibility of Cp*Co^{III}-catalyzed C–H functionalization reactions for the production of synthetically useful compounds.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b00846.

Experimental procedures, characterization data, and copy of NMR spectrum (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: tyoshino@pharm.hokudai.ac.jp.

*E-mail: smatsuna@pharm.hokudai.ac.jp.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported in part by ACT-C from JST, Scientific Research on Innovative Area from MEXT, Grant-in-Aid for Research Activity Start-up from JSPS, and Asahi Glass Foundation. Y.S. thanks JSPS for the fellowship.

■ REFERENCES

- (1) Trost, B. M. *Science* **1991**, 254, 1471.
- (2) Wender, P. A.; Miller, B. L. *Nature* **2009**, 460, 197.
- (3) Selected recent reviews and accounts for transition-metal-catalyzed C–H bond functionalization: (a) Ackermann, L. *Org. Process Res. Dev.* **2015**, 19, 260. (b) Rouquet, G.; Chatani, N. *Angew. Chem., Int. Ed.* **2013**, 52, 11726. (c) Wencel-Delord, J.; Glorius, F. *Nat. Chem.* **2013**, 5, 369. (d) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem., Int. Ed.* **2012**, 51, 8960. (e) Neufeldt, S. R.; Sanford, M. S. *Acc. Chem. Res.* **2012**, 45, 936. (f) Engle, K. M.; Mei, T.-S.; Wasa, M.; Yu, J.-Q. *Acc. Chem. Res.* **2012**, 45, 788. (g) Cho, S. H.; Kim, J. Y.; Kwak, J.; Chang, S. *Chem. Soc. Rev.* **2011**, 40, 5068.
- (4) (a) Magid, R. M. *Tetrahedron* **1980**, 36, 1901. (b) Pigge, F. C. *Synthesis* **2010**, 2010, 1745. Recent selected examples: (c) Farmer, J. L.; Hunter, H. N.; Organ, M. G. *J. Am. Chem. Soc.* **2012**, 134, 17470. (d) Yang, Y.; Buchwald, S. L. *J. Am. Chem. Soc.* **2013**, 135, 10642 and references therein.
- (5) Although direct allylation of aromatic C–H bonds is possible via Friedel–Crafts allylation, low regioselectivity, over-allylation, and limited substrate scope are often problematic: Rueping, M.; Nachtsheim, B. J. *Beilstein J. Org. Chem.* **2010**, 6, doi: 10.3762/bjoc.6.6.
- (6) (a) Oi, S.; Tanaka, Y.; Inoue, Y. *Organometallics* **2006**, 25, 4773. (b) Kuninobu, Y.; Ohta, K.; Takai, K. *Chem. Commun.* **2011**, 47, 10791. (c) Feng, C.; Feng, D.; Loh, T.-P. *Chem. Commun.* **2015**, 51, 342.
- (7) (a) Fan, S.; Chen, F.; Zhang, X. *Angew. Chem., Int. Ed.* **2011**, 50, 5918. (b) Wang, H.; Schröder, N.; Glorius, F. *Angew. Chem., Int. Ed.* **2013**, 52, 5386. (c) Zhang, S.-S.; Wu, J.-Q.; Lao, Y.-X.; Liu, X.-G.; Liu, Y.; Lv, W.-X.; Tan, D.-H.; Zeng, Y.-F.; Wang, H. *Org. Lett.* **2014**, 16, 6412. (d) Sharma, S.; Shin, Y.; Mishra, N. K.; Park, J.; Han, S.; Jeong, T.; Oh, Y.; Lee, Y.; Choi, M.; Kim, I. S. *Tetrahedron* **2015**, 71, 2435.
- (8) (a) Yao, T.; Hirano, K.; Satoh, T.; Miura, M. *Angew. Chem., Int. Ed.* **2011**, 50, 2990. (b) Makida, Y.; Ohmiya, H.; Sawamura, M. *Angew. Chem., Int. Ed.* **2012**, 51, 4122. (c) Cong, X.; Li, Y.; Wei, Y.; Zeng, X. *Org. Lett.* **2014**, 16, 3926. (d) Jo, H.; Han, S.; Park, J.; Choi, M.; Han, S. H.;

Jeong, T.; Lee, S.-Y.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *Tetrahedron* **2016**, *72*, 571.

(9) (a) Aihara, Y.; Chatani, N. *J. Am. Chem. Soc.* **2013**, *135*, 5308. (b) Yu, Y.-B.; Fan, S.; Zhang, X. *Chem. - Eur. J.* **2012**, *18*, 14643. (c) Xie, W.; Chang, S. *Angew. Chem., Int. Ed.* **2016**, *55*, 1876. (d) Dai, H.; Yu, C.; Lu, C.; Yan, H. *Eur. J. Org. Chem.* **2016**, 2016, 1255.

(10) (a) Asako, S.; Ilies, L.; Nakamura, E. *J. Am. Chem. Soc.* **2013**, *135*, 17755. (b) Yu, S.; Li, X. *Org. Lett.* **2014**, *16*, 1200.

(11) Reviews and accounts for Cp*Rh^{III}-catalyzed C–H bond functionalization: (a) Satoh, T.; Miura, M. *Chem. - Eur. J.* **2010**, *16*, 11212. (b) Patureau, F. W.; Wencel-Delord, J.; Glorius, F. *Aldrichimica Acta* **2012**, *45*, 31. (c) Song, G.; Wang, F.; Li, X. *Chem. Soc. Rev.* **2012**, *41*, 3651. (d) Chiba, S. *Chem. Lett.* **2012**, *41*, 1554. (e) Kuhl, N.; Schröder, N.; Glorius, F. *Adv. Synth. Catal.* **2014**, *356*, 1443. (f) Song, G.; Li, X. *Acc. Chem. Res.* **2015**, *48*, 1007. (g) Ye, B.; Cramer, N. *Acc. Chem. Res.* **2015**, *48*, 1308.

(12) Cp*Rh-catalyzed C–H allylation using allenes: (a) Zeng, R.; Fu, C.; Ma, S. *J. Am. Chem. Soc.* **2012**, *134*, 9597. (b) Ye, B.; Cramer, N. *J. Am. Chem. Soc.* **2013**, *135*, 636.

(13) (a) Yoshino, T.; Ikemoto, H.; Matsunaga, S.; Kanai, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 2207. (b) Ikemoto, H.; Yoshino, T.; Sakata, K.; Matsunaga, S.; Kanai, M. *J. Am. Chem. Soc.* **2014**, *136*, 5424. (c) Sun, B.; Yoshino, T.; Matsunaga, S.; Kanai, M. *Adv. Synth. Catal.* **2014**, *356*, 1491. (d) Sun, B.; Yoshino, T.; Kanai, M.; Matsunaga, S. *Angew. Chem., Int. Ed.* **2015**, *54*, 12968.

(14) Recent review: Moselage, M.; Li, J.; Ackermann, L. *ACS Catal.* **2016**, *6*, 498.

(15) Selected examples: (a) Hummel, J. R.; Ellman, J. A. *J. Am. Chem. Soc.* **2015**, *137*, 490. (b) Patel, P.; Chang, S. *ACS Catal.* **2015**, *5*, 853. (c) Li, J.; Ackermann, L. *Angew. Chem., Int. Ed.* **2015**, *54*, 3635. (d) Zhao, D.; Kim, J. H.; Stegemann, L.; Strasser, C. A.; Glorius, F. *Angew. Chem., Int. Ed.* **2015**, *54*, 4508. (e) Zhang, Z.-Z.; Liu, B.; Wang, C.-Y.; Shi, B.-F. *Org. Lett.* **2015**, *17*, 4094. (f) Sauermaann, N.; González, M. J.; Ackermann, L. *Org. Lett.* **2015**, *17*, 5316. (g) Park, J.; Chang, S. *Angew. Chem., Int. Ed.* **2015**, *54*, 14103. (h) Sen, M.; Kalsi, D.; Sundararaju, B. *Chem. - Eur. J.* **2015**, *21*, 15529. (i) Barsu, N.; Sen, M.; Premkumar, J. R.; Sundararaju, B. *Chem. Commun.* **2016**, *52*, 1338. (j) Prakash, S.; Muralirajan, K.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2016**, *55*, 1844. (k) Lerchen, A.; Vázquez-Céspedes, S.; Glorius, F. *Angew. Chem., Int. Ed.* **2016**, *55*, 3208. (l) Liang, Y.; Jiao, N. *Angew. Chem., Int. Ed.* **2016**, *55*, 4035. High-valent cobalt catalysis using bidentate directing groups: (m) Grigorjeva, L.; Daugulis, O. *Angew. Chem., Int. Ed.* **2014**, *53*, 10209. (n) Wu, X.; Yang, K.; Zhao, Y.; Sun, H.; Li, G.; Ge, H. *Nat. Commun.* **2015**, *6*, 6462. (o) Zhang, J.; Chen, H.; Lin, C.; Liu, Z.; Wang, C.; Zhang, Y. *J. Am. Chem. Soc.* **2015**, *137*, 12990. (p) Kalsi, D.; Sundararaju, B. *Org. Lett.* **2015**, *17*, 6118. (q) Gandeepan, P.; Rajamalli, P.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2016**, *55*, 4308.

(16) Reviews on inexpensive first-row transition metal catalyzed C–H bond functionalization: (a) Kulkarni, A. A.; Daugulis, O. *Synthesis* **2009**, 4087. (b) Yamaguchi, J.; Muto, K.; Itami, K. *Eur. J. Org. Chem.* **2013**, 2013, 19. (c) Yoshikai, N. *Yuki Gosei Kagaku Kyokaiishi* **2014**, *72*, 1198.

(17) Yu, D.-G.; Gensch, T.; de Azambuja, F.; Vázquez-Céspedes, S.; Glorius, F. *J. Am. Chem. Soc.* **2014**, *136*, 17722.

(18) Moselage, M.; Sauermaann, N.; Koeller, J.; Liu, W.; Gelman, D.; Ackermann, L. *Synlett* **2015**, 26, 1596.

(19) Gensch, T.; Vázquez-Céspedes, S.; Yu, D.-G.; Glorius, F. *Org. Lett.* **2015**, *17*, 3714.

(20) Suzuki, Y.; Sun, B.; Sakata, K.; Yoshino, T.; Matsunaga, S.; Kanai, M. *Angew. Chem., Int. Ed.* **2015**, *54*, 9944.

(21) Under Cp*Rh^{III}-catalyzed conditions with a stoichiometric oxidant, the β -hydride elimination pathway is dominant: Shi, Z.; Bouladakis-Arapinis, M.; Glorius, F. *Chem. Commun.* **2013**, 49, 6489.

(22) Kumar, G. S.; Kapur, M. *Org. Lett.* **2016**, *18*, 1112.

(23) (a) Guo, H.-M.; Jiang, L.-L.; Niu, H.-Y.; Rao, W.-H.; Liang, L.; Mao, R.-Z.; Li, D.-Y.; Qu, G.-R. *Org. Lett.* **2011**, *13*, 2008. (b) Guo, H.-M.; Rao, W.-H.; Niu, H.-Y.; Jiang, L.-L.; Meng, G.; Jin, J.-J.; Yang, X.-N.; Qu, G.-R. *Chem. Commun.* **2011**, 47, 5608. (c) Lakshman, M. K.; Deb, A. C.; Chamala, R. R.; Pradhan, P.; Pratap, R. *Angew. Chem., Int. Ed.* **2011**, *50*, 11400. (d) Chamala, R. R.; Parrish, D.; Pradhan, P.; Lakshman, M. K.

J. Org. Chem. **2013**, *78*, 7423. (e) Kim, H. J.; Ajitha, M. J.; Lee, Y.; Ryu, J.; Kim, J.; Lee, Y.; Jung, Y.; Chang, S. *J. Am. Chem. Soc.* **2014**, *136*, 1132.

(f) Patel, P.; Chang, S. *ACS Catal.* **2015**, *5*, 853. (g) Pawar, A. B.; Chang, S. *Org. Lett.* **2015**, *17*, 660. (h) Liang, Y.; Liang, Y.-F.; Tang, C.; Yuan, Y.; Jiao, N. *Chem. - Eur. J.* **2015**, *21*, 16395. (i) Wang, S.; Hou, J.-T.; Feng, M.-L.; Zhang, X.-Z.; Chen, S.-Y.; Yu, X.-Q. *Chem. Commun.* **2016**, *52*, 2709. (j) Pawar, A. B.; Lade, D. M. *Org. Biomol. Chem.* **2016**, *14*, 3275.

(24) (a) Hocek, M.; Holý, A.; Votruba, I.; Dvořáková, H. *J. Med. Chem.* **2000**, *43*, 1817. (b) Gundersen, L.-L.; Nissen-Meyer, J.; Spilsberg, B. *J. Med. Chem.* **2002**, *45*, 1383. (c) Bakkestuen, A. K.; Gundersen, L.-L.; Utenova, B. T. *J. Med. Chem.* **2005**, *48*, 2710. (d) Hocek, M.; Nauš, P.; Pohl, R.; Votruba, I.; Furman, P. A.; Tharnish, P. M.; Otto, M. J. *J. Med. Chem.* **2005**, *48*, 5869.

(25) The similar high reactivity in the fluorinated alcohol solvents was observed in several reactions under high-valent cobalt catalysis. See refs 15d–f, h, i, k, m, p, q, and 23j.

(26) (a) Lapointe, D.; Fagnou, K. *Chem. Lett.* **2010**, *39*, 1118. (b) Ackermann, L. *Chem. Rev.* **2011**, *111*, 1315 and references therein.

(27) See Table S1 in the [Supporting Information](#) for more details.

(28) Bis-allylated byproducts were observed in 17% yield for **4a** and 19% yield for **4b** by ¹H NMR analysis of the crude mixture.

(29) (a) Nahm, S.; Weinreb, S. M. *Tetrahedron Lett.* **1981**, *22*, 3815.

(b) Balasubramaniam, S.; Aidhen, I. S. *Synthesis* **2008**, 2008, 3707 and references therein.

(30) Examples of Weinreb amide directed C–H functionalization: (a) Yang, F.; Ackermann, L. *Org. Lett.* **2013**, *15*, 718. (b) Wang, Y.; Li, C.; Li, Y.; Yin, F.; Wang, X.-S. *Adv. Synth. Catal.* **2013**, *355*, 1724. (c) Davis, T. A.; Hyster, T. K.; Rovis, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 14181. (d) Wang, Y.; Zhou, K.; Lan, Q.; Wang, X.-S. *Org. Biomol. Chem.* **2015**, *13*, 353.

(31) β -Hydroxy elimination was proposed in some transition-metal-catalyzed reactions: (a) Walters, A. J. C.; Troeppner, O.; Ivanović-Burmazović, I.; Tejel, C.; del Río, M. P.; de Bruin, B. *Angew. Chem., Int. Ed.* **2012**, *51*, 5157. (b) Gumrukcu, Y.; de Bruin, B.; Reek, J. N. H. *Chem. - Eur. J.* **2014**, *20*, 10905.