

## Correction to "Metal-Free Remote C—H Bond Amidation of 8-amidoquinolines on the C5 Position under Mild Conditions"

Dezhong Ji, Xin He, Yizhu Xu, Ziyue Xu, Yuchen Bian, Weixing Liu, Qihua Zhu,\* and Yungen Xu\* Org. Lett. 2016, 18 (18), 4478–4481. DOI: 10.1021/acs.orglett.6b01980

## Supporting Information

The solvent used in the reaction was incorrectly identified throughout this paper. The correct solvent used in the reaction is glycol dimethyl ether (DME). The corrections for the solvent changes are provided below:

The solvent dimethyl ether should be changed to glycol dimethyl ether (DME) in the footnotes of Schemes 2 and Scheme 3.

The Table of Contents graphic is corrected as follows:

The graphic in Scheme 2 is corrected as follows:

$$\begin{array}{c} R_2 \\ \\ N \\ \\ HN \\ R_1 \end{array} + NH(SO_2Ph)_2 \\ \\ \frac{Phl(OAc)_2}{glycol\ dimethyl\ ether} \\ R_2 \\ \\ N \\ \\ N \\ R_1 \end{array}$$

The graphic in Scheme 2 is corrected as follows:

## ASSOCIATED CONTENT

## **S** Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b02867.

Revised file with the correct solvent listed in the Experimental Section on page S2 (PDF)

Received: October 7, 2016

Published: October 24, 2016