

Aryl Ketone Catalyzed Radical Allylation of C(sp³)—H Bonds under Photoirradiation

Shin Kamijo,*,† [6] Kaori Kamijo,† Kiyotaka Maruoka,‡ and Toshihiro Murafuji

[†]Graduate School of Sciences and Technology for Innovation and [‡]Department of Biology and Chemistry, Yamaguchi University, Yamaguchi 753-8512, Japan

Supporting Information

ABSTRACT: The catalytic introduction of an allyl group at nonacidic $C(sp^3)$ –H bonds was achieved under photoirradiation, in which 1,2-bis(phenylsulfonyl)-2-propene acts as an allyl source and 5,7,12,14-pentacenetetrone (PT) works as a C–H bond-cleaving catalyst. A variety of substances, including alkanes, carbamates, ethers, sulfides, and alcohols, were chemoselectively allylated in a single step under neutral conditions. The present transformation is catalyzed solely by an organic molecule, PT, and

proceeds smoothly even under visible light irradiation (425 nm) in the case of alkanes as a starting substance.

he allyl group is one of the most important carbon units in organic chemistry because of its synthetic versatility. For instance, the allyl group provides multiple opportunities for further bond-forming transformations since the olefinic moiety can be easily converted to electrophilic functional groups, such as aldehydes and epoxides, by oxidative manipulations. The synthetic utility of the allyl group has been further accentuated by the emergence of metathesis, the formation of a new olefinic product by the recombination of two olefins.² For this reason, intensive investigations have been carried out on allylation reactions from a wide array of aspects, such as the development of new allylating agents, stererochemical control of allylation, and catalytic allylation as well as their mechanistic investigations. Among the substitutive introductions of the allyl group at C(sp³)-H bonds, the majority of methods involve anionic allylation after generation of a carbanion by making use of the acidic nature of C-H bonds adjacent to electron-withdrawing groups (E, Scheme 1-1). LDA and LHMDS are representative bases for deprotonation to generate the carbanion (Scheme 1-1a),³ and an allyl carbonate-Pd(0) catalyst is a standard combination for allylation of active methine/methylene moieties (Scheme 1-1b). Despite numerous examples for allylation of acidic $C(sp^3)$ -H bonds, the methods for substitutive introduction of an allyl group at nonacidic C(sp³)-H bonds are still underdeveloped.5,

We therefore investigated allylation of unreactive $C(sp^3)$ —H bonds by applying a photochemical C—H functionalization strategy (Scheme 1-2).^{7,8} The keys to realizing such a transformation are to find a suitable aryl ketone as a precursor for an oxyl radical species to cleave the C—H bonds, and the design of an allyl source to trap the in situ generated carbon radical A. We herein report the photoinduced catalytic allylation of nonacidic $C(sp^3)$ —H bonds, in which 1,2-bis(phenylsulfonyl)-2-propene acts as the allyl source and 5,7,12,14-pentacenetetrone (PT) works as the C—H bond-cleaving catalyst. The characteristic features of the present transformation are that the reaction is

Scheme 1. Strategies for Substitutive Allylation of $C(sp^3)-H$ Bonds

1. anionic allylation [acidic C(sp3)-H bond]

(a) base-promoted allylation

(b) transition metal-catalyzed allylation

$$\begin{array}{c|c} & OCO_2R \\ \hline E & C \\ \hline C & H \\ \hline & RO-H \\ \hline \end{array} \begin{array}{c} E \\ \hline C \\ \hline & Pd(0) \\ \hline \end{array} \begin{array}{c} E \\ \hline & Pd(0) \\ \hline \end{array} \begin{array}{c} E \\ \hline & Pd(0) \\ \hline \end{array}$$

2. radical allylation [non-acidic C(sp3)-H bond] this work

catalyzed solely by the organic molecule, PT, and the allylation of alkanes smoothly proceeds even under irradiation of visible light at 425 nm wavelength.

At the outset of our investigation, we screened aryl ketones that are potentially effective for C–H bond cleavage of cyclooctane with employment of 1,2-bis(phenylsulfonyl)-2-propene as the allyl source 9,10 (Table 1). The reaction was catalyzed by benzophenone (Ph₂CO, 10 mol %), and the allylation product 2a was obtained in 52% yield (entry 1). This is a rare example of a photoinduced substitutive C–H function-

Received: December 1, 2016

Published: December 8, 2016

Organic Letters Letter

Table 1. Optimization of Reaction Conditions

entry	aryl ketone	mol %	h $ u$, nm	solvent	yield, % ^{b,c}
1	Ph_2CO	10	365	MeCN	52 ^d
2	4-BzPy	10	365	MeCN	30^d
3	2-ClAQ	10	365	MeCN	61
4	PT	10	365	MeCN	58 ^d
5	PT	10	425	MeCN	55
6	PT	10	455	MeCN	34 ^d
7	PT	10	425	CH_2Cl_2	59
8	PT	5	425	CH_2Cl_2	55
9 ^e	PT	5	425	CH_2Cl_2	60
10 ^e	PT	5	365	CH_2Cl_2	63
11 ^e	PT	1	425	CH_2Cl_2	38

^aConditions: cyclooctane **1a** (10 equiv), allyl source (1 equiv), aryl ketone (1–10 mol %), solvent (0.05 M), photoirradiation using an LED lamp at rt for 24 h. ^bIsolated yields unless otherwise noted. ^cRecovery of the allyl source was observed in general. ^dYield based on ¹H NMR analysis of the crude mixture. ^eThe reaction was conducted with addition of K_2CO_3 (1.1 equiv).

alization that was efficiently promoted even with a catalytic amount of aryl ketone. The use of 4-benzoylpyridine (4-BzPy)^{7h,i,11} reduced the product yield (30%, entry 2), and that of quinone-based ketones, 2-chloroanthraquinone (2-ClAQ) and PT, improved the yield (61% and 58% yield, entries 3 and 4). The application of PT enabled the use of visible light at 425 nm wavelength, and 2a was formed in practically the same yield as observed by the reaction under 365 nm light irradiation (entry 5). 12 The use of 455 nm light, however, appeared to retard the reaction progress (entry 6). A halogenated solvent, CH₂Cl₂, slightly improved the solubility of PT (entry 7), and its amount could be reduced (5 mol %, entry 8). Turther investigations revealed that the addition of K2CO3 slightly increased the yield of 2a (60%, entry 9). 14 The irradiation of 365 nm light again gave almost the same yield of 2a under the optimized conditions (63%, entry 10), whereas the reaction with 1 mol % PT decreased the product yield (38%, entry 11).

We then explored the applicability of the catalytic allylation of nonacidic C-H bonds by treating various alkanes 1b-j with the allyl source (1 equiv) in the presence of 5 mol % PT and 1.1 equiv of K₂CO₃ under photoirradiation in CH₂Cl₂ (Scheme 2). The reactions of cyclohexane 1b and cyclododecane 1c using 425 nm light afforded the corresponding products 2b and 2c in 69% and 58% yield, respectively. Irradiation of 365 nm light promoted the allylation as well for these cycloalkanes (63% and 42% yield, respectively). The reaction of adamantane 1d predominantly provided the 1-allyladamantane 2d along with a minute amount of 2-allyladamantane **2d**′. ¹⁵ This result clarified that the reactivity of the methine C-H bond is higher than that of the methylene one. The reaction of functionalized adamantanes 1e and 1f, with tosylamide and hydroxy groups, proceeded selectively at the methine carbon to furnish the respective adducts 2e and 2f. It is noteworthy that allylation at the amide and hydroxy functional groups was not detected at all under the present conditions. The benzoylated adamantanol 1g and the adamantanemethanol derivative 1h furnished the expected allylation products 2g and 2h. The allylation of adamantaneethanol 1i took place at the adamantane core as well. The acyclic compound 1j was chemoselectively allylated at the methine C-H bond to furnish

Scheme 2. Allylation of Alkanes^{a,b}

"Conditions: alkane 1 (5 equiv), allyl source (1 equiv), PT (5 mol %), K_2CO_3 (1.1 equiv), CH_2Cl_2 (0.05 M), photoirradiation using a 425 nm LED lamp at rt for 24 h unless otherwise noted. ^bIsolated yield. The alkanes 1b and 1d (10 equiv) were used. ^dIsolated yield of reaction using a 365 nm LED lamp. ^eMinute amounts of the methylene adducts were observed (<4%). ^fThe reaction was conducted without addition of K_2CO_3 . ^gThe reaction was conducted using 1f (1 equiv) and the allyl source (1.2 equiv). ^hYield based on ¹H NMR analysis of the crude mixture.

the product 2j in 31% yield. Although the yield of 2j remained moderate, an alternative method for one-step conversion of 1j to 2j is not readily available at the moment.

We next examined the catalytic allylation of heteroatomcontaining substances 1k-s (Scheme 3). The reaction of

Scheme 3. Allylation of Heteroatom-Containing Substances a,b

^aConditions: starting substance 1 (5 equiv), allyl source (1 equiv), PT (5 mol %), K₂CO₃ (1.1 equiv), CH₂Cl₂ (0.05 M), photoirradiation using a 365 nm LED lamp at rt for 24 h unless otherwise noted. ^bIsolated yield. ^cThe heterocycle 1k or 1p (10 equiv) was used. ^dIrradiated using a 425 nm LED lamp. ^eThe reaction using a 425 nm LED lamp gave <26% yield based on ¹H NMR analysis of the crude mixture.

tetrahydrofuran (THF) 1k revealed that the allyl group was chemoselectively introduced to the C-center adjacent to the O-atom to form the product 2k in 64% yield with 365 nm light and 73% yield with 425 nm light. The allylation of ambroxide 1l proceeded with the same chemoselectivity without touching potentially reactive methine C—H bonds. In the case of allylation

Organic Letters Letter

of *N*-Boc pyrrolidine **1m**, the irradiation of 365 nm light (76%) was more effective than that of 425 nm light (<26% yield determined by NMR analysis). The allylation of the morpholine derivative **1n** clarified that the C–H bond proximal to the *N*-Boc group has higher reactivity than that next to the O-atom (2n). The acyclic starting material 10 derived from butylamine furnished the expected adduct 20 as well. The cyclic sulfides, such as tetrahydrothiophene 1p and thioxane 1q, were allylated at the C-H bond adjacent to the S-atom to produce the corresponding adducts 2p (81%) and 2q (57%). The allylation of cyclohexanol 1r took place at the hydroxylated C-center to give the tertiary homoallyl alcohol 2r. The construction of a tetrasubstituted C-center via selective allylation at the most sterically congested methine C-H bond is of note. The allyl group was similarly introduced to the primary alcohol 1s to provide the secondary alcohol 2s. This result illustrates that the reacting C-H bond can be controlled by the proper choice of alcohol protecting group (see Scheme 2, $1j \rightarrow 2j$).

Having succeeded in the photoinduced catalytic substitutive introduction of an allyl group to a variety of substances, we turned our attention to obtaining mechanistic information on the present transformation. We conducted the allylation of cyclooctane 1a in the presence of TEMPO (see Supporting Information for details). The analysis of the crude mixture revealed the formation of the TEMPO adduct 3a along with a trace amount of the allylated product 2a and recovery of the allyl source. This result strongly indicated the generation of carbon radical A during the course of the reaction (Schemes 1 and 4).

Scheme 4. Proposed Mechanism for the Photoinduced Catalytic Radical Allylation of Nonacidic C(sp³)—H Bonds

We then measured the kinetic isotope effect by treating a mixture of cyclohexane ${\bf 1b}$ and its deuterated analogue ${\bf 1b}$ -d with the allyl source under the standard conditions (see Supporting Information for details). The value of the kinetic isotope effect (KIE) was determined to be 3.6, indicating that $C(sp^3)$ -H bond cleavage is involved in the rate-determining step.

Taking all this information into account, we propose a reaction mechanism as illustrated in Scheme 4. The photoexcited aryl ketone, PT, having biradical reactivity, abstracts a hydrogen from the starting substance 1 to generate the carbon radical intermediate $\bf A$ and the semiquinone-type radical $\bf B$. The generation of the carbon radical $\bf A$ was strongly supported by the TEMPO trap (3a), and this step was suggested to be the rate-

determining step based on the KIE experiment. The derived radical A then adds to the electron-deficient vinyl sulfone moiety of the allyl source to furnish the radical C.^{7e,j} Subsequent elimination of the sulfonyl radical furnishes the allylated product 2, and the released sulfonyl radical most probably accepts the hydrogen from B to regenerate PT, ¹⁸ completing the catalytic cycle for the present substitutive allylation of nonacidic C–H bonds.

To highlight the superiority of the present one-step allylation of nonacidic $C(sp^3)$ -H bonds for rapid assembly of several molecules and for construction of a cyclic molecular framework, we performed the following two transformations (Scheme 5).

Scheme 5. Transformations of Allylated Products

NHTs
$$Ph_2CO (1 \text{ equiv})$$
 $NHTs OH$ SO_2Ph $PrOH (0.05 M)$ SO_2Ph $PrOH (0.05 M)$ SO_2Ph $PrOH (0.05 M)$ SO_2Ph $Ph_2CO (1 \text{ equiv})$ SO_2Ph SO_2Ph

The first one is the reaction of the allylated adduct **2e** with isopropyl alcohol by the photoinduced Michael-type radical addition (**3e**). Thus, two different molecules could be successfully linked with 1,2-bis(phenylsulfonyl)-2-propene as a tethering precursor utilizing the two different types of photoinduced C—H functionalization reactions. The second one is the preparation of azacycles **3o** and **4o**²⁰ from acyclic homoallylamine **2o**. The pyrrolidine core could be constructed in merely two steps from *N*-Boc butylamide **1o** as a starting substance.

In conclusion, we have developed the photoinduced catalytic allylation of nonacidic C(sp³)-H bonds of a variety of substances, including alkanes, carbamates, ethers, sulfides, and alcohols, in a chemoselective manner. The reaction is proposed to proceed through a radical mechanism, in which the homolytic cleavage of nonacidic C(sp³)-H bonds is effected solely by photoexcited 5,7,12,14-pentacenetetrone (PT), and the allyl group is delivered from 1,2-bis(phenylsulfonyl)-2-propene. In this study, we showed for the first time that (1) the organic molecule, PT, acts as an excellent C-H bond cleaving agent; (2) the direct allylation of nonacidic C(sp³)-H bonds can be efficiently catalyzed by PT; and (3) PT allows the use of visible light (425 nm) for the functionalization of alkanes. The newly developed radical allylation of nonacidic C(sp³)-H bonds complements conventional methods for anionic introduction of an allyl group at acidic $C(sp^3)$ —H bonds under basic conditions. In addition, the present C-H allylation provides rapid access to structurally complex substances by taking advantage of the electrophilic nature of the installed vinyl sulfone moiety for further transformations. The protocol developed in this study, therefore, should serve as a unique tool for carbon chain extension stemming from nonacidic $C(sp^3)$ -H bonds.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b03586.

Additional experimental details (PDF)

Organic Letters Letter

AUTHOR INFORMATION

Corresponding Author

*E-mail: kamijo@yamaguchi-u.ac.jp.

ORCID ®

Shin Kamijo: 0000-0002-5662-5371

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This research was supported by the Program to Disseminate Tenure Tracking System (MEXT, Japan) and the research grant of Astellas Foundation for Research on Metabolic Disorders. We thank Prof. Masayuki Inoue and Dr. Yuuki Amaoka at The University of Tokyo for valuable suggestions on design of the allyl source. We also acknowledge Prof. Masahiro Terada at Tohoku University for continuous encouragement.

REFERENCES

- (1) For selected reviews on allylation, see: (a) Yamamoto, Y.; Asao, N. Chem. Rev. 1993, 93, 2207. (b) Yus, M.; González-Gómez, J. C.; Foubelo, F. Chem. Rev. 2013, 113, 5595.
- (2) For selected reviews on olefin metathesis, see: (a) *Handbook of Metathesis*; Grubbs, R. H., Wenzel, A. G., O'Leary, D. J., Khosravi, E., Eds.; Wiley-VCH: Weinheim, 2005. (b) Hoveyda, A. H.; Malcolmson, S. J.; Meek, S. J.; Zhugralin, A. R. *Angew. Chem., Int. Ed.* **2010**, 49, 34. (c) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. *Angew. Chem., Int. Ed.* **2005**, 44, 4490.
- (3) For a book Chapter on alkylation of enols and enolates, see: Stoltz, B. M.; Bennett, N. B.; Duquette, D. C.; Goldberg, A. F. G.; Liu, Y.; Loewinger, M. B.; Reeves, C. M. In *Comprehensive Organic Synthesis*, 2nd ed.; Knochel, P., Molander, G. A., Eds.; Elsevier: Amsterdam, 2014; Vol. 3, pp 1–55.
- (4) For selected reviews on π -allylpalladium chemistry, see: (a) Tsuji, J. *Palladium Reagents and Catalysts*; Wiley: Chichester, 2004; Chapter 4, pp 431–517. (b) Trost, B. M.; Crawley, M. L. *Chem. Rev.* **2003**, *103*, 2921.
- (5) For pioneering works on the allylation of nonacidic $C(sp^3)$ -H bonds, see: (a) Xiang, J.; Evarts, J.; Rivkin, A.; Curran, D. P.; Fuchs, P. L. *Tetrahedron Lett.* **1998**, 39, 4163. (b) Tanko, J. M.; Sadeghipour, M. *Angew. Chem., Int. Ed.* **1999**, 38, 159. (c) Zhang, J.; Li, Y.; Zhang, F.; Hu, C.; Chen, Y. *Angew. Chem., Int. Ed.* **2016**, 55, 1872.
- (6) For recent representative reports on the C(sp³)—H functionalization using a tungsten photocatalyst, see: (a) Ravelli, D.; Protti, S.; Fagnoni, M. Acc. Chem. Res. 2016, 49, 2232. (b) Murphy, J. J.; Bastida, D.; Paria, S.; Fagnoni, M.; Melchiorre, P. Nature 2016, 532, 218. (c) Yamada, K.; Okada, M.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Org. Lett. 2015, 17, 1292. (d) Okada, M.; Fukuyama, T.; Yamada, K.; Ryu, I.; Ravelli, D.; Fagnoni, M. Chem. Sci. 2014, 5, 2893.
- (7) (a) Kamijo, S.; Hoshikawa, T.; Inoue, M. Tetrahedron Lett. 2011, 52, 2885. (b) Kamijo, S.; Hoshikawa, T.; Inoue, M. Org. Lett. 2011, 13, 5928. (c) Hoshikawa, T.; Yoshioka, S.; Kamijo, S.; Inoue, M. Synthesis 2013, 45, 874. (d) Hoshikawa, T.; Kamijo, S.; Inoue, M. Org. Biomol. Chem. 2013, 11, 164. (e) Amaoka, Y.; Nagatomo, M.; Watanabe, M.; Tao, K.; Kamijo, S.; Inoue, M. Chem. Sci. 2014, 5, 4339. (f) Kamijo, S.; Hirota, M.; Tao, K.; Watanabe, M.; Murafuji, T. Tetrahedron Lett. 2014, 55, 5551. (g) Kamijo, S.; Tao, K.; Takao, G.; Murooka, H.; Murafuji, T. Tetrahedron Lett. 2015, 56, 1904. (h) Kamijo, S.; Watanabe, M.; Kamijo, K.; Tao, K.; Murafuji, T. Synthesis 2016, 48, 115. (i) Kamijo, S.; Takao, G.; Kamijo, K.; Hirota, M.; Tao, K.; Murafuji, T. Angew. Chem., Int. Ed. 2016, 55, 9695. (j) Kamijo, S.; Takao, G.; Kamijo, K.; Tsuno, T.; Ishiguro, K.; Murafuji, T. Org. Lett. 2016, 18, 4912.
- (8) For related examples of C(sp³)—H functionalization using a photoexcited ketone, see: (a) Hoshikawa, T.; Inoue, M. *Chem. Sci.* **2013**, *4*, 3118. (b) Nagatomo, M.; Yoshioka, S.; Inoue, M. *Chem. Asian J.* **2015**, *10*, 120. (c) Xia, J.-B.; Zhu, C.; Chen, C. *J. Am. Chem. Soc.* **2013**,

- 135, 17494. (d) Xia, J.-B.; Zhu, C.; Chen, C. Chem. Commun. 2014, 50, 11701. (e) Cantillo, D.; de Frutos, O.; Rincón, J. A.; Mateos, C.; Kappe, C. O. J. Org. Chem. 2014, 79, 8486. (f) Ota, E.; Mikame, Y.; Hirai, G.; Nishiyama, S.; Sodeoka, M. Synlett 2016, 27, 1128. (g) Lipp, A.; Lahm, G.; Opatz, T. J. Org. Chem. 2016, 81, 4890. (h) For a recent review, see: Chen, C. Org. Biomol. Chem. 2016, 14, 8641.
- (9) For reports using allyl sulfones as an allyl source in radical chemistry, see: (a) Baldwin, J. E.; Adlington, R. M.; Birch, D. J.; Crawford, J. A.; Sweeney, J. B. J. Chem. Soc., Chem. Commun. 1986, 1339. (b) Padwa, A.; Murphree, S. S.; Yeske, P. E. Tetrahedron Lett. 1990, 31, 2983. (c) Kim, S.; Lim, C. J. Angew. Chem., Int. Ed. 2002, 41, 3265. (d) Lee, S.; Lim, C. J.; Kim, S. Bull. Korean Chem. Soc. 2004, 25, 1611. (e) Lee, J. Y.; Kim, S. Bull. Korean Chem. Soc. 2006, 27, 189. (f) Schaffner, A.-P.; Renaud, P. Angew. Chem., Int. Ed. 2003, 42, 2658. (g) Xu, G.; Lüthy, M.; Habegger, J.; Renaud, P. J. Org. Chem. 2016, 81, 1506. (h) De Dobbeleer, C.; Pospišil, J.; De Vleeschouwer, F.; De Proft, F.; Markó, I. E. Chem. Commun. 2009, 2142. (i) Quinn, R. K.; Schmidt, V. A.; Alexanian, E. J. Chem. Sci. 2013, 4, 4030. (j) Corcé, V.; Chamoreau, L.-M.; Derat, E.; Goddard, J.-P.; Ollivier, C.; Fensterbank, L. Angew. Chem., Int. Ed. 2015, 54, 11414. (k) Cui, L.; Chen, H.; Liu, C.; Li, C. Org. Lett. 2016, 18, 2188. (1) Heitz, D. R.; Rizwan, K.; Molander, G. A. J. Org. Chem. 2016, 81, 7308. (m) Qi, L.; Chen, Y. Angew. Chem., Int. Ed. 2016, 55, 3312.
- (10) For protocols of the allyl source preparation, see: (a) Padwa, A.; Kline, D. N.; Norman, B. H. *Tetrahedron Lett.* **1988**, 29, 265. (b) Padwa, A.; Kline, D. N.; Murphree, S. S.; Yeske, P. E. *J. Org. Chem.* **1992**, 57, 298. (c) Chang, M.-Y.; Wu, M.-H. *Synlett* **2014**, 25, 411.
- (11) Kamijo, S.; Tao, K.; Takao, G.; Tonoda, H.; Murafuji, T. *Org. Lett.* **2015**, *17*, 3326.
- (12) Baxter, I.; Cameron, D. W.; Titman, R. B. J. Chem. Soc. C 1971, 1253.
- (13) The reactions in acetone, benzene, and t-BuOH gave lower yields.
- (14) Inorganic salts, such as Li_2CO_3 , Na_2CO_3 , Cs_2CO_3 , and $CaCO_3$, as well as 2,6-di(tert-butyl)pyridine, gave similar results. Accordingly, it is presumed that these bases neutralize in situ generated PhSO₂H to suppress undesired side reactions. A similar base effect was observed in ref 8g.
- (15) 2-Allyladamantane **2d'** was prepared for unambiguous structure confirmation; see: (a) Wang, G.-Z.; Jiang, J.; Bu, X.-S.; Dai, J. J.; Xu, J.; Fu, Y.; Xu, H.-J. *Org. Lett.* **2015**, *17*, 3682. (b) Inomata, K.; Sasaoka, S.-i.; Kobayashi, T.; Tanaka, Y.; Igarashi, S.; Ohtani, T.; Kinoshita, H.; Kotake, H. *Bull. Chem. Soc. Jpn.* **1987**, *60*, 1767.
- (16) For a UVspectrum of PT, see: Baxter, I.; Cameron, D. W.; Titman, R. B. J. Chem. Soc. C 1971, 1253.
- (17) For the structure of the anion radical of PT, see: (a) Forkner, M. W.; Miller, L. L.; Rak, S. F. *Synth. Met.* **1990**, 36, 65. (b) Almlöf, J. E.; Feyereisen, M. W.; Jozefiak, T. H.; Miller, L. L. *J. Am. Chem. Soc.* **1990**, 112, 1206.
- (18) Although the possibility of a radical chain mechanism cannot be exclusively ruled out at present, we confirmed that the allylation took place only when the light was irradiated (see Supporting Information for details) and that the recovery of a majority of PT was observed after the reaction. More examinations are required to account for the precise reaction mechanism, see: Cismesia, M. A.; Yoon, T. P. Chem. Sci. 2015, 6, 5426.
- (19) Pioneering work on the photoinduced alkylation of isopropanol: Schenck, G. O.; Koltzenburg, G.; Grossmann, H. *Angew. Chem.* **1957**, *69*, 177.
- (20) Coldham, I.; Leonori, D. J. Org. Chem. 2010, 75, 4069.