

Copper-Catalyzed P–H Insertions of α -Imino Carbenes for the Preparation of 3-Phosphinoylindoles


Lisheng Wang,[†] Yi Wu,[†] Yang Liu,[†] Hua Yang,[†] Xu Liu,[†] Jianyi Wang,[†] Xiaoxun Li,[‡] and Jun Jiang^{*,†}

[†]School of Chemistry and Chemical Engineering, Guangxi University, Nanning 530004, P. R. China

[‡]Department of Chemistry, Stanford University, Stanford, California 94305-5580, United States

S Supporting Information

ABSTRACT: A highly efficient P–H insertion of α -imino copper carbene into H-phosphine oxides, leading to 3-phosphinoylindoles with a broad substrate diversity and good chemoselectivity, is established. This methodology provides a rapid and efficient approach for the C(sp²)–P bond formation via the P–H insertion strategy. The stereoselective P–H insertion of α -imino copper carbene represents a unique example of asymmetric P–H insertion, affording P-stereogenic 3-phosphinoylindoles with high stereoselectivities.




The highly efficient construction of carbon–heteroatom bonds is one of the most pursued goals in modern organic synthesis. Transition-metal-catalyzed insertion of carbenes and carbenoids generated in situ from α -diazo carbonyl compounds into heteroatom–hydrogen bonds (X–H, where X = O, N, S, P, etc.) is an efficient strategy for the construction of $\text{C}(\text{sp}^3)$ –X bonds.¹ Recently, α -imino rhodium carbene, generated in situ from *N*-sulfonyl 1,2,3-triazoles, has emerged as an increasingly valuable intermediate possessing novel reactivity and synthetic potential.² Significant progress has been made by the research groups of Fokin,³ Gevorgyan,⁴ Davies,⁵ Sarpong,⁶ Shi,⁷ and others. In particular, α -imino rhodium carbenes insertions have led to a novel synthetic method toward the formation of C–X bonds, including $\text{C}(\text{sp}^2)$ –X bonds. For example, high regio- and stereoselectivities have been achieved with the insertion of α -imino rhodium carbene into O–H and N–H bonds, affording enamide products (Scheme 1a).^{3a}

Indole and its derivatives are important bioactive heterocycles in natural products and pharmaceuticals.⁸ More recently, indole-embedded α -imino metal carbene in situ generated from 3-diazoindolin-2-imine has been identified as a reactive intermediate in carbene insertions to furnish a variety of indole derivatives.⁹ The transformation of rhodium-catalyzed N–H insertion was reported to build 3-substituted indole derivatives with the $\text{C}(\text{sp}^2)$ –N bond formation (Scheme 1b).^{9c} Despite progress in α -imino carbene insertion transformations, the insertion reactions of indole-embedded α -imino metal carbenes for the generation of $\text{C}(\text{sp}^2)$ –X bonds is much less developed.


Aryl and heteroaryl phosphorus compounds have been of particular interest due to their biological significance and synthetic value.¹⁰ 3-Phosphoindoles represent novel second-generation NNRTIs (non-nucleoside reverse transcriptase inhibitors) which show remarkable effect on inhibiting the replication of HIV-1 virus (IDX899).¹¹ In consideration of the

Scheme 1. Insertion Reactions of α -Imino Metal Carbenes for the Construction of $\text{C}(\text{sp}^2)$ –Heteroatom Bonds


a. O–H and N–H insertions of α -imino rhodium carbene



b. N–H insertion of indole-embedded α -imino rhodium carbene



c. P–H insertion of indole-embedded α -imino copper carbene (this work)




importance of 3-phosphoindoles in pharmaceuticals and inspired by these previous findings shown in Scheme 1, herein we describe our efforts toward the development of the copper-catalyzed P–H insertion of novel α -imino copper carbene to access a variety of 3-phosphoindole derivatives (Scheme 1c). This method will not only expand the repertoire of transformations of α -imino metal carbenes but also provide a straightforward route to $\text{C}(\text{sp}^2)$ –P bond formation and creation of new phosphoindoles. To the best of our knowledge, the P–H insertion reactions of α -imino metal carbene for the construction of C–P bond has never been reported.

Received: December 16, 2016

Published: February 1, 2017

We began to explore the P–H insertion of 3-diazoindolin-2-imine (**1a**) and diphenylphosphine oxide (**2a**) with the common catalysts used for the decomposition of triazoles and diazo compounds, including $\text{Rh}_2(\text{oct})_4$, $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$, $(\eta^3\text{-C}_3\text{H}_5)_2\text{Pd}_2\text{Cl}_2$, and $\text{Ni}(\text{cod})_2$ (Table 1, entries 1–4).

Table 1. Optimization of the Reaction Conditions^a




entry	catalyst	solvent	temp (°C)	time (h)	yield ^b (%)
1	$\text{Rh}_2(\text{oct})_4$	DCE	35	24	0
2	$[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$	DCE	35	24	0
3	$(\eta^3\text{-C}_3\text{H}_5)_2\text{Pd}_2\text{Cl}_2$	DCE	35	24	0
4	$\text{Ni}(\text{cod})_2$	DCE	35	24	0
5	$\text{Cu}(\text{OTf})_2$	DCE	35	24	20
6	$\text{Cu}(\text{OAc})_2$	DCE	35	24	15
7	$\text{Cu}(\text{acac})_2$	DCE	35	24	15
8	Cu_2O	DCE	35	24	23
9	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	DCE	35	24	42
10	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	DCE	35	1	42
11	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	DCE	50	1	56
12	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	DCE	15	1	trace
13		DCE	50	1	0
14	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	DCM	50	1	26
15	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	CHCl_3	50	1	82
16	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	1, 4-dioxane	50	1	64
17	$\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$	toluene	50	1	52

^aReaction conditions: **1a** (0.2 mmol), **2a** (0.2 mmol), catalyst (0.01 mmol), solvent (0.8 mL). ^bIsolated yield.

Unfortunately, no desired product but a significant amount of starting material was observed. To our delight, the reaction gave **3a** in 20% yield in the presence of 5 mol % of $\text{Cu}(\text{OTf})_2$ in DCE at 35 °C (Table 1, entry 5). Encouraged by this initial result, we then screened other copper catalysts, of which $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ proved superior (Table 1, entries 5–9). By screening the reaction temperature and reaction time, the optimal reaction temperature and reaction time were found to be 50 °C and 1 h, respectively. A trace amount of the desired **3a** was exactly detected when the reaction was carried out at 15 °C (entry 12). A control reaction in the absence of metal catalyst afforded no desired product, indicating the critical role of the copper catalyst in generating the α -imino metal carbene intermediate (entry 11 vs 13). The reaction was subjected to notable solvent effects. Chloroform (CHCl_3) was a better solvent over $\text{ClCH}_2\text{CH}_2\text{Cl}$ (DCE), CH_2Cl_2 (DCM), 1, 4-dioxane, and toluene, providing **3a** in 82% yield (entries 15–17).

With the optimal reaction conditions, we then explored the substrate scope with respect to the 3-diazoindolin-2-imines. As illustrated in Scheme 2, the alkyl group on the 1-position of indole could be varied from methyl, ethyl, isopropyl, allyl, and benzyl to produce **3a**, **3b**, **3c**, **3d**, and **3e** in 82%, 76%, 71%, 79%, and 58% yields, respectively. The structure of **3e** was unambiguously determined by single-crystal X-ray analysis (as shown in the Supporting Information). No insertion product was observed by using *N*-unprotected 3-diazoindolin-2-imines. Various 3-diazoindolin-2-imines with substituents on the 5-

Scheme 2. Scope of 3-Diazoindolin-2-imines^{a,b}




^aReaction conditions: $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.01 mmol), **1** (0.20 mmol) and $\text{HP}(\text{O})\text{Ph}_2$ (0.20 mmol) in CHCl_3 (0.8 mL) at 50 °C for 1 h. ^bIsolated yield.


position of indole were feasible substrates in this reaction and delivered **3f–i** in 53–74% yields. The process was tolerant to 3-diazoindolin-2-imines bearing different sulfonyl groups ($\text{R}^3 = 4\text{-BrC}_6\text{H}_4$, $4\text{-CF}_3\text{C}_6\text{H}_4$, $2,4,6\text{-i-Pr}_3\text{C}_6\text{H}_2$), thus giving the corresponding products **3j–l** with high chemoselectivities (65–88% yield).

We next investigated the scope with respect to the H-phosphine oxides (Scheme 3). H-Phosphine oxides **2** under the optimal reaction conditions and reacted with **1** generally proceeded in good yields. Both electron-poor and electron-rich substituted groups in H-phosphine oxides (**2d,e**) could be tolerated and afforded **3o** and **3p** in 86% and 88% yield, respectively. Notably, heterocyclic H-phosphine oxides such as 9,10-dihydro-9-oxa-10-phosphaphenanthrene 10-oxide (DOPO, **2f**) could be introduced to the insertion product. Moreover, ethyl phenylphosphinate (**2g**) was also viable for this transformation, giving the expected product **3s** in 42% yield. In the case where butyl(phenyl)phosphine oxide (**2h**) was used as a substrate, the desired product was isolated in 57% yield. The tosyl group can be removed with concentrated sulfuric acid to afford stable 2-free amino indole **6** in 91% yield (Supporting Information).

Inspired by the success in the P–H insertions of α -imino copper carbene, we then turned our attention to the asymmetric synthesis in an attempt to obtain a chiral 3-phosphoindoles. (1*R*,2*S*,5*R*)-(–)-Menthoxypheylphosphinate (**4**) was employed as the chiral reagent for the chirality induction (Scheme 4). The P–H insertion of α -imino copper carbene proceeded smoothly to give the chiral 3-phosphoindoles (**5a–f**) in moderate yield with very high diastereoselectivity. The absolute configuration of **5f** was established by X-ray single-crystal analysis (Figure 1).

Scheme 3. Scope of H-Phosphine Oxides^{a,b}

^aReaction conditions: $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.01 mmol), **1a** (0.20 mmol) and **2** (0.20 mmol) in CHCl_3 (0.8 mL) at 50 °C for 1 h.
^bIsolated yield.

Scheme 4. Stereoselective Synthesis of Chiral 3-Methoxyarylphosphindoles^{a,b}

^aReaction conditions: $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.01 mmol), **1a** (0.20 mmol), and $\text{HP}(\text{O})(\text{OMe})\text{Ph}$ (0.20 mmol) in CHCl_3 (0.8 mL) at 50 °C for 8 h. ^bIsolated yield and the diastereomeric ratio were detected by ^1H NMR (600 MHz) spectroscopy.


Figure 1. X-ray structure of **5f**.

P-Stereogenic phosphorus compounds have attracted considerable attention due to their wide applications as chiral ligands for asymmetric synthesis.¹² The stereoselective synthesis of *P*-stereogenic compounds is one of the most

important and challenging subjects in synthetic organic chemistry.^{13,14} In this asymmetric P–H insertion, the H–P bond in (1*R*,2*S*,5*R*)-(–)-menthoxyphenylphosphinate (**4**) was converted into a $\text{C}(\text{sp}^2)\text{--P}$ bond to give chiral 3-phosphindoles with high setereoselectivities. Although there are several examples of P–H insertions reported in the formation of the $\text{C}(\text{sp}^3)\text{--P}$ bond,¹⁵ no asymmetric P–H insertion reaction has been achieved to date. The stereoselective P–H insertion of α -imino copper carbene represents a unique example of asymmetric P–H insertion, affording chiral 3-phosphindoles with a *P*-stereogenic center.

On the basis of the preliminary results of control reactions (as shown in Table 1, entry 11 vs 13) and previous work,^{3a,16} we envisioned that the insertion of α -imino copper carbene into P–H bonds may proceed in two plausible pathways (Scheme 5). The α -imino copper carbene intermediate derived from 3-

Scheme 5. Proposed Mechanism



diazindolin-2-imines may undergo direct 1,1-insertion into the P–H bond, followed by tautomerization leading to product of 3-phosphindoles (path a). Alternatively, the insertion may proceed by the formation of zwitterionic P-ylide intermediate. Subsequent dissociation of copper results in a proton transfer from the phosphorus to imine moiety, finally furnishing the 1,3-insertion product of 3-phosphindoles (path b).

In summary, we have developed P–H insertions of α -imino copper carbenes for the construction of $\text{C}(\text{sp}^2)\text{--P}$ bonds. The protocol provides an efficient approach to 3-phosphindoles from 3-diazindolin-2-imines and H-phosphine oxides with good chemoselectivity. In addition, the stereoselective P–H insertion was also successfully developed with high diastereoselectivity. The investigations on the biological activities of the new 3-phosphindoles are in progress.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b03752.

Experimental procedures, characterization data, and NMR spectra of all new compounds (PDF)

X-ray crystallography of **3e** (CIF)

X-ray crystallography of **5f** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: jiangjun@gxu.edu.cn.

ORCID

Jun Jiang: 0000-0002-5542-7592

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful for financial support from Natural Science Foundation of China (21402032), Guangxi Natural Science Foundation (2014GXNSFBA118031, 2015GXNSFDA139008), and the Scientific Research Foundation of Guangxi University (XQZ130869)

■ REFERENCES

- (1) (a) Doyle, M. P.; McKervey, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*; Wiley, 1998. (b) Liu, L.; Zhang, J. L. *Chem. Soc. Rev.* **2016**, 45, 506. (c) Zhu, S. F.; Zhou, Q. L. *Acc. Chem. Res.* **2012**, 45, 1365. (d) Zhang, Z. H.; Wang, J. B. *Tetrahedron* **2008**, 64, 6577. (e) Ye, T.; McKervey, M. A. *Chem. Rev.* **1994**, 94, 1091.
- (2) For recent reviews, see: (a) Davies, H. M.; Alford, J. S. *Chem. Soc. Rev.* **2014**, 43, 5151. (b) Gulevich, A. V.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2013**, 52, 1371. (c) Chattopadhyay, B.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2012**, 51, 862.
- (3) (a) Chuprakov, S.; Worrell, B. T.; Selander, N.; Sit, R. K.; Fokin, V. V. *J. Am. Chem. Soc.* **2014**, 136, 195. (b) Zibinsky, M.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2013**, 52, 1507. (c) Chuprakov, S.; Kwok, S. W.; Fokin, V. V. *J. Am. Chem. Soc.* **2013**, 135, 4652. (d) Selander, N.; Worrell, B. T.; Chuprakov, S.; Velaparthi, S.; Fokin, V. V. *J. Am. Chem. Soc.* **2012**, 134, 14670. (e) Selander, N.; Fokin, V. V. *J. Am. Chem. Soc.* **2012**, 134, 2477. (f) Chuprakov, S.; Malik, J. A.; Zibinsky, M.; Fokin, V. V. *J. Am. Chem. Soc.* **2011**, 133, 10352. (g) Grimster, N.; Zhang, L.; Fokin, V. V. *J. Am. Chem. Soc.* **2010**, 132, 2510. (h) Horneff, T.; Chuprakov, S.; Chernyak, N.; Gevorgyan, V.; Fokin, V. V. *J. Am. Chem. Soc.* **2008**, 130, 14972.
- (4) (a) Gulevich, A. V.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2013**, 52, 1371. (b) Shi, Y.; Gevorgyan, V. *Org. Lett.* **2013**, 15, 5394. (c) Chattopadhyay, B.; Gevorgyan, V. *Org. Lett.* **2011**, 13, 3746.
- (5) (a) Parr, B. T.; Davies, H. M. L. *Angew. Chem., Int. Ed.* **2013**, 52, 10044. (b) Spangler, J. E.; Davies, H. M. L. *J. Am. Chem. Soc.* **2013**, 135, 6802. (c) Parr, B. T.; Green, S. A.; Davies, H. M. L. *J. Am. Chem. Soc.* **2013**, 135, 4716. (d) Alford, J. S.; Davies, H. M. L. *Org. Lett.* **2012**, 14, 6020.
- (6) (a) Lindsay, V. N.; Viart, H. M.; Sarpong, R. *J. Am. Chem. Soc.* **2015**, 137, 8368. (b) Schultz, E. E.; Lindsay, V. N. G.; Sarpong, R. *Angew. Chem., Int. Ed.* **2014**, 53, 9904. (c) Schultz, E. E.; Sarpong, R. *J. Am. Chem. Soc.* **2013**, 135, 4696.
- (7) (a) Jiang, Y.; Wang, Q.; Sun, R.; Tang, X. Y.; Shi, M. *Org. Chem. Front.* **2016**, 3, 744. (b) Jiang, Y.; Tang, X. Y.; Shi, M. *Chem. Commun.* **2015**, 51, 2122. (c) Tang, X. Y.; Zhang, Y. S.; He, L.; Wei, Y.; Shi, M. *Chem. Commun.* **2015**, 51, 133. (d) Tang, X. Y.; Zhang, Y. S.; He, L.; Wei, Y.; Shi, M. *Chem. Commun.* **2015**, 51, 133. (e) Jiang, Y.; Sun, R.; Wang, Q.; Tang, X. Y.; Shi, M. *Chem. Commun.* **2015**, 51, 16968. (f) Chen, K.; Zhu, Z. Z.; Zhang, Y. S.; Tang, X. Y.; Shi, M. *Angew. Chem., Int. Ed.* **2014**, 53, 6645. (g) Yang, J. M.; Zhu, C. Z.; Tang, X. Y.; Shi, M. *Angew. Chem., Int. Ed.* **2014**, 53, 5142.
- (8) (a) Ishikura, M.; Abe, T.; Choshi, T.; Hibino, S. *Nat. Prod. Rep.* **2010**, 27, 1630. (b) Gribble, G. W., Ed. In *Heterocyclic Scaffolds II: Reactions and Applications of Indoles*; Topics in Heterocyclic Chemistry; Springer: Heidelberg, 2010; Vol. 26. (c) Kochanowska-Karamyan, A. J.; Hamann, M. T. *Chem. Rev.* **2010**, 110, 4489. (d) Burgett, A. W. G.; Li, Q. Y.; Wei, W.; Harran, P. G. *Angew. Chem., Int. Ed.* **2003**, 42, 4961. (e) Crich, D.; Banerjee, A. *Acc. Chem. Res.* **2007**, 40, 151.
- (9) (a) Qian, J.; Sheng, G. R.; Huang, K.; Liu, S. J.; Lu, P.; Wang, Y. G. *Org. Lett.* **2016**, 18, 3682. (b) Wang, C.; Zhang, H.; Lang, B.; Ren, A.; Lu, P.; Wang, Y. *Org. Lett.* **2015**, 17, 4412. (c) Sheng, G. R.; Huang, K.; Ma, S. C.; Qian, J.; Lu, P.; Wang, Y. G. *Chem. Commun.* **2015**, 51, 11056. (d) Xing, Y. P.; Sheng, G. R.; Wang, J.; Lu, P.; Wang, Y. G. *Org. Lett.* **2014**, 16, 1244. (e) Sheng, G. R.; Huang, K.; Chi, Z. H.; Ding, H. L.; Xing, Y. P.; Lu, P.; Wang, Y. G. *Org. Lett.* **2014**, 16, 5096.
- (10) (a) Monge, S.; David, G. *Phosphorus-Based Polymers from Synthesis to Applications*; The Royal Society of Chemistry: Cambridge, 2014. (b) Quin, L. D. *A Guide to Organophosphorus Chemistry*; Wiley Interscience: New York, 2000. (c) Montchamp, J. L. *Acc. Chem. Res.* **2014**, 47, 77. (d) Helmchen, G.; Pfaltz, A. *Acc. Chem. Res.* **2000**, 33, 336. (e) Corbridge, D. E. C. *Phosphorus: Chemistry, Biochemistry and Technology*, 6th ed.; CRC Press: New York, 2013. (f) Pereira, M. M.; Calvete, M. J. F.; Carrilho, R. M. B.; Abreu, A. R. *Chem. Soc. Rev.* **2013**, 42, 6990. (g) Lim, C. W.; Tissot, O.; Mattison, A.; Hooper, M. W.; Brown, J. M.; Cowley, A. R.; Hulmes, D. I.; Blacker, A. J. *Org. Process Res. Dev.* **2003**, 7, 379.
- (11) (a) Storer, R.; Dousson, C.; Alexandre, F. R.; Roland, A. WOPCT Int Appl. 054182, 2006. (b) Zhou, X. J.; Garner, R. C.; Nicholson, S.; Kissling, C. J.; Mayers, D. J. *Clin. Pharmacol.* **2009**, 49, 1408. (c) La Regina, G.; Coluccia, A.; Silvestri, R. *Antiviral Chem. Chemother.* **2010**, 20, 213. (d) Zhou, X. J.; Pietropaolo, K.; Dampousse, D.; Belanger, B.; Chen, J.; Sullivan-Bolyai, J.; Mayers, D. *Antimicrob. Agents Chemother.* **2009**, 53, 1739.
- (12) (a) Xie, J. H.; Zhou, Q. L. *Acc. Chem. Res.* **2008**, 41, 581. (b) Tang, W.; Zhang, X. *Chem. Rev.* **2003**, 103, 3029.
- (13) For reviews on catalytic stereoselective synthesis of P-stereogenic compounds: (a) Harvey, J. S.; Gouverneur, V. *Chem. Commun.* **2010**, 46, 7477. (b) Glueck, D. S. *Chem. - Eur. J.* **2008**, 14, 7108. (c) Glueck, D. S. *Synlett* **2007**, 2007, 2627. (d) Grabulosa, A.; Granell, J.; Muller, G. *Coord. Chem. Rev.* **2007**, 251, 25–90. (e) Wozniak, L. A.; Okruszek, A. *Chem. Soc. Rev.* **2003**, 32, 158.
- (14) Recent reports: (a) Sun, Y.; Cramer, N. *Angew. Chem., Int. Ed.* **2017**, 56, 364. (b) Lin, Z. Q.; Wang, W. Z.; Yan, S. B.; Duan, W. L. *Angew. Chem., Int. Ed.* **2015**, 54, 6265. (c) Du, Z. J.; Guan, J.; Wu, G. J.; Xu, P.; Gao, L. X.; Han, F. S. *J. Am. Chem. Soc.* **2015**, 137, 632. (d) Nikitin, K.; Rajendran, K. V.; Müller-Bunz, H.; Gilheany, D. G. *Angew. Chem., Int. Ed.* **2014**, 53, 1906. (e) Huang, Y. H.; Li, Y. X.; Leung, P. H.; Hayashi, T. *J. Am. Chem. Soc.* **2014**, 136, 4865. (f) Toda, Y.; Pink, M.; Johnston, J. N. *J. Am. Chem. Soc.* **2014**, 136, 14734. (g) Han, Z. S.; Goyal, N.; Herbage, M. A.; Sieber, J. D.; Qu, B.; Xu, Y.; Li, Z.; Reeves, J. T.; Desrosiers, J. N.; Ma, S.; Grinberg, N.; Lee, H.; Mangunuru, H. P. R.; Zhang, Y.; Krishnamurthy, D.; Lu, B. Z.; Song, J. J.; Wang, G.; Senanayake, C. H. *J. Am. Chem. Soc.* **2013**, 135, 2474. (h) Ding, B.; Zhang, Z.; Xu, Y.; Liu, Y.; Sugiyama, M.; Imamoto, T.; Zhang, W. *Org. Lett.* **2013**, 15, 5476.
- (15) (a) Chen, Z. S.; Zhou, Z. Z.; Hua, H. L.; Duan, X. H.; Luo, J. Y.; Wang, J.; Zhou, P. X.; Liang, Y. M. *Tetrahedron* **2013**, 69, 1065. (b) Jiang, H. L.; Jin, H. M.; Abdulkader, A.; Lin, A. J.; Cheng, Y. X.; Zhu, C. J. *Org. Biomol. Chem.* **2013**, 11, 3612. (c) Bartrum, H. E.; Blakemore, D. C.; Moody, C. J. *Tetrahedron* **2013**, 69, 2276. (d) Miao, W. J.; Gao, Y. Z.; Li, X. Q.; Gao, Y. X.; Tang, G.; Zhao, Y. F. *Adv. Synth. Catal.* **2012**, 354, 2659. (e) Wu, L.; Zhang, X.; Chen, Q. Q.; Zhou, A. K. *Org. Biomol. Chem.* **2012**, 10, 7859.
- (16) (a) Miura, T.; Biyajima, T.; Fujii, T.; Murakami, M. *J. Am. Chem. Soc.* **2012**, 134, 194. (b) Miura, T.; Tanaka, T.; Biyajima, T.; Yada, A.; Murakami, M. *Angew. Chem., Int. Ed.* **2013**, 52, 3883.