


## A Stereoselective Arylative-Cyclopropanation Process

Siomenan Coulibali, Elsa Deruer, Elizabeth Godin, and Sylvain Canesi\*<sup>ID</sup>

Laboratoire de Méthodologie et Synthèse de Produits Naturels, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville, Montréal, H3C 3P8, Québec, Canada


## S Supporting Information

**ABSTRACT:** A new stereoselective arylative cyclopropanation process involving treatment of halogenated dienone systems in the presence of a Michael donor containing a nitro-aryl-sulfone has been developed. This transformation enables production of an arylated cyclopropane under mild conditions and occurs via a Michael–Smiles ring closure cascade process, reflecting the concepts of green chemistry and atom economy.



Cyclopropanation is an important process in organic synthesis, yielding three-membered ring structures that are useful as precursors for the elaboration of advanced intermediates. Cyclopropanes are formed in several processes such as the Simmons–Smith<sup>1</sup> method or its variants mediated by diazo compounds or iodonium ylides,<sup>2</sup> Corey–Chaykovsky<sup>3</sup> or Kulinkovich transformations,<sup>4</sup> or intramolecular alkylation.<sup>5</sup> There is increasing sentiment that transformations should be atom economical<sup>6</sup> and should respect the concept of “green chemistry.” To this end, we developed a methodology enabling the formation of functionalized scaffolds **3** mediated by a Michael–Smiles tandem process.<sup>7</sup> In this paper we describe a new application of this approach on a halogenated enone to develop an arylative cyclopropanation strategy, [Scheme 1](#).


Scheme 1. Michael–Smiles Tandem Process Avenues



One advantage of the Smiles rearrangement<sup>8</sup> is its capacity for rapid reconfiguration of a structure into a more elaborated one under slightly basic and mild thermal conditions. A judicious combination of this process with a tandem Michael addition leads to  $\alpha$ – $\beta$ -disubstituted cyclohexanones **3**.<sup>7</sup> Indeed, a malonate derivative containing a nitro-aryl sulfonyl moiety<sup>9</sup> **2** may be used as an activating group, enabling an initial Michael addition on enone moiety **1** to produce an enolate and triggering a Smiles rearrangement yielding **3** ([Scheme 1](#)) in which SO<sub>2</sub> serves as a leaving group and is released as the sole byproduct. During our efforts to extend the scope of this

process, we observed that the presence of enone **6** resulted in formation of an interesting cyclopropanated byproduct **11** with very good diastereoselectivity instead of **3**. The yield of this transformation product was increased in the presence of oxygen and decreased in the presence of TEMPO, suggesting a free radical process<sup>10</sup> via oxygen-mediated electron transfer was responsible. A potential mechanism is described in [Scheme 2](#).

Scheme 2. Radical-Based-Cyclopropanation Mechanism




It should be noted that compound **11** represents a polyfunctionalized core obtained in a diastereoselective and environmentally benign manner in one step from simple starting materials. However, we were unable to extend this method to other substrates and only one other cyclopropanated example **13** was obtained from **12** under similar conditions, [Scheme 3](#).

We therefore investigated alternative methods of generating this cyclopropane scaffold, beginning with introduction of a bromine on the enone moiety **14** to replace the uncertain oxidative cyclopropanation step with an intramolecular alkylation. We were pleased to observe that cyclopropane **11** was produced in 80% yield by this strategy, [Scheme 4](#).


Received: January 23, 2017

Published: February 16, 2017

Scheme 3. Intramolecular-Radical-Based Cyclopropanation




Scheme 4. Halide-Mediated Cyclopropanation Alternative



We assume that the mechanism is similar to the mechanism described in Scheme 2 and the prototypical transformation involves a Michael–Smiles cyclopropanation cascade process. We suppose that the aryl migration occurs with retention of configuration leading to 18 from 17 and enabling an antiperiplanar attack to produce cyclopropane 19. One interesting aspect is the potential intramolecular nucleophilic substitution that would occur on a tertiary alkyl bromide with inversion of configuration. We suspect that a partial carbocation leading to a potential enolonium ion<sup>11</sup> could be involved due to the presence of a conjugated system; this species could be generated in the presence of a cesium ion acting as a Lewis acid. When radical processes were excluded by performing the reaction in the absence of oxygen or in the presence of TEMPO, the cyclopropane 19 was still obtained in comparable yields, Scheme 5.

Scheme 5. Michael–Smiles Cyclopropanation Mechanism



Encouraged by this result, we investigated different halides as nucleofuges. These attempts have been performed on dienone 16, which represents a good Michael acceptor obtained by direct oxidation of halogenated phenols 15 in the presence of methanol and a hypervalent iodine reagent.<sup>12</sup> Iodine and bromine provided slightly improved results, Scheme 6. Unfortunately, this process was not efficient with a simple halogenated enone.

The transformation was subsequently performed on several bromo-dienones 21, derivatives of 16. Bromine was selected as the halogen atom because it was easily introduced on the phenol precursor. The *ortho*- and *para*-nosyl derivatives 20 have both been tested. An interesting aspect of this process is the good stereoselectivity observed, with only one diastereomer observed in NMR spectra.<sup>13</sup> The corresponding systems 22 that were obtained are described in Table 1.

Scheme 6. Rapid Elaboration of Functionalized Tricycles




Table 1. Michael–Smiles Cyclopropanation Process

entry	NO <sub>2</sub>	R <sub>1</sub>	R <sub>2</sub>	yield (%)
a	ortho	Me	Me	64
b	para	Me	Me	68
c	ortho	Et	H	61
d	para	Me	H	70
e	ortho	<i>t</i> -Bu	H	74
f	para	<i>t</i> -Bu	H	73
g	ortho	Me	H	63
h	para	allyl	H	70
i	para	Br	H	50
j	para	Et	H	71


In order to produce a highly substituted cyclopropane subunit, the process was extended to compound 23, a methyl derivative of 20. We were pleased to observe that, once again, only one diastereomer was observed by NMR<sup>13</sup> in the substituted cyclopropane 25.<sup>14</sup> The diastereoselectivity was further verified using NMR NOE. Some examples are described in Table 2.

Table 2. Formation of Highly Substituted Cyclopropanes

entry	R <sub>1</sub>	yield (%)
a	<i>t</i> -Bu	67
b	allyl	61
c	Et	60
d	Me	49


Interestingly, in the presence of a methyl-substituted malonate derivative 23, the ester functionality was *anti* to the nitro-aryl segment. The difference of diastereoselectivity observed between the formation of cyclopropane 22 and its more substituted analogous 25 could be explained by their enolate precursors 26. Indeed, if R<sub>1</sub> is a hydrogen, we supposed that conformer 26a was favored and led to compound 22. However, in the presence of a methyl group (R<sub>1</sub> = Me) a potential steric interaction between the aryl and methyl groups would favor conformer 26b yielding cyclopropane 25, Scheme 7.

Scheme 7. Diastereoselectivity Issues



An asymmetric variant containing an Evans auxiliary<sup>15</sup> was developed, and the corresponding cyclopropanes **29** were obtained in 41% and 44% yields with an excellent stereoselectivity of 95% based on NMR.<sup>13</sup> A potential approach suggesting an opened transition state to explain the observed stereoselectivity is depicted in Scheme 8.

Scheme 8. Asymmetric Avenue



In summary, we have developed a novel stereoselective arylative cyclopropanation transformation mediated by a Michael–Smiles ring closure cascade process. These methods occur under slightly basic conditions at moderate temperatures with sulfur dioxide as the only byproduct. Very good diastereoselectivity is observed during this multistep transformation even during the formation of highly substituted cyclopropane moieties. We are currently looking for novel applications for these transformations.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.7b00248.

Experimental procedures and spectral data for key compounds (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: canesi.sylvain@uqam.ca.

### ORCID

Sylvain Canesi: 0000-0002-0639-7796

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We are very grateful to the Natural Sciences and Engineering Research Council of Canada (NSERC), the Canada Foundation for Innovation (CFI), and the provincial government of

Quebec (FQRNT and CCVC) for their precious financial support in this research. S.C. thanks the “Programme Canadien de Bourses de la Francophonie” for a scholarship.

## ■ REFERENCES

- (a) Simmons, H. E.; Smith, R. D. *J. Am. Chem. Soc.* **1958**, *80*, 5323. (b) Simmons, H. E.; Smith, R. D. *J. Am. Chem. Soc.* **1959**, *81*, 4256. (c) Charette, A. B.; Beauchemin, A. *Org. React.* **2001**, *58*, 1. (d) Hoveyda, A. H.; Evans, D. A.; Fu, G. C. *Chem. Rev.* **1993**, *93*, 1307. (e) Pellissier, H. *Tetrahedron* **2008**, *64*, 7041. (f) Lebel, H.; Marcoux, J. F.; Molinaro, C.; Charette, A. B. *Chem. Rev.* **2003**, *103*, 977. (g) Furukawa, J.; Kawabata, N. *Adv. Organomet. Chem.* **1974**, *12*, 83.
- (a) Doyle, M. P.; McKerver, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides*; Wiley: New York, 1998. (b) Moreau, B.; Charette, A. B. *J. Am. Chem. Soc.* **2005**, *127*, 18014. (c) Marcoux, D.; Goudreau, S. R.; Charette, A. B. *J. Org. Chem.* **2009**, *74*, 8939. (d) Nani, R. R.; Reisman, S. E. *J. Am. Chem. Soc.* **2013**, *135*, 7304. (e) Deng, C.; Wang, L.; Zhu, J.; Tang, Y. *Angew. Chem., Int. Ed.* **2012**, *51*, 11620. (f) Lindsay, V. N. G.; Fiset, D.; Gritsch, P. J.; Azzi, S.; Charette, A. B. *J. Am. Chem. Soc.* **2013**, *135*, 1463. (g) Lindsay, V. N. G.; Fiset, D.; Gritsch, P. J.; Azzi, S.; Charette, A. B. *J. Am. Chem. Soc.* **2013**, *135*, 1463. (h) Duan, Y. N.; Zhang, Z.; Zhang, C. *Org. Lett.* **2016**, *18*, 6176.
- (a) Johnson, A. W.; LaCount, R. B. *J. Am. Chem. Soc.* **1961**, *83*, 417. (b) Corey, E. J.; Chaykovsky, M. *J. Am. Chem. Soc.* **1965**, *87*, 1353. (c) Li, A. H.; Dai, L. X.; Aggarwal, V. K. *Chem. Rev.* **1997**, *97*, 2341.
- (a) Kulinkovich, O. G.; Sviridov, S. V.; Vasilevskii, D. A.; Pritytskaya, T. S. *Zh. Org. Khim.* **1989**, *25*, 2244. (b) Kulinkovich, O. G.; de Meijere, A. *Chem. Rev.* **2000**, *100*, 2789.
- (a) Papageorgiou, C. D.; Ley, S. V.; Gaunt, M. J. *Angew. Chem., Int. Ed.* **2003**, *42*, 828. (b) Liao, W. W.; Li, K.; Tang, Y. *J. Am. Chem. Soc.* **2003**, *125*, 13030. (c) Bremeyer, N. S.; Smith, S. C.; Ley, S. V.; Gaunt, M. J. *Angew. Chem., Int. Ed.* **2004**, *43*, 2681. (d) Kunz, R. K.; MacMillan, D. W. *J. Am. Chem. Soc.* **2005**, *127*, 3240. (e) Ibrahim, I.; Zhao, G. L.; Rios, R.; Vesely, J.; Sundén, H.; Dziedzic, P.; Córdova, A. *Chem. - Eur. J.* **2008**, *14*, 7867. (f) Gao, L.; Hwang, G. S.; Ryu, D. H. *J. Am. Chem. Soc.* **2011**, *133*, 20708. (g) Ferrary, T.; David, E.; Milanole, G.; Besset, T.; Jubault, P.; Pannecoucke, X. *Org. Lett.* **2013**, *15*, 5598. (h) Aitken, L. S.; Hammond, L. E.; Sundaram, R.; Shankland, K.; Brown, G. D.; Cobb, A. *Chem. Commun.* **2015**, *51*, 13558. (i) Shim, S. Y.; Kim, J. Y.; Nam, M.; Hwang, G. S.; Ryu, D. H. *Org. Lett.* **2016**, *18*, 160.
- (a) Trost, B. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 259. (b) Baran, P. S.; Maimone, T. J.; Richter, J. M. *Nature* **2007**, *446*, 404.
- Coulbali, S.; Godou, T.; Canesi, S. *Org. Lett.* **2016**, *18*, 4348.
- (a) Levy, A. A.; Rains, H. C.; Smiles, S. *J. Chem. Soc.* **1931**, *0*, 3264. (b) Truce, W. E.; Kreider, E. M.; Brand, W. W. *Organic Reactions* **1970**, *18*, 99.
- Kan, T.; Fukuyama, T. *Chem. Commun.* **2004**, 353.
- Boyd, R. J.; Gupta, A.; Langer, R. F.; Lownie, S. P.; Pincock, J. A. *Can. J. Chem.* **1980**, *58*, 331.
- Arava, S.; Kumar, J. N.; Maksymenko, S.; Iron, M. A.; Parida, K. N.; Frisrup, P.; Szpilman, A. M. *Angew. Chem., Int. Ed.* **2017**, *129*, doi: 10.1002/anie.201610274.
- (a) Tamura, Y.; Yakura, T.; Haruta, J.; Kita, Y. *J. Org. Chem.* **1987**, *52*, 3927. (b) Kita, Y.; Tohma, H.; Hatanaka, K.; Takada, T.; Fujita, S.; Mitoh, S.; Sakurai, H.; Oka, S. *J. Am. Chem. Soc.* **1994**, *116*, 3684. (c) Dohi, T.; Yamaoka, N.; Kita, Y. *Tetrahedron* **2010**, *66*, 5775. (d) Singh, J.; Rehbein, F. V.; Wirth, T. *ChemistryOpen* **2012**, *1*, 245. (e) Farid, U.; Malmedy, F.; Claveau, R.; Albers, L.; Wirth, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 7018. (f) Zhdankin, V. V. *Hypervalent Iodine Chemistry: Preparation, Structure, and Synthetic Applications of Polyvalent Iodine Compounds*; John Wiley & Sons: Chichester, U.K., 2013. (g) Pouységu, L.; Deffieux, D.; Quideau, S. *Tetrahedron* **2010**, *66*, 2235. (h) Jacquemot, G.; Canesi, S. *J. Org. Chem.* **2012**, *77*, 7588. (i) Jacquemot, G.; Menard, M. A.; L'Homme, C.; Canesi, S. *Chem. Sci.* **2013**, *4*, 1287. (j) Beaulieu, M. A.; Ottenwaelde, X.; Canesi, S. *Chem.*

*Eur. J.* **2014**, *20*, 7581. (k) L'Homme, C.; Menard, M. A.; Canesi, S. *J. Org. Chem.* **2014**, *79*, 8481. (l) Desjardins, S.; Maertens, G.; Canesi, S. *Org. Lett.* **2014**, *16*, 4928. (m) Jacquemot, G.; Maertens, G.; Canesi, S. *Chem. - Eur. J.* **2015**, *21*, 7713. (n) Guérard, K. C.; Hamel, V.; Guérinot, A.; Bouchard-Aubin, C.; Canesi, S. *Chem. - Eur. J.* **2015**, *21*, 18068.

(13) Only one diastereomer was observed by NMR, and the stereoselectivity was verified by nuclear Overhauser effect (NOE). The coupling constants for the cyclopropane hydrogens **22** were close to 6 Hz and in agreement with a *trans* isomer.

(14) Unfortunately, this process was not efficient in the presence of an alkyl-substituted malonate derivative more sterically hindered.

(15) Evans, D. A.; Bartroli, J.; Shih, T. L. *J. Am. Chem. Soc.* **1981**, *103*, 2127.