

This article was downloaded by:

On: 29 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

PHENYLATION OF PLATINUM(II) THIOETHER COMPLEXES BY TETRAPHENYLBORATE(III) IN SOLID STATE AND NITROMETHANE SOLUTION

Vadim Yu^a; Kukushkin^{ab}; Karin Löqvist^a; Bertil Norén^a; Åke Oskarsson^a; Lars Ivar Elding^a

^a Inorganic Chemistry 1, Chemical Center, University of Lund, Lund, Sweden ^b Department of Chemistry, St. Petersburg State University, Stary Petergof, Russia

To cite this Article Yu, Vadim , Kukushkin, Löqvist, Karin , Norén, Bertil , Oskarsson, Åke and Elding, Lars Ivar(1992) 'PHENYLATION OF PLATINUM(II) THIOETHER COMPLEXES BY TETRAPHENYLBORATE(III) IN SOLID STATE AND NITROMETHANE SOLUTION', Phosphorus, Sulfur, and Silicon and the Related Elements, 73: 1, 253 — 255

To link to this Article: DOI: 10.1080/10426509208034450

URL: <http://dx.doi.org/10.1080/10426509208034450>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Communication

PHENYLATION OF PLATINUM(II) THIOETHER COMPLEXES BY TETRAPHENYLBORATE(III) IN SOLID STATE AND NITROMETHANE SOLUTION

VADIM YU. KUKUSHKIN,[‡] KARIN LÖVQVIST, BERTIL NORÉN,
ÅKE OSKARSSON and LARS IVAR ELDING[†]


*Inorganic Chemistry 1, Chemical Center, University of Lund, P.O. Box 124,
S-221 00 Lund, Sweden*

(Received August 21, 1992; in final form September 26, 1992)

Platinum(II) complexes of the type $[\text{Pt}(\text{thioether})_3\text{Cl}]^+$ (thioether = dimethyl sulfide, thioxane) are capable of abstracting a phenyl group from the BPh_4^- counterion with formation of *trans*- $[\text{Pt}(\text{thioether})_2\text{ClPh}]$ compounds. Thermal reactions proceed both in the solid phase and in nitromethane solution at elevated temperature and have preparative importance. Phenylation of the Pt(II) centre also occurs in reaction between $[\text{Pt}(\text{Me}_2\text{S})_2\text{Cl}_2]$ and AgBPh_4 in CH_2Cl_2 suspension. Brief X-ray crystallographic structural data for *trans*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{ClPh}]$ and $[\text{Pt}(\text{thioether})_3\text{Cl}]\text{X}$ (thioether = Me_2S , tx, X = SO_3CF_3 ; thioether = Me_2S , X = PF_6) are reported.

Key words: Platinum(II) thioether complexes; phenyl abstraction; tetraphenylborate ion; synthesis of thioether complexes; silver tetraphenylborate; X-ray structures.

The ability of platinum(II) complexes to abstract a phenyl group from the tetraphenylborate(III) anion, BPh_4^- , is well documented.¹ Phenyl migration reactions, however, are not common. In all cases so far studied phenyl abstraction proceeded with or via formation of electrophilic complexes containing weakly coordinated solvento ligands or anions of strong acids. We have found that complexes of the type $[\text{Pt}(\text{thioether})_3\text{Cl}]^+$, where thioether = dimethyl sulfide (Me_2S) or 1,4-thioxane (tx), which do not contain weakly coordinated ligands² are nevertheless sufficiently electrophilic to abstract a phenyl group from the tetraphenylborate counterion with displacement of one of the thioether ligands, probably in *trans*-position:




Studies of the kinetics and mechanism of this reaction in nitromethane solution are in progress in our laboratory. The thermal phenyl migration reaction is of preparative importance. It proceeds with a good yield (*ca.* 60%) both in the solid phase (within 3 h at 95°C for $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}](\text{BPh}_4)$ and within 2 h at 105°C for $[\text{Pt}(\text{tx})_3\text{Cl}](\text{BPh}_4)$) and in boiling nitromethane solution.

[†] Author to whom correspondence should be addressed.

[‡] On leave from Department of Chemistry, St. Petersburg State University, 198904 Stary Petergof, Russia.

Reaction between $[\text{Pt}(\text{Me}_2\text{S})_4]^{2+}$ (prepared by halide abstraction from *trans*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{Cl}_2]$ with $\text{AgNO}_3 \cdot \text{Me}_2\text{S}^3$ in 1:2 molar ratio) and NaBPh_4 in acetone-ethanol solution leads to formation of a precipitate, which is insoluble in water, dimethyl formamide, acetone and dichloromethane. Incubation of the precipitate in CD_2Cl_2 for 2 days followed by filtration leads to the product *cis*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{Ph}_2]$, as identified by its ^1H NMR spectrum.

Interaction between $\text{K}_2[\text{PtCl}_4]$ and an excess of Me_2S in water at 25°C for 2 days yields a solution which contains $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}]^+$ and a precipitate which is a mixture of *cis*- and *trans*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{Cl}_2]$. Filtration of the precipitate followed by the addition of KPF_6 to the filtrate gives a precipitate of $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}](\text{PF}_6)$ (yield 19%). Metathesis of a water-acetone solution of $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}](\text{PF}_6)$ with an ethanolic solution of NaBPh_4 in a 1:1 molar ratio results in precipitation of the previously characterized $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}](\text{BPh}_4)$ compound.³ The latter product as well as $[\text{Pt}(\text{tx})_3\text{Cl}](\text{BPh}_4)$ can also be obtained according to the following scheme (References, 4–6):



Our preliminary experiments clearly show that the heterogeneous reaction of $[\text{Pt}(\text{Me}_2\text{S})_2\text{Cl}_2]$ with one mole of AgBPh_4 in dichloromethane suspension results in the formation of *trans*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{ClPh}]$. This means that AgBPh_4 can serve as a mild phenylation reagent for platinum(II) thioether chloro complexes.

Characterization of compounds. Analytical data for *trans*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{ClPh}]$: M.P. 137°C . Anal. Calc. for $\text{C}_{10}\text{H}_{17}\text{ClPtS}_2$: C, 27.8%; H, 4.0%; Cl, 8.2%. Found: C, 27.4%; H, 4.0%; Cl, 8.4%. ^1H NMR spectrum (CD_2Cl_2) δ_{SMe} , ppm: 2.33 ($^3J_{\text{PtH}}$ 57 Hz). IR spectrum in nujol, cm^{-1} : 332 $\nu(\text{Pt-Cl})$. We have also characterized the structure of this compound by X-ray crystallography: space group $P2_1/n$ with $a = 10.106(1)$, $b = 13.046(3)$, $c = 20.939(2)$ Å, $V = 2752.4(6)$ Å³, $\beta = 94.39(1)^\circ$, $D_x = 2.08 \text{ g cm}^{-3}$, $R = 0.038$.

Analytical data for *trans*- $[\text{Pt}(\text{tx})_2\text{ClPh}]$: M.P. 110°C . Anal. Calc. for $\text{C}_{14}\text{H}_{21}\text{ClO}_2\text{PtS}_2$: Cl, 6.9%; Pt, 37.8%. Found: Cl, 6.7%; Pt, 37.7%. IR spectrum in nujol, cm^{-1} : 335 $\nu(\text{Pt-Cl})$.

^1H NMR data for *cis*- $[\text{Pt}(\text{Me}_2\text{S})_2\text{Ph}_2]$ δ_{SMe} , ppm: 2.13 ($^3J_{\text{PtH}}$ 24 Hz) [*lit.*: 2.1 ppm ($^3J_{\text{PtH}}$ 24 Hz)⁷, 2.09 ppm ($^3J_{\text{PtH}}$ 23.6 Hz)⁸].

Analytical data for $[\text{Pt}(\text{Me}_2\text{S})_3\text{Cl}](\text{PF}_6)$: Anal. Calc. for $\text{C}_6\text{H}_{18}\text{ClF}_6\text{PPtS}_3$: C, 12.8%; H, 3.2%; S, 17.1%. Found: C 12.6%; H, 3.2%; S, 17.3%. M.P. $126\text{--}130^\circ\text{C}$ (capillary). IR spectrum (KBr pellet, cm^{-1}): 336 s $\nu(\text{Pt-Cl})$. ^1H NMR spectrum (acetone- d_6 ; δ , ppm): 2.78 ($^3J_{\text{PtH}}$ 51.5 Hz, 3H) and 2.70 ($^3J_{\text{PtH}}$ 60.5 Hz, 6H). X-ray single-crystal diffraction data, two independent crystal determinations: space

group $P2_1/a$ with $a = 8.263(2)/8.258(2)$, $b = 23.970(5)/23.937(6)$, $c = 8.544(2)/8.540(1)$ Å, $V = 1642.5(4)/1638.1(4)$ Å³, $\beta = 103.93(1)/103.96(2)^\circ$, $D_x = 2.27/2.28$ g cm⁻³, $R = 0.068/0.073$.

Compounds of the type [Pt(thioether)₃Cl](SO₃CF₃) have been structurally characterized. [Pt(Me₂S)₃Cl](SO₃CF₃) crystallizes in space group $P\bar{1}$ with $a = 8.2320(9)$, $b = 9.207(1)$, $c = 23.670(6)$ Å, $V = 1719.3(4)$ Å³, $\alpha = 91.54(2)$, $\beta = 98.37(1)$, $\gamma = 103.898(9)^\circ$, $D_x = 2.19$ g cm⁻³, $R = 0.030$. [Pt(tx)₃Cl](SO₃CF₃) crystallizes in space group $P\bar{1}$ with $a = 9.865(1)$, $b = 10.3469(8)$, $c = 11.568(1)$ Å, $V = 1114.8(1)$ Å³, $\alpha = 97.131(7)$, $\beta = 97.144(9)$, $\gamma = 105.299(7)$, $D_x = 2.062$ g cm⁻³, $R = 0.044$.

ACKNOWLEDGEMENT

V. Yu. Kukushkin is grateful to the Royal Swedish Academy of Sciences and the Academy of Sciences of Russia for financial support of his stay at the Chemical Center of the University of Lund. This research was also supported by a grant from the Swedish Natural Science Research Council.

REFERENCES

1. K. Siegmann, P. S. Pregosin and L. M. Venanzi, *Organometallics*, **8**, 2659 (1989); B. Crociani, F. Di Bianca, P. Uguagliati, L. Canovese and A. Berton, *J. Chem. Soc., Dalton Trans.*, **1991**, 71; P.-K. F. Chin and F. R. Hartley, *Inorg. Chem.*, **15**, 982 (1976); R. A. Alcock, F. R. Hartley and D. E. Rogers, *J. Chem. Soc., Dalton Trans.*, **1973**, 1070.
2. J. A. Davies and F. R. Hartley, *Chem. Rev.*, **81**, 79 (1981); V. Yu. Kukushkin and Yu. N. Kukushkin, *Theory and Practice of Coordination Compounds Synthesis*, Nauka: Leningrad, 1990.
3. H. Morita and J. C. Bailar, Jr., *Inorg. Synth.*, **22**, 126 (1983).
4. For synthesis of [Pt(NCET)₃Cl₂] see: V. V. Lebedinskii and V. A. Golovnya, *Ann. secteur platine, Inst. chim. gén.*, **18**, 38 (1945); *Chem. Abstr.*, **41**: 6187d.
5. For synthesis of [Pt(NCET)₄](SO₃CF₃)₂ see: V. Yu. Kukushkin, Å. Oskarsson and L. I. Elding, *Inorg. Synth.*, submitted for publication.
6. X-ray crystal structures of [Pt(Me₂S)₄](SO₃CF₃)₂, [Pt(thioxane)₄](SO₃CF₃)₂ and ¹H NMR spectra of these compounds were reported: Z. Bugarcic, B. Norén, Å. Oskarsson, C. Stålhandske and L. I. Elding, *Acta Chem. Scand.*, **45**, 361 (1991); U. Frey, S. Elmroth, B. Moullet, L. I. Elding and A. E. Merbach, *Inorg. Chem.*, **30**, 5033 (1991).
7. N. Hadj-Bagheri and R. J. Puddephatt, *Polyhedron*, **7**, 2695 (1988).
8. G. Alibrandi, G. Bruno, S. Lanza, D. Minniti, R. Romeo and M. Tobe, *Inorg. Chem.*, **26**, 185 (1987).