


This article was downloaded by:

On: 29 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

INTRAMOLECULAR LIGAND EXCHANGE OF PENTACOORDINATED ANIONIC SILICATES, RSiF_4^- , VIA ^{29}Si AND ^{19}F NMR SPECTROSCOPY, SOLUTION AND SOLID-STATE STRUCTURES

Stephen E. Johnson^a, Roberta O. Day^a, Robert R. Holmes^a

^a Contribution from the Department of Chemistry, University of Massachusetts, Amherst, Massachusetts

To cite this Article Johnson, Stephen E. , Day, Roberta O. and Holmes, Robert R.(1995) 'INTRAMOLECULAR LIGAND EXCHANGE OF PENTACOORDINATED ANIONIC SILICATES, RSiF_4^- , VIA ^{29}Si AND ^{19}F NMR SPECTROSCOPY, SOLUTION AND SOLID-STATE STRUCTURES', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 99: 1, 165 – 183

To link to this Article: DOI: 10.1080/10426509508031344

URL: <http://dx.doi.org/10.1080/10426509508031344>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

INTRAMOLECULAR LIGAND EXCHANGE OF PENTACOORDINATED ANIONIC SILICATES, RSiF_4^- , VIA ^{29}Si AND ^{19}F NMR SPECTROSCOPY, SOLUTION AND SOLID-STATE STRUCTURES^{1,2}

STEPHEN E. JOHNSON,³ ROBERTA O. DAY, and
ROBERT R. HOLMES*

*Contribution from the Department of Chemistry, University of Massachusetts,
Amherst, Massachusetts 01003*

(Received April 25, 1989)

New pentacoordinated anionic silicates were synthesized as potassium 18-crown-6 salts, $[\text{RSiF}_4][\text{K}-18\text{-crown-6}]$, where R = 2,4,6-tri-*tert*-butylphenyl (TTBP) (1), mesityl (2), *n*-propyl (3), 1,2-ethanediyl-bis(tetrafluorosilicate) (4), *tert*-butyl (5), cyclohexyl (6), benzyl (7), *p*-chlorophenyl (8), *p*-tolyl (9), *m*-tolyl (10), and *o*-tolyl (11), by fluoride addition to the respective fluorosilane, RSiF_3 , ^{19}F and ^{29}Si variable-temperature NMR studies provide the first example of an anionic RSiF_4^- derivative whose intramolecular ligand exchange is “stopped” in solution. This occurs for 1 and indicates the magnitude of steric effects required to accomplish this. A comparison is made with substituent effects introduced in isoelectronic phosphoranes, RPF_4 , to stop ligand exchange on the NMR time scale. NMR data of all other silicates indicate rapid intramolecular exchange down to the lowest temperatures studied. A barrier energy for fluorine exchange in 1 of 12.8 kcal/mol was obtained. The molecular structures of 1 and 2 are reported and indicate the effects of steric crowding on distortion of the resulting trigonal bipyramidal geometries, more so with 1 than 2. The latter agrees with trends in ^{19}F and ^{29}Si chemical shifts comparisons within the series RSiF_4^- and RSiF_3 . Silicate 1 crystallizes in the orthorhombic space group $P2_12_12$, with $a = 17.563$ (4) Å, $b = 22.528$ (5) Å, $c = 9.800$ (2) Å, and $Z = 4$. The mesityl derivative 2 crystallizes in the monoclinic space group $P2_1/m$ with $a = 8.158$ (3) Å, $b = 14.847$ (7) Å, $c = 10.664$ (1) Å, $\beta = 91.24$ (2)°, and $Z = 2$. The final conventional unweighted residuals are 0.041 (1) and 0.040 (2).

INTRODUCTION

Analogous to anionic diorganotrifluorosilicates,^{4a} R_2SiF_3^- , the related tetrafluorosilicates, RSiF_4^- , have been increasingly invoked as the principal intermediate in accounting for a variety of organic transformations. In a group-transfer reaction, Kira *et al.*^{4b} proposed activation of allyltrifluorosilane by fluoride ion to form the anionic allyltetrafluorosilicate, which in turn facilitates allyl transfer to the aldehyde substrate. We have carried out ab initio molecular orbital calculations on model hydridofluorosilanes, $\text{SiH}_x\text{F}_{4-x}$, and anionic hydridofluorosilicates, $[\text{SiH}_x\text{F}_{5-x}]^-$.⁵ The study indicates that enhanced reactivity of pentacoordinated species is associated with an increase in leaving group ability relative to that in tetracoordinate members.

A study of the structural features and properties of isolatable members of the tetrafluorosilicate series RSiF_4^- should be pursued to gain an understanding of steric and electronic effects that may be controlling specific reaction mechanisms.

Reprinted with permission from *Inorg. Chem.*, 28, 3182 (1989). Copyright 1989 American Chemical Society.

However, a limited amount of such information is available on this class of compounds. Klanberg and Muetterties⁶ characterized tetraalkylammonium salts of the anions MeSiF_4^- , $n\text{-BuSiF}_4^-$, and PhSiF_4^- . From their ^1H and ^{19}F NMR spectra, trigonal-bipyramidal geometries undergoing rapid intramolecular fluorine exchange were proposed.

An IR study⁷ of $[\text{PhSiF}_4][\text{Et}_4\text{N}]$ and an IR matrix-isolation study of $[\text{MeSiF}_4][\text{Cs}]^8$ also are consistent with a trigonal-bipyramidal geometry. Confirmation of this stereochemistry was provided by Schomburg,⁹ who performed an X-ray diffraction study of $[\text{PhSiF}_4][n\text{-Pr}_4\text{N}]$ and found the anion to be trigonal bipyramidal with the phenyl group occupying an equatorial site. A hydrolysis study^{1b} of dimesityldifluorosilane in the presence of $\text{Et}_4\text{NF} \cdot 2\text{H}_2\text{O}$, afforded $[\text{MeSiF}_4][\text{Et}_4\text{N}]$ as one of the products. Like the derivatives Klanberg and Muetterties⁶ isolated, NMR spectra of the MeSiF_4^- anion at room temperature^{1b} and below² indicated only the presence of equivalent fluorine atoms that exhibited spin–spin coupling to silicon.

Marat and Janzen¹⁰ further examined fluorine exchange in $[\text{MeSiF}_4][n\text{-Pr}_4\text{N}]$ and $[\text{PhSiF}_4][n\text{-Pr}_4\text{N}]$ and confirmed the exchange to be rapid on the NMR time scale under all experimental conditions. It was postulated that the lowest energy pathway of exchange involved Lewis acid–base interactions and rapid equilibration of five- and six-coordinate geometries.^{10,11} The study, however, provided no direct evidence in favor of this mechanism, nor did it provide any direct evidence that would rule out an intramolecular exchange via a Berry pseudorotation process.¹²

The Berry pseudorotation process is the most plausible low-energy pathway to account for ligand exchange in the anions, RSiF_4^- , as it is the only exchange mechanism receiving extensive verification for main-group elements.¹³ This is particularly meaningful for comparison with the isoelectronic phosphoranes.¹⁴ Both phosphoranes^{15,16} and anionic five-coordinated silicates exhibit a continuous range of solid-state structures ranging from trigonal-bipyramidal to rectangular pyramidal along the Berry coordinate.^{17–21}

In order to provide a firm basis to discuss mechanistic criteria for reactions involving RSiF_4^- species, the present study was undertaken. Herein, the synthesis, structural characterization, and dynamic ^1H , ^{19}F , and ^{29}Si NMR behavior for a series of 10 new monoorganotetrafluorosilicate anions as potassium 18-crown-6 salts, $[\text{RSiF}_4][\text{K-18-crown-6}]$, are reported. Similar studies on $[\text{PhSiF}_4][\text{K-18-crown-6}]$ are presented. The molecular structures of $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ (TTBP = 2,4,6-tri-*tert*-butyl-phenyl) and $[\text{MesSiF}_4][\text{K-18-crown-6}]$ are provided by X-ray diffraction analysis and serve to give insight into the solution-state NMR interpretations, particularly regarding implied steric effects. This study provides the first example, the $[(\text{TTBP})\text{SiF}_4]^-$ anion, where intramolecular fluorine exchange is “stopped” on the NMR time scale for a monoorganotetrafluorosilicate anion. An energy of activation for the exchange process also is obtained.

EXPERIMENTAL

Vacuum-line and Schlenk techniques were used for the preparation, purification, and manipulation of reactants and products. All solvents were freshly distilled and stored over Ar or N₂. Dichloromethane (Fisher) was dried over CaH₂. Toluene (Fisher) was dried over P₂O₅. Ethyl ether and tetrahydrofuran were dried over Na and benzophenone. CDCl₃ and CD₂Cl₂ were obtained from Aldrich. Grignard reagents prepared and purchased (Aldrich) were stored under Ar prior to use. Organotrichlorosilanes

(R = Mes, *o*-Tol, *m*-Tol) were either purchased (Petrarch Systems) or synthesized by reaction of the corresponding Grignard reagent with a 4-fold excess of SiCl_4 .

^{29}Si (59.59 MHz), ^{19}F (282.2 MHz), and ^1H (299.1 MHz) pulse Fourier transform NMR spectra were recorded on a Varian Associates Corp. XL-300 spectrometer. ^1H and ^{29}Si chemical shifts are reported relative to TMS in ppm (internal standard, ^1H ; external standard, ^{29}Si). ^{19}F chemical shifts were referenced relative to CFCl_3 in ppm (internal standard). NMR parameters were determined at room temperature unless otherwise stated. Variable-temperature experiments were carried out in CDCl_3 and CD_2Cl_2 solutions. Temperature calibration was accomplished by using a standard ethylene glycol sample, with a calibration error of ± 0.5 °C. ^{29}Si NMR experiments were performed with proton decoupling, using standard ^{13}C pulse programs and/or INEPT programs.²² Calculated spectra for $[(\text{TTBP})\text{SiF}_4]\text{[K-18-crown-6]}$ were produced with the program DNMR-4²³ using a Fortran H compiler on a microcomputer. The ADPLOT²³ program was rewritten, with a Hewlett Packard plotter driver, to produce simulated spectral plots.²⁴ The pseudo-first-order rate constants for exchange were obtained by fitting the calculated to the experimental spectra.

Syntheses. (a) *(Tri-tert-butylphenyl)trifluorosilane*, $(\text{TTBP})\text{SiF}_4$. Preparation of this organofluorosilane was reported recently^{25,26} from the reaction of SiF_4 with a solution of 2,4,6-tri-*tert*-butylphenyl bromide²⁷ and *n*-BuLi in THF: mp 65–68 °C (lit.²⁶ mp 71 °C). ^1H (CDCl_3): 1.31 (s, *p*-*t*-Bu, 9 H); 1.46 (s, *o*-*t*-Bu, 18 H); 7.44 (s, aromatic ring protons, 2 H). ^{19}F (CDCl_3): –124.4 (d_{Si} , $J_{\text{SiF}} = 266$ Hz). ^{29}Si (CDCl_3): –61.30 (q_{F} , $J_{\text{SiF}} = 262$ Hz).

(b) *Potassium 18-Crown-6 (Tri-tert-butylphenyl)tetrafluorosilicate* [$\text{K-18-crown-6}/[(\text{TTBP})\text{SiF}_4]$] (1). *Tri-tert-butyltrifluorosilane* (1.96 g, 5.93 mmol), 18-crown-6 (1.57 g, 5.93 mmol), and KF (0.344 g, 5.93 mmol) were reacted in 20 mL of toluene at 25 °C. The reactants were stirred rapidly for 24 h. Precipitate formation was observed shortly after the reactants were combined. The resulting white precipitate was collected by vacuum filtration. Recrystallization from methylene chloride afforded 3.11 g (80% yield) of [$\text{K-18-crown-6}/[(\text{TTBP})\text{SiF}_4]$] as colorless crystals, mp 184–188 °C dec. The crystals lose solvent (CH_2Cl_2) on exposure to air, forming a powder, mp 188 °C dec. Anal. Calcd for $\text{C}_{30}\text{H}_{53}\text{SiF}_4\text{O}_6\text{K}$: C, 50.45; H, 7.53. Found: C, 50.37; H, 7.40. ^1H (CD_2Cl_2): 1.28, 1.31 (s, *o*-*t*-Bu, 18 H); 1.55 (s, *p*-*t*-Bu, 9 H); 2.29 (s, CH_2Cl_2 , 2 H); 3.57 (s, 18-crown-6, 24 H), 7.11–7.38 (m's, aromatic ring protons, 2 H). ^{19}F (CD_2Cl_2 , 21.0 °C): –95.08 (s). ^{19}F (CD_2Cl_2 , –87.8 °C): –77.98 (t_{F} , $J_{\text{FeF}_3} = 29.6$ Hz, 2 F_{ax} , $J_{\text{SiF}_{\text{ax}}} = 248$ Hz); –112.7 (t_{F} , $J_{\text{FeF}_3} = 29.6$ Hz, 2 F_{eq} , $J_{\text{SiF}_{\text{eq}}} = 191$ Hz). ^{29}Si (CD_2Cl_2 , 21.0 °C): –114.1 (p_{F} , $J_{\text{SiF}_{\text{av}}} = 224$ Hz). ^{29}Si (CD_2Cl_2 , –107.2 °C): –114.1 (t of t, $J_{\text{SiF}_{\text{ax}}} = 248$ Hz, $J_{\text{SiF}_{\text{eq}}} = 193$ Hz).

(c) *Mesityltrichlorosilane*, MesSiCl_3 . MesSiCl_3 was first prepared by the reaction of MesLi with 2 equiv of SiCl_4 (48% yield).²⁸ MesSiCl_3 was prepared for this study by the reaction of MesMgBr with excess SiCl_4 (51% yield): bp 96–102 °C (1.5 mm); mp 60–62 °C (lit.²⁸ mp 55–59 °C). Anal. Calcd for $\text{C}_9\text{H}_{11}\text{SiCl}_3$: C, 42.61; H, 4.38. Found: C, 42.49; H, 4.50. ^1H (CDCl_3): 2.29 (s, *p*-Me, 3 H); 2.66 (s, *o*-Me, 6 H); 6.90 (s, aromatic ring protons, 2 H). ^{29}Si (CDCl_3): –3.391 (s).

(d) *Mesityltrifluorosilane*, MesSiF_3 . A solution of CuF_2 (7.87 g, 77.5 mmol) in 50 mL of ethyl ether was prepared. A solution of MesSiCl_3 (13.1 g, 51.6 mmol) in 20 mL of Et_2O was slowly added (dropwise) to the solution containing CuF_2 . The reaction flask was immersed in an ice bath (0 °C) during the addition (0.5 h). The reaction was stirred at 25 °C for 24 h. The reaction mixture was distilled directly at atmospheric pressure to remove excess ether. The residue was distilled at reduced pressure to produce 7.12 g (67% yield) of MesSiF_3 , bp 37–40 °C (0.10 mm). Anal. Calcd for $\text{C}_9\text{H}_{11}\text{SiF}_3$: C, 52.91; H, 5.44. Found: C, 53.50; H, 5.39. ^1H (CDCl_3): 2.26 (s, *p*-Me, 3 H); 2.43 (s, *o*-Me, 6 H); 6.78 (s, aromatic ring protons, 2 H). ^{19}F (CDCl_3): –130.6 (d_{Si} , $J_{\text{SiF}} = 274$ Hz). ^{29}Si (CDCl_3): –69.19 (q_{F} , $J_{\text{SiF}} = 274$ Hz).

(e) *Potassium 18-Crown-6-Mesityltetrafluorosilicate*, [$\text{K-18-crown-6}/[\text{MesSiF}_4]$] (2). *Mesityltrifluorosilane* (2.37 g, 11.6 mmol), 18-crown-6 (3.06 g, 11.6 mmol), and KF (0.673 g, 11.6 mmol) were reacted in 30 mL of toluene at 25 °C. The reaction mixture was vigorously stirred for 24 h. Precipitate formation was observed shortly after the reactants were combined (15 min). The white precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 78–81 °C. Anal. Calcd for $\text{C}_{21}\text{H}_{35}\text{SiF}_4\text{O}_6\text{K}$: C, 47.88; H, 6.71. Found: C, 47.45; H, 6.82. ^1H (CDCl_3): 2.17 (s, *p*-Me, 3 H); 2.48 (s, *o*-Me, 6 H); 3.52 (s, 18-crown-6, 24 H); 6.68 (s, aromatic ring protons, 2 H). ^{19}F (CDCl_3 , 21.0 °C): –111.8 (d_{Si} , $J_{\text{SiF}_{\text{av}}} = 219$ Hz). ^{19}F (CD_3CN , 21 °C): –109.0 (d_{Si} , $J_{\text{SiF}_{\text{av}}} = 217$ Hz). ^{29}Si (CDCl_3 , 21 °C): –120.1 (p_{F} , $J_{\text{SiF}_{\text{av}}} = 219$ Hz). ^{29}Si (CD_3CN , 21 °C): –121.3 (p_{F} , $J_{\text{SiF}_{\text{av}}} = 217$ Hz).

(f) *1,1-Difluorosilacyclobutane*, $(\text{CH}_2)_2\text{SiF}_2$. The preparation of this organofluorosilane is a modification of an earlier procedure,^{29a} using CuF_2 instead of SBF_3 . A solution of CuF_2 (10.8 g, 10.6 mmol) in 40 mL of diphenyl ether (Ph_2O) was prepared. A solution of 1,1-dichlorosilacyclobutane (10.0 g, 7.09

mmol) in 20 mL of $p\text{H}_2\text{O}$ was added to the CuF_2 solution (1 h). The reaction flask was immersed in a water bath (10 °C) during the addition. The reaction was warmed to 25 °C and then stirred overnight. The mixture was distilled directly to yield $(\text{CH}_2)_3\text{SiF}_2$, bp 39–41 °C (lit.^{29a} bp 36–38 °C (760 mm)). A total of 4.91 g (64% yield) of $(\text{CH}_2)_3\text{SiF}_2$ was collected. ^1H (CDCl_3): 1.760 (m, aliphatic ring protons, 6 H). ^{19}F (CDCl_3): –122.4 (d_{Si} , $J_{\text{SiF}} = 380$ Hz). Both the ^{19}F and ^1H NMR spectra recorded were consistent with the spectra obtained in a previous study.^{29b} ^{29}Si (CDCl_3): –19.16 (t_{F} , $J_{\text{SiF}} = 380$ Hz).

(g) *Potassium 18-Crown-6 n-Propyltetrafluorosilicate, [K-18-crown-6]/[n-PrSiF₄] (3)*. 1,1-Difluorosilacyclobutane (0.464 g, 4.29 mmol), 18-crown-6 (1.13 g, 4.29 mmol), and KF (0.249 g, 4.29 mmol) were reacted in 20 mL of toluene at 25 °C. The reactants were vigorously stirred for 24 h. Shortly after the reactants were mixed, precipitate formation was observed (0.5 h). The precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 196–198 °C dec. A total of 0.919 g of [K-18-crown-6]/[n-PrSiF₄] was collected (48% yield). Anal. Calcd for $\text{C}_{15}\text{H}_{31}\text{SiF}_4\text{O}_3\text{K}$: C, 39.98; H, 6.95. Found: C, 39.88; H, 6.80. ^1H (CDCl_3): 0.612 (t , CH_3 —, 3 H); 0.810 (t , — CH_2Si —, 2 H); 1.38 (m, — CH_2 —, 2 H); 3.48 (s, 18-crown-6, 24 H). ^{19}F (CDCl_3): –118.1 (d_{Si} , $J_{\text{SiFav}} = 224$ Hz). ^{29}Si (CDCl_3 , –78.2 °C): –110.8 (p_{F} , $J_{\text{SiFav}} = 225$ Hz).

(h) *1,2-Bis(trifluorosilyl)ethane, $\text{F}_3\text{SiCH}_2\text{CH}_2\text{SiF}_3$* . Preparation of this organofluorosilane had not been reported previously. A solution of CuF_2 (25.6 g, 25.3 mmol) in 40 mL of butyl ether ($n\text{-Bu}_2\text{O}$) was prepared. A second solution, containing $\text{Cl}_3\text{SiCH}_2\text{CH}_2\text{SiCl}_3$ (25.0 g, 8.42 mmol) in 15 mL of $n\text{-Bu}_2\text{O}$, was slowly added to the CuF_2 via an addition funnel. The reaction vessel was immersed in a water bath (0 °C) during the addition (1 h). After the addition was complete, the mixture was allowed to warm to 25 °C and then was stirred for 24 h. The reaction mixture was distilled directly to produce 9.76 g (58% yield) of $\text{F}_3\text{SiCH}_2\text{CH}_2\text{SiF}_3$, bp 45–48 °C. ^1H (CDCl_3): 1.19 (t_{F} — CH_2CH_2 —, $J_{\text{HF}} = 1.1$ Hz, 4 H). ^{19}F (CDCl_3): –139.7 (d_{Si} , $J_{\text{SiF}} = 282$ Hz). ^{29}Si (CDCl_3): –61.11 (q_{F} , $J_{\text{SiF}} = 281$ Hz).

(i) *Potassium 18-Crown-6 1,2-Ethanediylbis(tetrafluorosilicate), [K-18-crown-6]₂[$\text{F}_4\text{SiCH}_2\text{CH}_2\text{SiF}_4$] (4)*. Bis(trifluorosilyl)ethane (4.30 g, 21.7 mmol), 18-crown-6 (11.46 g, 43.4 mmol), and KF (2.52 g, 43.4 mmol) were reacted in toluene solution at 25 °C. The reactants were vigorously stirred for 24 h. Precipitate formation was observed shortly after mixing the reactants (0.5 h). The white precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to give colorless crystals, mp 89–93 °C. A total of 15.2 g of [K-18-crown]₂[$\text{F}_4\text{SiCH}_2\text{SiF}_4$] was isolated (83% yield). Anal. Calcd for $\text{C}_{26}\text{H}_{53}\text{F}_8\text{Si}_2\text{O}_{12}\text{K}_2$: C, 37.03; H, 6.23. Found: C, 36.83; H, 6.08. ^1H (CDCl_3): 0.767 (s, — CH_2CH_2 —, 4 H); 3.48 (s, 18-crown-6, 24 H). ^{19}F (CDCl_3 , –88.2 °C): –118.2 (d_{Si} , $J_{\text{SiFav}} = 227$ Hz). ^{29}Si (CDCl_3): –103.8 (br s). ^{29}Si (CDCl_3 , –98.2 °C): –109.0 (p_{F} , $J_{\text{SiFav}} = 227$ Hz).

(j) *tert-Butyltrifluorosilane, t-BuSiF₃*. This organofluorosilane was synthesized by a previous procedure based on the fluorination of *t*-Bu-SiCl₃ with CuF_2 ²⁵, bp 41–44 °C (lit.²⁵ bp 35–36 °C). ^1H (CDCl_3): 1.147 (q_{F} , $J_{\text{HF}} = 1.10$ Hz, 9 H). ^{19}F (CDCl_3): –149.8 (d_{Si} , $J_{\text{SiF}} = 305$ Hz). ^{29}Si (CDCl_3): –62.10 (q_{F} , $J_{\text{SiF}} = 304$ Hz).

(k) *Potassium 18-Crown-6 tert-Butyltetrafluorosilicate, [K-18-crown-6]/[t-BuSiF₄] (5)*. *tert*-Butyltrifluorosilane (1.46 g, 10.3 mmol), 18-crown-6 (2.71 g, 10.3 mmol), and KF (0.596 g, 10.3 mmol) were reacted in 30 mL of toluene at 25 °C with brisk stirring. Precipitate formation was observed shortly after mixing the reactants (1 h). The mixture was allowed to stir for 24 h. The white precipitate was collected by vacuum filtration. A total of 3.76 g of [K-18-crown-6]/[t-BuSiF₄] was collected (79% yield): mp 185 °C dec. The compound was recrystallized from methylene chloride to give colorless crystals: mp 92–94 °C; dec pt 185 °C. Anal. Calcd for $\text{C}_{16}\text{H}_{33}\text{SiF}_4\text{O}_6\text{K}$: C, 41.35; H, 7.17. Found: C, 41.21; H, 7.01. ^1H (CDCl_3): 1.07 (s, *t*-Bu, 9 H), 3.68 (s, 18-crown-6, 24 H). ^{19}F (CDCl_3 , –27.2 °C): –123.5 (d_{Si} , $J_{\text{SiFav}} = 235$ Hz). ^{29}Si (CDCl_3 , –58.2 °C): –114.8 (p_{F} , $J_{\text{SiFav}} = 237$ Hz).

(l) *Cyclohexyltrifluorosilane, C₆H₁₁SiF₃*. This organofluorosilane was first prepared by using SbF₃.^{30,31} A solution of CuF_2 (17.5 g, 17.2 mmol) in 50 mL of phenyl ether was prepared. A solution of C₆H₁₁SiCl₃ (25.0 g, 11.5 mmol) in 20 mL of Ph₂O was slowly added to the CuF_2 via an addition funnel. The reaction funnel was immersed in a water bath during the addition (0.5 h). The reaction was allowed to stir for 24 h. The mixture was directly distilled to produce 13.3 g (69% yield) of C₆H₁₁SiF₃, bp 85–89 °C (lit.³⁰ bp 105 °C). ^1H (CDCl_3): 1.18–1.91 (m's, cyclohexyl ring protons, 11 H). ^{19}F (CDCl_3): –144.2 (d_{Si} , $J_{\text{SiF}} = 296$ Hz). ^{29}Si (CDCl_3): –62.60 (q_{F} , $J_{\text{SiF}} = 296$ Hz).

(m) *Potassium 18-Crown-6 Cyclohexyltetrafluorosilicate, [K-18-crown-6]/[C₆H₁₁SiF₄] (6)*. Cyclohexyltrifluorosilane (1.22 g, 7.23 mmol), 18-crown-6 (1.91 g, 7.23 mmol), and KF (0.419 g, 7.23 mmol) were reacted in 30 mL of toluene at 25 °C with rapid stirring. Precipitate formation was observed shortly (1 h) after the reactants were mixed. The white precipitate was collected by vacuum filtration. The

compound was recrystallized from methylene chloride to yield colorless crystals: mp 139–143 °C, dec pt 188–190 °C. A total of 2.48 g (70% yield) of [K-18-crown-6][$\text{C}_6\text{H}_{11}\text{SiF}_4$] was collected. Anal. Calcd for $\text{C}_{18}\text{H}_{35}\text{SiF}_4\text{O}_6\text{K}$: C, 44.05; H, 7.20. Found: C, 44.97; H, 6.99. ^1H (CDCl_3): 0.85–1.95 (multiplets, cyclohexyl ring protons, 11 H), 3.64 (s, 18-crown-6, 24 H). ^{19}F (CDCl_3 , –58.3 °C): –123.8 (d_{Si} , $J_{\text{SiFav}} = 230$ Hz). ^{29}Si (CDCl_3 , –68.7 °C): –114.7 (p_{F} , $J_{\text{SiFav}} = 232$ Hz).

(n) *Benzyltrifluorosilane, BzSiF₃*. BzSiF₃ was prepared by fluorinating BzSiCl₃ with ZnF₂ in Et₂O solution by following a related literature method³²; bp 125–126 °C (lit.³² bp 134–135 °C). ^1H (CDCl_3): 2.51 (t_{F} , methylene protons, $J_{\text{HF}} = 0.9$ Hz, 2 H): 7.03–7.65 (m, aromatic ring protons, 4 H). ^{19}F (CDCl_3): –138.9 (d_{Si} , $J_{\text{SiF}} = 281$ Hz). ^{29}Si (CDCl_3): –64.13 (q_{F} , $J_{\text{SiF}} = 282$ Hz).

(o) *Potassium 18-Crown-6 Benzyltetrafluorosilicate, [K-18-crown-6]/[BzSiF₄]* (7). Benzyltrifluorosilane (1.85 g, 10.5 mmol), 18-crown-6 (2.78 g, 10.5 mmol), and KF (0.610 g, 10.5 mmol) were reacted in 20 mL of toluene at 25 °C. The reaction mixture was vigorously stirred for 24 h. Precipitate formation was observed immediately after the reactants were combined. The white precipitate was collected by vacuum filtration. A total of 4.82 g (92% yield) of [K-18-crown-6][BzSiF₄] was isolated. The compound was recrystallized from methylene chloride to produce colorless crystals, mp 169–172 °C dec. Anal. Calcd for $\text{C}_{19}\text{H}_{31}\text{SiF}_4\text{O}_6\text{K}$: C, 45.76; H, 6.28. Found: C, 45.13; H, 6.24. ^1H (CDCl_3): 2.24 (s, methylene protons, 2 H); 3.53 (s, 18-crown-6, 24 H); 6.76–7.43 (m, aromatic ring protons, 4 H). ^{19}F (CD_2Cl_2 , –98.2 °C): –117.3 (d_{Si} , $J_{\text{SiFav}} = 215$ Hz). ^{29}Si (CD_2Cl_2 , –98.2 °C): –116.4 (p_{F} , $J_{\text{SiFav}} = 218$ Hz).

(p) *(p-Chlorophenyl)trifluorosilane, p-ClC₆H₄SiF₃*. This fluorosilane was prepared from the reaction of boron trifluoride etherate with *p*-ClC₆H₄Si(OEt)₃ following a modification of a literature method³³; bp 137–141 °C (lit.³⁴ 132–134 °C). ^1H (CDCl_3): 7.37, 7.39, 7.68, 7.70 (A_2B_2 , aromatic ring protons, 4 H). ^{19}F (CDCl_3): –140.8 (d_{Si} , $J_{\text{SiF}} = 252$ Hz). ^{29}Si (CDCl_3): –73.12 (q_{F} , $J_{\text{SiF}} = 251$ Hz).

(q) *Potassium 18-Crown-6 (p-Chlorophenyl)tetrafluorosilicate, [K-18-crown-6]/[p-ClC₆H₄SiF₄]* (8). (p-Chlorophenyl)trifluorosilane (1.50 g, 7.63 mmol), 18-crown-6 (2.02 g, 7.63 mmol), and KF (0.443 g, 7.63 mmol) were reacted in 20 mL of toluene at 25 °C. The reactants were vigorously stirred for 24 h. Precipitate formation was observed shortly after the reactants were combined (0.5 h). The white precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 189–192 °C dec. Anal. Calcd for $\text{C}_{30}\text{H}_{28}\text{SiF}_4\text{ClO}_6\text{K}$: C, 54.33; H, 4.26. Found: C, 53.99; H, 4.21. ^1H (CDCl_3): 3.44 (s, 18-crown-6, 24 H); 7.12, 7.14, 7.93, 7.96 (A_2B_2 , aromatic ring protons, 4 H). ^{19}F (CDCl_3 , –38.2 °C): –120.9 (d_{Si} , $J_{\text{SiFav}} = 212$ Hz). ^{29}Si (CDCl_3 , –58.3 °C): –125.2 (p_{F} , $J_{\text{SiFav}} = 213$ Hz).

(r) *p-Tolyltrifluorosilane, p-TolSiF₃*. Preparation of this organofluorosilane previously reported^{35,36} from the reaction of CuF₂ with *p*-TolSiCl₃ in Et₂O solution was followed here: bp 75–77 °C (85 mm) (lit.³⁵ bp 129–132 °C). ^1H (CDCl_3): 2.36 (s, *p*-Me, 3 H), 7.27, 7.29, 7.58, 7.61 (A_2B_2 , aromatic ring protons, 4 H). ^{19}F (CDCl_3): –141.4 (d_{Si} , $J_{\text{SiF}} = 266$ Hz). ^{29}Si (CDCl_3): –74.00 (q_{F} , $J_{\text{SiF}} = 267$ Hz).

(s) *Potassium 18-Crown-6 p-Tolyltetrafluorosilicate, [K-18-crown-6]/[p-TolSiF₄]* (9). *p*-Tolyltrifluorosilane (1.50 g, 8.53 mmol), 18-crown-6 (2.26 g, 8.53 mmol), and KF (0.469 g, 8.53 mmol) were reacted in 30 mL of toluene at 25 °C, with rapid stirring. Precipitate formation was observed shortly (1 h) after the reactions were combined. The white precipitate was collected by vacuum filtration. A total 4.02 g (94% yield) of [K-18-crown-6][*p*-TolSiF₄] was isolated. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 160–165 °C. Anal. Calcd for $\text{C}_{19}\text{H}_{31}\text{SiF}_4\text{O}_6\text{K}$: C, 45.76; H, 6.28. Found: C, 45.35; H, 6.09. ^1H (CDCl_3): 2.26 (s, *p*-Me, 3 H), 3.2 (s, 18-crown-6, 24 H), 7.02, 7.05, 7.83, 7.86 (A_2B_2 , aromatic ring protons, 4 H). ^{19}F (CDCl_3): –122.4 (d_{Si} , $J_{\text{SiFav}} = 210$ Hz). ^{29}Si (CDCl_3 , –62.3 °C): –126.1 (p_{F} , $J_{\text{SiFav}} = 209$ Hz).

(t) *m-Tolyltrifluorosilane, m-TolSiF₃*. *m*-TolSiF₃ was obtained by a literature reaction similar to that used for the preparation of BzSiF₃; bp 127–130 °C (lit.³⁴ bp 123–125 °C). ^1H (CDCl_3): 2.47 (s, *m*-Me, 3 H); 7.21–7.92 (m, aromatic ring protons, 4 H). ^{19}F (CDCl_3): –140.2 (d_{Si} , $J_{\text{SiF}} = 268$ Hz). ^{29}Si (CDCl_3): –72.59 (q_{F} , $J_{\text{SiF}} = 268$ Hz).

(u) *Potassium 18-Crown-6 m-Tolyltetrafluorosilicate, [K-18-crown-6]/[m-TolSiF₄]* (10). *m*-Tolyltrifluorosilane (2.30 g, 13.1 mmol), 18-crown-6 (3.45 g, 13.1 mmol), and KF (0.758 g, 13.1 mmol) were reacted in 30 mL of toluene at 25 °C. The mixture was rapidly stirred for 24 h. Precipitate formation was observed shortly after the reactants were combined (0.5 h). The white precipitate was collected by vacuum filtration. A total of 5.66 g (87% yield) of [K-18-crown-6][*m*-TolSiF₄] was collected. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 141–145 °C. Anal. Calcd for $\text{C}_{19}\text{H}_{31}\text{SiF}_4\text{O}_6\text{K}$: C, 45.68; H, 6.28. Found: C, 45.62; H, 6.09. ^1H (CDCl_3): 2.39 (s, *m*-Me, 3

H); 6.80–7.75 (m's, aromatic ring protons, 4 H). ^{19}F (CDCl₃, –58.2 °C): –119.6 (d_{Si}, J_{SiFav} = 214 Hz). ^{29}Si (CDCl₃, –58.2 °C): –124.3 (p_F, J_{SiFav} = 214 Hz).

(v) *o-Tolyltrifluorosilane, o-TolSiF₃*. This silane was prepared by following a literature³² reaction similar to that used for the preparation of BzSiF₃; bp 125 °C (lit.³⁷ bp 130.8 °C). ^1H (CDCl₃): 2.55 (s, *o*-Me, 3 H); 7.20–7.80 (m's, aromatic ring protons, 4 H). ^{19}F (CDCl₃): –138.9 (d_{Si}, J_{SiF} = 270 Hz). ^{29}Si (CDCl₃): –71.14 (q_F, J_{SiF} = 270 Hz).

(w) *Potassium 18-Crown-6 o-Tolyltetrafluorosilicate, [K-18-crown-6][o-TolSiF₄] (11)*. *o*-Tolyltrifluorosilane (2.45 g, 13.9 mmol), 18-crown-6 (3.68 g, 13.9 mmol), and KF (0.809 g, 13.9 mmol) were reacted in 30 mL of toluene at 25 °C. The reaction mixture was stirred vigorously for 24 h. Precipitate formation was observed shortly after mixing the reactants (15 min). The white precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to yield colorless crystals, mp 120–122 °C. A total of 6.08 g of [K-18-crown-6][*o*-TolSiF₄] was isolated (88% yield). Anal. Calcd for C₁₉H₃₁SiF₄O₆K: C, 45.68; H, 6.28. Found: C, 45.34; H, 6.08. ^1H (CDCl₃): 2.47 (s, *o*-Me, 3 H); 3.45 (s, 18-crown-6, 24 H); 6.93–7.60 (m's, aromatic ring protons, 4 H). ^{19}F (CDCl₃, 21.0 °C): –116.0 (s) ^{19}F (CDCl₃, –48.2 °C): –116.0 (d_{Si}, J_{SiFav} = 216 Hz). ^{29}Si (CDCl₃, –58.2 °C): –121.9 (F, J_{SiFav} = 216 Hz).

(x) *Phenyltrifluorosilane, PhSiF₃*. The preparation of this organofluorosilane is well documented.^{38,39} The fluorination of PhSiCl₃ with CuF₂ in Et₂O was used: bp 102–105 °C (lit.³⁸ bp 101–102 °C). ^1H (CDCl₃): 7.34 (m, aromatic ring protons, 5 H). ^{19}F (CDCl₃): –141.4 (d_{Si}, J_{SiF} = 266 Hz). ^{29}Si (CDCl₃): –72.91 (q_F, J_{SiF} = 268 Hz).

(y) *Potassium 18-Crown-6 Phenyltetrafluorosilicate, [K-18-crown-6][PhSiF₄] (12)*. Phenyltrifluorosilane (1.90 g, 11.7 mmol), 18-crown-6 (3.10 g, 11.7 mmol), and KF (0.681 g, 11.7 mmol) were reacted in 30 mL of toluene at 25 °C. The reactants were stirred rapidly for 24 h. Precipitate formation was observed shortly after the reactants were combined. The white precipitate was collected by vacuum filtration. The compound was recrystallized from methylene chloride to afford colorless crystals, mp 164–166 °C dec. A total of 5.16 g (91% yield) of [K-18-crown-6][PhSiF₄] was isolated. Anal. Calcd for C₁₈H₂₉SiF₄O₆K: C, 44.60; H, 6.04. Found: C, 44.34; H, 5.99. ^1H (CDCl₃): 6.98–7.40 (m, aromatic ring protons, 5 H). ^{19}F (CDCl₃, –18.2 °C): –114.3 (d_{Si}, J_{SiFav} = 2.09 Hz). ^{29}Si (CDCl₃, –58.2 °C): –125.9 (p_F, J_{SiFav} = 210 Hz).

X-ray Studies. All X-ray crystallographic studies were done by using an Enraf-Nonius CAD4 diffractometer and graphite-monochromated molybdenum radiation ($\lambda(\text{K}\bar{\alpha}) = 0.71073 \text{ \AA}$) at an ambient temperature of 23 ± 2 °C. Details of the experimental procedures have been described previously.⁴⁰

Crystals were mounted in thin-walled glass capillaries that were sealed as a precaution against moisture sensitivity. Data were collected by using the θ -2 θ scan mode with $3^\circ \leq 2\theta_{\text{MoK}\bar{\alpha}} \leq 43^\circ$ for **1** and $3^\circ \leq 2\theta_{\text{MoK}\bar{\alpha}} \leq 50^\circ$ for **2**. The structures were solved by use of direct methods and difference Fourier techniques and were refined by full-matrix least-squares techniques.⁴¹ Nonhydrogen atoms were refined anisotropically. Hydrogen atom coordinates were obtained either by calculation in idealized positions or by regularizing coordinates obtained from difference Fourier syntheses. These atoms were included in the refinements as fixed isotropic scatterers. All computations were performed on a Microwax II computer using the Enraf-Nonius SDP system of programs.

X-ray Study for [(i-Bu)₃C₆H₅SiF₄]/[K-18-crown-6]-CH₂Cl₂ (1). Colorless crystals of **1** are large, distorted-square-prismatic needles with internal defects. The crystal used for the X-ray study was cut to dimensions of $0.35 \times 0.45 \times 0.48 \text{ mm}$.

Crystal data: [(C₄H₉)₃C₆H₅SiF₄][KO₆C₁₂H₂₄] \cdot CH₂Cl₂ (**1**) orthorhombic space group $P2_12_12$, [D₂⁴-No. 19].⁴² $a = 17.563$ (4) Å, $b = 22.528$ (5) Å, $c = 9.800$ (2) Å, $Z = 4$, and $\mu_{\text{MoK}\bar{\alpha}} = 3.585 \text{ cm}^{-1}$. A total of 2519 independent reflections (+ h , + k , + l) were measured. No corrections were made for absorption. The final agreement factors⁴³ (45 independent non-hydrogen atoms anisotropic, 55 independent hydrogen atoms fixed isotropic) were $R = 0.041$ and $R_{\text{w}} = 0.046$ for the 1875 reflections having $l \geq 2\sigma_l$.

X-ray Study for [Me₃C₆H₅SiF₄]/[K-18-crown-6] (2). The colorless crystal used for the X-ray study was cut from a polycrystalline mass of fused chunky crystals and was somewhat irregular with maximum dimensions of $0.40 \times 0.50 \times 0.68 \text{ mm}$.

Crystal data: [C₉H₁₁SiF₄][KO₆C₁₂H₂₄] (**2**) monoclinic space group $P2_1/m$ [C_{2h}-No. 11],⁴⁴ $a = 8.158$ (3) Å, $b = 14.847$ (7) Å, $c = 10.664$ (1) Å, $\beta = 91.24$ (2)°, $Z = 2$, and $\mu_{\text{MoK}\bar{\alpha}} = 3.075 \text{ cm}^{-1}$. A total of 2356 independent reflections (+ h , + k , $\pm l$) were measured. An empirical absorption correction

based on ψ scans was applied (from 0.956 to 1.000 on I). The fluorine atoms of the anion are disordered about a mirror plane to which the rest of the atoms conform. Attempts to remove the disorder by refining in the lower symmetry space group $P2_1$ led to the usual strong correlation between parameters of atoms related by the mirror plane and to a higher R factor. The final agreement factors⁴³ (22 independent non-hydrogen atoms anisotropic, four fluorine atoms in half-occupancy in general positions; 18 independent hydrogen atoms riding and isotropic) were $R = 0.040$ and $R_w = 0.059$ for the 1942 reflections having $l > 3\sigma_l$.

RESULTS

The atom-labeling scheme for **1** is given in the ORTEP plot of Figure 1. Atomic coordinates are given in Table I while selected bond lengths and angles are given in Table II. The corresponding information for **2** is given in Figure 2 and in Tables III and IV. Thermal parameters, additional bond lengths and angles, and hydrogen atom parameters for both compounds are provided as supplementary material.

DISCUSSION

Syntheses

All the pentacoordinated anions were prepared by reaction of the corresponding organotrifluorosilane with stoichiometric amounts of KF and 18-crown-6 in toluene solution, except $[(n\text{-PrSiF}_4)[\text{K-18-crown-6}]] \cdot \text{CH}_2\text{Cl}_2$. In accord with observations by Damrauer and Danahey,⁴⁵ the K-18-crown-6-salts that formed were relatively nonhy-




FIGURE 1 ORTEP plot of the cation-anion pair in $[(t\text{-Bu})_3\text{C}_6\text{H}_2\text{SiF}_4][\text{K-18-crown-6}]\cdot\text{CH}_2\text{Cl}_2$ (**1**) with thermal ellipsoids at the 30% probability level. Hydrogen atoms are omitted for clarity.

TABLE I
Atomic Coordinates in Crystalline
[(*t*-Bu)₃C₆H₂SiF₄][K-18-crown-6]·CH₂Cl₂ (1)^a

atom ^b	<i>x</i>	<i>y</i>	<i>z</i>	<i>B</i> _{equiv.} ^c Å ²
K	0.95658 (8)	0.93357 (6)	0.8147 (2)	4.11 (3)
C11	0.4146 (1)	0.3008 (1)	0.3193 (3)	9.51 (7)
C12	0.5158 (2)	0.2969 (1)	0.0901 (2)	8.23 (6)
Si	0.8401 (1)	0.83191 (7)	1.0241 (2)	3.76 (4)
F1	0.7601 (2)	0.8373 (2)	1.1129 (4)	5.41 (9)
F2	0.8893 (2)	0.8517 (2)	1.1555 (4)	6.1 (1)
F3	0.9215 (2)	0.8290 (1)	0.9313 (4)	4.57 (8)
F4	0.8188 (2)	0.8922 (1)	0.9407 (4)	5.27 (9)
O1	0.8830 (3)	1.0420 (2)	0.8666 (5)	6.3 (1)
O2	0.8675 (3)	0.9893 (2)	0.6084 (5)	6.4 (1)
O3	0.9899 (4)	0.9171 (2)	0.5207 (5)	7.5 (1)
O4	1.0768 (3)	0.8620 (2)	0.7241 (5)	6.4 (1)
O5	1.0917 (3)	0.9153 (2)	0.9834 (5)	5.9 (1)
O6	0.9729 (3)	0.9904 (2)	1.0703 (5)	6.2 (1)
C1	0.8214 (3)	0.7505 (2)	0.9681 (6)	2.9 (1)
C2	0.8500 (3)	0.7016 (3)	1.0445 (6)	3.4 (1)
C3	0.8509 (3)	0.6447 (3)	0.9862 (6)	3.6 (1)
C4	0.8229 (3)	0.6324 (2)	0.8568 (6)	3.7 (1)
C5	0.7887 (3)	0.6791 (3)	0.7894 (6)	3.7 (1)
C6	0.7858 (3)	0.7376 (2)	0.8412 (6)	3.1 (1)
C7	0.8846 (4)	0.7046 (3)	1.1909 (7)	4.7 (2)
C8	0.8940 (6)	0.6426 (4)	1.2551 (8)	8.4 (2)
C9	0.8337 (4)	0.7385 (3)	1.2917 (6)	5.6 (2)
C10	0.9660 (4)	0.7321 (4)	1.1825 (8)	6.5 (2)
C11	0.8308 (4)	0.5695 (3)	0.7972 (7)	5.5 (2)
C12	0.7836 (6)	0.5275 (4)	0.881 (1)	11.5 (3)
C13	0.9120 (5)	0.5506 (4)	0.790 (1)	11.8 (3)
C14	0.7998 (8)	0.5668 (4)	0.653 (1)	13.0 (4)
C15	0.7413 (3)	0.7818 (3)	0.7494 (6)	3.8 (1)
C16	0.6944 (4)	0.7513 (3)	0.6369 (8)	6.8 (2)
C17	0.7984 (4)	0.8221 (3)	0.6727 (7)	5.3 (2)
C18	0.6837 (4)	0.8190 (3)	0.8311 (8)	5.6 (2)
C20	0.8169 (5)	1.0469 (4)	0.781 (1)	8.4 (3)
C21	0.8383 (6)	1.0454 (4)	0.636 (1)	8.9 (3)
C22	0.8919 (5)	0.9844 (4)	0.4697 (8)	8.0 (2)
C23	0.9212 (6)	0.9242 (4)	0.4453 (8)	8.8 (2)
C24	1.0223 (7)	0.8597 (4)	0.5047 (8)	10.0 (3)
C25	1.0948 (6)	0.8566 (4)	0.582 (1)	10.0 (3)
C26	1.1432 (4)	0.8548 (4)	0.807 (1)	8.5 (3)
C27	1.1192 (4)	0.8568 (3)	0.953 (1)	7.5 (2)
C28	1.0670 (5)	0.9212 (4)	1.1223 (7)	7.5 (2)
C29	1.0406 (5)	0.9836 (4)	1.1417 (8)	7.4 (2)
C30	0.9443 (6)	1.0488 (3)	1.0810 (8)	8.0 (2)
C31	0.8699 (6)	1.0510 (4)	1.006 (1)	9.1 (3)
C100	0.5048 (5)	0.2805 (3)	0.2633 (8)	6.0 (2)

^a Numbers in parentheses are estimated standard deviations.

^b Atoms are labeled to agree with Figure 1. ^c Equivalent isotropic thermal parameters are calculated as $(4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$.

grosscopic compared to tetraalkylammonium salts that had been reported previously for related anionic fluorosilicates.⁶ We observed a similar enhancement in hydrolytic stability in the case of five-coordinated cyclic pentoxy silicates.⁴⁶

The formation of [*n*-PrSiF₄][K-18-crown-6] by a ring-cleavage reaction represents

TABLE II
Selected Distances (\AA) and Angles (deg) for
[(*t*-Bu)₃C₆H₅SiF₄][K-18-crown-6]·CH₂Cl₂ (1)^a

Distances			
Si-F1	1.657 (4)	K-O1	2.809 (5)
Si-F2	1.613 (4)	K-O2	2.848 (5)
Si-F3	1.696 (4)	K-O3	2.963 (5)
Si-F4	1.630 (4)	K-O4	2.801 (5)
Si-C1	1.942 (6)	K-O5	2.921 (5)
K-F3	2.690 (4)	K-O6	2.828 (5)
K-F4	2.872 (4)		

Angles			
F1-Si-F2	90.9 (2)	F2-Si-F4	107.1 (2)
F1-Si-F3	177.9 (2)	F2-Si-C1	125.2 (2)
F1-Si-F4	90.4 (2)	F3-Si-F4	87.5 (2)
F1-Si-C1	94.3 (2)	F3-Si-C1	87.4 (2)
F2-Si-F3	89.2 (2)	F4-Si-C1	127.3 (2)

^aEstimated standard deviations in parentheses. The atom-labeling scheme is shown in Figure 1.




FIGURE 2 ORTEP plot of $[\text{Me}_3\text{C}_6\text{H}_5\text{SiF}_4][\text{K-18-crown-6}]$ (2) with thermal ellipsoids at the 30% probability level. Atoms with the same labels are related by a crystallographic mirror plane ($x, \frac{1}{2} - y, z$). For purposes of clarity, only one set of positions is shown for the fluorine atoms and hydrogen atoms are omitted.

a potential new route for the preparation of such anions.⁴⁷ The process is indicated in eq 1 although the




TABLE III
Atomic Coordinates in Crystalline
[Me₃C₆H₅SiF₄][K-18-crown-6] (2)^a

atom ^b	x	y	z	B _{equiv} , Å ²
K	0.84529 (8)	0.250	0.14590 (5)	4.40 (1)
Si	0.34765 (9)	0.250	0.18506 (7)	4.64 (2)
F1 ^c	0.1644 (3)	0.1905 (2)	0.1868 (2)	6.97 (7)
F2 ^c	0.4136 (3)	0.1591 (2)	0.1040 (2)	6.67 (6)
F3 ^c	0.5279 (3)	0.2922 (2)	0.1722 (2)	7.21 (6)
F4 ^c	0.2610 (4)	0.3226 (2)	0.1089 (3)	8.53 (8)
O1	0.8845 (3)	0.250	0.4043 (2)	6.93 (6)
O2	0.8216 (2)	0.0855 (1)	0.2812 (2)	6.66 (4)
O3	0.8168 (2)	0.0856 (1)	0.0125 (2)	6.60 (4)
O4	0.7311 (3)	0.250	-0.1018 (2)	6.61 (6)
C1	0.8395 (3)	0.1707 (3)	0.4675 (2)	8.95 (8)
C2	0.9051 (3)	0.0927 (2)	0.3978 (2)	9.08 (7)
C3	0.8858 (3)	0.0158 (2)	0.2082 (3)	8.69 (8)
C4	0.7832 (3)	0.0098 (2)	0.0910 (3)	9.01 (8)
C5	0.7082 (3)	0.0925 (2)	-0.0910 (3)	8.87 (6)
C6	0.7586 (3)	0.1703 (3)	-0.1709 (2)	9.28 (9)
C11	0.3595 (3)	0.250	0.3634 (2)	3.87 (5)
C12	0.3604 (2)	0.1689 (1)	0.4318 (2)	4.11 (4)
C13	0.3489 (2)	0.1703 (1)	0.5628 (2)	4.62 (4)
C14	0.3381 (3)	0.250	0.6293 (3)	4.68 (6)
C15	0.3762 (3)	0.0782 (2)	0.3686 (2)	5.52 (5)
C16	0.3143 (4)	0.250	0.7693 (3)	6.44 (8)

^a Numbers in parentheses are estimated standard deviations. Values with no esd's were fixed. ^b Atoms are labeled to agree with Figure 2.

^c Equivalent isotropic thermal parameters are calculated as $(4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$.

^d Half-occupancy.


TABLE IV
Selected Distances (Å) and Angles (deg) for
[Me₃C₆H₅SiF₄][K 18-crown-6] (2)^a

Distances			
Si-F1	1.737 (3)	K-O3	2.832 (2)
Si-F2	1.696 (3)	K-O4	2.782 (2)
Si-F3	1.607 (3)	K-O2	2.845 (2)
Si-F4	1.516 (3)	K-F1''	2.775 (2)
Si-C11	1.902 (3)	K-F3	2.684 (2)
K-O1	2.767 (2)		

Angles			
F1-Si-F2	83.3 (1)	F2-Si-F4	116.2 (1)
F1-Si-F3	171.3 (1)	F2-Si-C11	120.0 (1)
F1-Si-F4	88.6 (2)	F3-Si-F4	95.4 (2)
F1-Si-C11	90.8 (1)	F3-Si-C11	93.4 (1)
F2-Si-F3	88.0 (1)	F4-Si-C11	123.3 (1)

^a Estimated standard deviations in parentheses. The atom-labeling scheme is shown in Figure 2. Double prime denotes the symmetry operation $1 + x, y, z$.

presence of C₆H₅CH₂⁻ was not verified. Scheme I represents a likely mechanism for the formation of *n*-PrSiF₄⁻. A two-step process is suggested, with the first step involving nucleophilic attack by fluoride ion resulting in an opening of the standard four-membered ring with concomitant proton capture from the solvent. The or-



SCHEME I

ganotrifluorosilane formed then undergoes nucleophilic attack by fluoride ion to produce the five-coordinated anion. Ring cleavage of the highly strained 1,1-disubstituted silacyclobutanes has been shown to be facile in the presence of Lewis acids.²⁹ Apparently, ring strain is too severe in the disubstituted anion, $(\text{CH}_2)_3\text{SiF}_3^-$, making the ring opening process leading to the monosubstituted anion the favored reaction pathway.

Ligand Exchange

Both the variable-temperature ^{19}F and ^{29}Si NMR solution-state spectra of $[(\text{TTBP})\text{SiF}_4^-][\text{K-18-crown-6}]$ are representative of a system in which rapid intramolecular fluorine exchange is sufficiently slowed at reduced temperatures, with the resulting spectrum reflecting the ground-state structure. The spectra are displayed in Figures 3 and 4, respectively, and represent the first example for which ligand exchange has been “stopped” for a RSiF_4^- derivative.

At 314 K, a single fluorine signal, centered at -95.08 ppm, is observed (Figure 3), indicating rapid exchange of all four fluorine nuclei. As the temperature decreases, the fluorine signal broadens and two different fluorine signals centered at -77.98 and -112.7 ppm appear. The signals sharpen and ^{29}Si – ^{19}F spin–spin coupling is observed near the low-temperature limit. At the low-temperature limit (205 K), both fluorine resonances exhibit fine structure arising from ^{19}F – ^{19}F spin–spin coupling. This results in a triplet splitting of each resonance (Figure 5).

The same temperature-dependent spectral behavior is observed when the ^{29}Si NMR of the anion is examined (Figure 4). At 254 K, the silicon resonance is split into a pentuplet, indicating rapid exchange of all four fluorine nuclei. As the temperature decreases, the pentuplet splitting of the ^{29}Si signal decomposes into a new multiplet. At the low-temperature limit studied, the silicon signal consists of a triplet of triplets, indicative of stopped exchange. Retention of ^{29}Si – ^{19}F spin–spin coupling throughout the temperature range studied adds further evidence to indicate the exchange process is, indeed, an intramolecular process. The exchange




FIGURE 3 Variable-temperature ^{19}F NMR spectra of $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ in CD_2Cl_2 from the region of "stopped" exchange, -68°C , to the fast exchange region, 41°C . The spectra are referenced to CFCl_3 (internal standard).




FIGURE 4 Variable-temperature ^{29}Si NMR spectra of $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ showing retention of $^{29}\text{Si}-^{19}\text{F}$ spin-spin coupling from the region of "stopped" exchange, -107°C , to the fast exchange region at -19°C .




FIGURE 5 ^{19}F NMR spectra of $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ in CD_2Cl_2 at -73°C . $\delta(^{19}\text{F}_{\text{ax}}) = -77.98$ ppm; $J_{\text{SiF}} = 248$ Hz; $\delta(^{19}\text{F}_{\text{eq}}) = -112.7$ ppm; $J_{\text{SiF}} = 191$ Hz; $J_{\text{FeF}_4} = 29.6$ Hz.

behavior exhibited by the fluorosilicate anion is analogous to its isoelectronic phosphorus derivative, $(\text{TTBP})\text{PF}_4$.⁴⁸

Both the ^{19}F and ^{29}Si NMR spectra of $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ are consistent with a trigonal-bipyramidal geometry for the anion in solution with the TTBP substituent occupying an equatorial position. The X-ray structure, likewise is trigonal bipyramidal. With the knowledge that both phosphoranes^{13,15,16} and five-coordinated anionic silicates^{13,17-21} form a series of structures extending along the Berry coordinate¹² from the trigonal bipyramidal to the square pyramid, there is little doubt that the solution-state intramolecular exchange process for anionic silicates, like phosphoranes, is represented by this coordinate.

The energy barrier for ligand exchange of **1** is compared with values available for monosubstituted phosphoranes, RPF_4 ,^{49,50} in Table V. MeSPF_4 also shows a low-temperature NMR pattern consistent with stopped exchange.^{51,52} Rotational hindrance contributes to the slowing of the exchange process for this substance as well as that for the R_2NPF_4 derivatives^{49,50} while steric effects undoubtedly reduce exchange rates for **1** and the related phosphorane, $(\text{TTBP})\text{PF}_4$.⁴⁸ The magnitude

TABLE V
Energy Barriers for Intramolecular Exchange in
 $[(\text{TTBP})\text{SiF}_4][\text{K-18-crown-6}]$ (**1**) and RPF_4 Phosphoranes

compound	ΔG^\ddagger , ^a kcal/mol	ref
CIPF_4	4.2	49
Me_2NPF_4	8.8	49
$i\text{-Pr}_2\text{NPF}_4$	7.5	50
$(\text{TTBP})\text{SiF}_4^-$	12.8*	this work

* E_a value.

of the steric effect required to stop exchange for silicates apparently needs to be large. Even the mesityl derivative, **2**, showed no evidence for significant slowing of fluorine exchange down to -100°C .

Further insight into the specific steric effects that may be encountered in solution is gained upon examination of the molecular structure of **1** discussed in the next section.

Structural Details

The molecular geometry about the silicon atom in both **1** and **2** can be referred to a trigonal bipyramidal where the aryl substituent is, as expected, in an equatorial position.

For **1**, the potassium ion is in close proximity with both an axial and an equatorial fluorine atom ($\text{K}-\text{F3}_{\text{ax}} = 2.690$ (4) Å and $\text{K}-\text{F4}_{\text{eq}} = 2.872$ (4) Å, compared to the $\text{K}-\text{F}$ distance in potassium fluoride⁵³ of 2.67 Å). While the $\text{Si}-\text{F}_{\text{ax}}$ bond lengths are longer than the $\text{Si}-\text{F}_{\text{eq}}$ bond lengths, as expected, the interaction of the potassium ion with F3 and F4 causes the $\text{Si}-\text{F}$ bonds to these atoms to be the longest of their kinds: $\text{Si}-\text{F3}_{\text{ax}} = 1.696$ (4) Å compared to $\text{Si}-\text{F1}_{\text{ax}} = 1.657$ (4) Å and $\text{Si}-\text{F4}_{\text{eq}} = 1.630$ (4) Å compared to $\text{Si}-\text{F2}_{\text{eq}} = 1.613$ (4) Å.

In **2**, the potassium atom is not associated with one axial and one equatorial fluorine atom in the same atom, as was the case for **1**. Rather, it is positioned between the axial F3 and the axial F1 of a translationally related anion (atom" = $1 + x, y, z$), forming a cation-anion chain along *a*. The distances $\text{K}-\text{F3}$ and $\text{K}-\text{F1}''$ are 2.684 (2) and 2.775 (2) Å, respectively.

The most surprising structural features in **1** reflect the extreme crowding in the molecule. The $\text{F}_{\text{eq}}-\text{Si}-\text{F}_{\text{eq}}$ angle of 107.1 (2) $^{\circ}$ is very small. The phenyl group is *not* rotated out of the equatorial plane, but is tipped up out of it in such a way that there is a local mirror plane containing Si , F1 , F3 , C1 , and C4 . The dihedral angle between the equatorial plane ($\text{Si}, \text{F2}, \text{F4}$, and C1 coplanar to within ± 0.030 Å) and the plane of the phenyl group is 13.8 $^{\circ}$. However, the torsion angles $\text{F2}-\text{Si}-\text{C1}-\text{C2}$ and $\text{F4}-\text{Si}-\text{C1}-\text{C6}$ have values of 1.3 and 0.5 $^{\circ}$, respectively. These torsion angles would be expected to approach the dihedral angle if rotation of the phenyl group about the $\text{Si}-\text{C1}$ bond had occurred. Additional data which reflect the fact that the phenyl group is tipped out of the equatorial plane are the dispositions of C4 and C11 , which would be in this plane if no tipping had occurred. In fact they are displaced from this plane by 0.685 and 1.150 Å, respectively, in a direction toward F3 . It is interesting to note that the phenyl group is tipped up toward the longest of the $\text{Si}-\text{F}$ axial bonds.

The driving force for this deformation is apparent in the contacts between the axial F atoms and the methyl groups of the *o-t*-Bu groups. F3 is in van der Waals contact with C10 and C17 ($\text{F3}-\text{C10} = 3.381$ (9) Å and $\text{F3}-\text{C17} = 3.358$ Å compared to the van der Waals sum of 3.35 Å). F1 , however, is in even closer proximity to C9 and C18 ($\text{F1}-\text{C9} = 3.114$ (8) Å and $\text{F1}-\text{C18} = 3.098$ Å). In the absence of the aforementioned deformation, these contacts would be even shorter. This arrangement in the molecule also serves to stagger the methyl groups relative to the equatorial fluorine atoms: $\text{F2}-\text{C9} = 3.039$ (8) Å and $\text{F2}-\text{C10} = 3.023$ (9) Å; $\text{F4}-\text{C17} = 3.085$ (8) Å and $\text{F4}-\text{C18} = 3.083$ (8) Å. It is worth noting that these close F-Me contacts occur in the presence of the extremely small $\text{F2}-\text{Si}-\text{F4}$ angle and

serves to hinder an opening up of this angle which would occur during Berry pseudorotation.¹² For the latter process to occur, a 90° rotation of the phenyl group must accompany a concurrent interchange of the equatorial and axial fluorine atoms, brought about by a synchronous bond bending of the respective F–Si–F angles. Thus, the “stoppage” of intramolecular fluorine atom exchange postulated to take place by the pseudorotation route in interpreting the dynamic NMR behavior seems reasonable. Assuming that the molecular structure observed for **1** is maintained in solution, tilting of the phenyl group during the course of its rotation toward the newly lengthened Si–F_{ax} bond is required as it assumes its final position in the pseudorotated conformation.

In **2**, there is a crystallographic mirror plane that passes through Si, C11, C14, and C16 of the anion and through O1, O4, and K⁺ of the cation. The fluorine atoms of the anion are disordered about the mirror plane. It is likely that the disorder in the fluorine atoms has had a deleterious effect on the determination of the Si–F bond lengths. This is reflected in the usually short Si–F4_{eq} distance of 1.516 (3) Å.

The F_{eq}–Si–F_{eq} angle in **2** (116.2 (1)°) is only slightly smaller than the idealized value of 120° and reflects less crowding in the anion than for **1**. In **2**, there is a rotation of the phenyl group about an axis coincident with the Si–C11 bond of about 25°, as evidenced by the F2–Si–C11–C12 torsion angle of –24.8°. The atoms of the equatorial plane (Si, F₂, F₄, and C11) are coplanar to within ± 0.035 Å. The dihedral angle between this plane and the plane of the phenyl group is 28.4° and reflects the rotation of the phenyl group in large measure. However, the phenyl group in this case is also tipped up out of the equatorial plane towards the longer Si–F1 bond but to a lesser extent than in **1**. This can be seen in the distances of C14 and C16 from the equatorial plane, which are 0.399 and 0.692 Å, respectively. The disposition of the mesityl group serves to stagger the *o*-methyl groups between axial–equatorial pairs of fluorine atoms: C15–F1 = 3.062 (3) Å and C15–F2 = 3.088 (3) Å; C15’–F3 = 3.120 (3) Å and C15’–F4 = 3.257 (4) Å.

In Table VI, selected structural parameters for **1** and **2** are compared with those

TABLE VI
Comparison of Bond Lengths and Angles for
Trigonal-Bipyramidal Organotetrafluorosilicate Anions

	[PhSiF ₄] ^{–a}	[MesSiF ₄] [–]	[(TTBP)- SiF ₄] [–]
Si–F _{ax} (av), Å	1.669 (3)	1.672 (3)	1.677 (4)
Si–F _{eq} (av), Å	1.601 (3)	1.606 (3)	1.622 (4)
Si–C, Å	1.871 (4)	1.902 (3)	1.942 (6)
F _{ax} –Si–F _{ax} , ^b deg	174.6 (1)	171.3 (1)	177.9 (2)
F _{eq} –Si–F _{eq} , ^b deg	121.4 (2)	116.2 (1)	107.1 (2)
displacement of Si atom out of plane of aromatic ring, Å	0.029	0.251	0.454
τ , ^c deg	54.3	28.4	
Δ , ^d deg	6.8	12.5	15.0

^a Reference 9. ^b These values are for angles that are on the side of the axial atoms opposite the pivotal equatorial phenyl group.

^c Rotation of aromatic ring out of the equatorial plane formed by the C atom bonded to silicon, Si, and the two F_{eq} atoms. ^d Δ represents the sum of the displacements of the bond angles, F_{ax}–Si–F_{ax} and F_{eq}–Si–F_{eq}, from 180 and 120°, respectively, from the ideal TBP structure.

for the related, but less crowded, $[\text{PhSiF}_4]^-$ anion.⁹ As the degree of crowding increases in the series from the phenyl, through the mesityl, to the tri-*tert*-butyl-phenyl anionic silicates, there is a steady increase in the Si-C bond lengths, a less pronounced increase in the average Si-F bond lengths, and a decrease in the F_{eq} -Si-F_{eq} angle, which is very pronounced for the tri-*tert*-butyl compound **1**. Perhaps a good measure of the steric effect is found in the sum of the axial and equatorial deviations from 180 and 120° angles, respectively, that are present in the ideal TBP structure. These are the Δ values listed in Table VI. They increase in the order $\text{PhSiF}_4^- < \text{MesSiF}_4^- < (\text{TTBP})\text{SiF}_4^-$. Previously, the largest value for this parameter was obtained for the $\text{Xyl}_2\text{SiF}_3^-$ derivative, 11.8°.^{54,55}

There is also a correlation between the degree to which the phenyl group is tipped up and the degree of crowding that can be measured by looking at the displacement of the silicon atom out of the plane of the aromatic ring. In the phenyl derivative, the silicon atom is essentially coplanar with the ring with a displacement of 0.029 Å. This value progressively increases for the more sterically crowded molecules in the order **1** > **2**. Finally, as steric crowding increases, Ph to Mes to TTBP, the rotation (τ) of the phenyl ring out of the equatorial plane decreases.

NMR Parameters

¹⁹F and ²⁹Si NMR data for organotrifluorosilanes, RSiF_3 , and organotetrafluorosilicates, RSiF_4^- , are summarized in Tables VII and VIII, respectively. With a wide range of alkyl and aromatic substituents, rather narrow ranges for ¹⁹F and ²⁹Si chemical shifts, as well as Si-F coupling constants, result in each of the series studied. However as amplified in Table IX, the ¹⁹F chemical shifts for the pentacoordinated derivatives, RSiF_4^- , are on average 24 ppm downfield from those for the respective silanes, RSiF_3 , while ²⁹Si chemical shifts are upfield by about 51 ppm on average for the pentacoordinated members compared to the silanes. Also,

TABLE VII
¹⁹F and ²⁹Si Chemical Shifts and Coupling Constants of
Organofluorosilanes, RSiF_3 ^a

R	$\delta(^{19}\text{F})$, ppm	J_{SiF} , Hz	$\delta(^{29}\text{Si})$, ppm	J_{SiF} , Hz
Me ^b	-139.2	260		
F ₃ SiCH ₂ CH ₃	-139.7	282	-61.11	281
<i>n</i> -Bu ^b	-140.4	292		
<i>t</i> -Bu	-149.8	305	-62.10	304
C ₆ H ₁₁	-144.2	296	-62.6	296
Bz	-138.9	281	-64.13	282
Ph	-143.0	266	-72.91	268
<i>p</i> -Tol	-141.4	266	-74.00	267
<i>m</i> -Tol	-140.2	268	-72.59	268
<i>o</i> -Tol	-138.9	270	-71.14	270
<i>p</i> -ClPh	-140.8	252	-73.12	251
Mes	-130.6	274	-69.19	274
TTBP	-124.4	266	-61.3	262

^aAll the NMR data were obtained in this study unless otherwise noted. Data from this study were recorded at room temperature in CDCl_3 solvent. ^b¹⁹F NMR data recorded on neat liquids were taken from ref 6.

TABLE VIII
 ^{19}F and ^{29}Si Chemical Shifts and Coupling Constants of
 Organotetrafluorosilicates, $[\text{RSiF}_4]^-$ ^a

R	t, °C	$\delta^{(29)\text{Si}}$, ppm		
		$\delta^{(19)\text{F}}$, ppm	J_{SiF} , Hz	t, °C
Me ^c	-60	-110.9	218	
n-Pr	-118.1	224		-78.2
F_3SiCH_2	-88.2	-118.2	227	-98.2
CH ₃				-109.9
n-Bu ^c	-60	-116.6	223	
t-Bu	-27.2	-123.5	235	-58.2
C ₆ H ₁₁	-58.3	-123.8	230	-68.7
Bz	-98.2	-117.3	215	-98.2
Ph ^d	35	-119.2	206	
Ph	-18.2	-114.3	209	-58.2
Ph ^e	25	-116.8	205	
p-Tol		-122.4	210	-62.3
m-Tol	-58.2	-119.6	214	-58.2
o-Tol	-48.2	-116.0	216	-58.2
p-CIPh	-38.2	-120.9	212	-58.3
Mes	21.0	-111.8	219	21.0
TTBP ^f	-87.8	-95.34	220	-107.2
				-114.1
				221

^a [K-18-crown-6]⁺ salt. CDCl_3 solvent except for R = Bz and TTBP where CD_2Cl_2 was used. All the NMR data were obtained in this study unless otherwise noted. ^b Average values. ^c $[\text{n-Pr}_4\text{N}]^+$ salt, CH_2Cl_2 solvent. ^d $[\text{n-Pr}_4\text{N}]^+$ salt, CHCl_3 solvent. ^e $[\text{Me}_2\text{N}]^+$ salt, Me_2SO solvent. ^f At low temperature where exchange is stopped. Average values are listed however.

TABLE IX
 ^{19}F and ^{29}Si Chemical Shift Differences $|\delta(\text{RSiF}_4^-) - \delta(\text{RSiF}_3)|$ ^a
 and Si-F Coupling Constant Differences $J_{\text{RSiF}_3} - J_{\text{RSiF}_4}$ ^b

R	$\Delta\delta^{(19)\text{F}}$, ppm	$\Delta\delta^{(29)\text{Si}}$, ppm	ΔJ_{SiF} , Hz
Me	28.3		42
$\text{F}_3\text{SiCH}_2\text{CH}_3$	21.5	47.9	55
n-Bu	23.8		69
t-Bu	26.3	52.7	69
C ₆ H ₁₁	20.4	52.1	65
Bz	21.6	53.6	65
Ph	28.7	53.0	58
p-Tol	19.0	52.1	57
m-Tol	20.6	51.7	54
o-Tol	22.9	50.8	54
p-CIPh	19.6	52.1	39
Mes	18.8	50.9	55
TTBP	29.1	52.8	44

^a The difference in chemical shifts is an absolute value. The ^{19}F chemical shift moves downfield upon pentacoordination while the ^{29}Si chemical shift moves upfield upon pentacoordination. ^b The difference in coupling constants is a positive value. ^c Average values.

it is seen that the Si-F coupling constants are larger for the silanes compared to the anionic siliconates.

While the ranges are narrow, there is a discernible trend associated with the substituent effect on ^{29}Si chemical shifts of essentially σ -donor ligands compared to those that also are capable of π bonding. For both the tetracoordinated and

pentacoordinated silicon compounds, the aryl substituents, except for the mesityl and tri-*tert*-butylphenyl groups, cause about a 10 ppm upfield shift relative to the alkyl substituents. The latter difference follows Oldfield's⁵⁶ correlation for ligand types attached to silicon and ²⁹Si chemical shifts for mineral silicates and aluminosilicates. No such differences are seen in the ¹⁹F chemical shifts in the two series. This may be reasonable since changes in the ¹⁹F chemical shifts are regarded as a secondary effect of substitution. However, as with the ²⁹Si chemical shifts, there is a drop off in the ¹⁹F chemical shifts to more deshielded values for the most sterically encumbered substituents, mesityl and 2,4,6-tri-*tert*-butylphenyl. This general feature must be a manifestation of the steric effects introduced in the respective molecules in each series. These chemical shift changes are greater for the TTBP group than for the mesityl group. This implies a greater steric effect, attributable to the TTBP, that is consistent with conclusions based on the ability of this group to stop intramolecular ligand exchange compared to the lack of this ability for the mesityl group and that causes larger distortions in the molecular structures of the TTBP derivative compared to the mesitylsilicate.

ACKNOWLEDGEMENT

The support of this research by the National Science Foundation (Grant CHE85-04737) is gratefully acknowledged.

Registry No. **1**, 121351-71-1; **2**, 121351-72-2; **3**, 121351-74-4; **4**, 121351-76-6; **5**, 121351-78-8; **6**, 121351-80-2; **7**, 121351-82-4; **8**, 121351-84-6; **9**, 121351-86-8; **10**, 121351-88-0; **11**, 121351-90-4; **12**, 102307-91-5; (CH₃)₂SiF₅, 13883-11-9; F₃SiCH₂CH₂SiF₃, 121330-06-1; C₆H₁₁SiF₃, 368-46-7; MesSiCl₃, 17902-75-9; MesSiF₃, 121330-07-2; Cl₃SiCH₂CH₂SiCl₃, 2504-64-5; C₆H₁₁SiCl₃, 98-12-4; tri-*tert*-butyl-phenyltrifluorosilane, 99789-89-6; 1,1-dichlorosilacyclobutane, 2351-33-9; *tert*-butyltrifluorosilane, 60556-38-9; benzyltrifluorosilane, 658-32-2; (*p*-chlorophenyl)(trifluorosilane, 349-92-8; *p*-tolyltrifluorosilane, 13688-78-3; *m*-tolyltrifluorosilane, 13688-77-2; *o*-tolyltrifluorosilane, 363-86-0; phenyltrifluorosilane, 368-47-8.

Supplementary Material Available: Thermal parameters, additional bond lengths and angles, and hydrogen atom parameters for **1** (Tables S1–S3) and for **2** (Tables S4–S6) (12 pages); tables of calculated and observed structure factors (16 pages). Ordering information is given on any current masthead page.

REFERENCES AND NOTES

- (1) (a) Pentacoordinated Molecules. 77. (b) Part 76: Johnson, S. E., Deiters, J. A.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1989**, *111*, 3250.
- (2) Presented in part at the Third Chemical Congress of North America, Toronto, Canada, June 1988, paper INOR 31.
- (3) This work represents a portion of the Ph.D. Thesis of S. E. Johnson, University of Massachusetts, Amherst, MA, 1990.
- (4) (a) Corriu, R. J. P.; Guerin, C.; Henner, B. J. L.; Wong Chi Man, W. W. C. *Organometallics* **1988**, *7*, 237. (b) Kira, M.; Kobayashi, M.; Sakurai, H. *Tetrahedron Lett.* **1987**, *35*, 4081.
- (5) Deiters, J. A.; Holmes, R. R. *Abstracts of Papers*, 19th National Meeting of the American Chemical Society, Dallas, TX: American Chemical Society: Washington, DC, 1989; INOR423. See also: Deiters, J. A.; Holmes, R. R.; Holmes, J. M. *J. Am. Chem. Soc.* **1988**, *110*, 7672.
- (6) Klanberg, F.; Muetterties, E. L. *Inorg. Chem.* **1968**, *7*, 155.
- (7) Kuroda, K.; Ishikawa, N. *Nippon Kagaku Zasshi* **1977**, *41*, 77.
- (8) Ault, B. S.; Tandoc, U. *Inorg. Chem.* **1981**, *20*, 1937.
- (9) Schomburg, D. J. *Organomet. Chem.* **1981**, *221*, 137.
- (10) Marat, R. K.; Janzen, A. F. *Can. J. Chem.* **1977**, *55*, 3845.
- (11) Gibson, J. A.; Ibott, D. G.; Janzen, A. F. *Can. J. Chem.* **1973**, *51*, 3203.
- (12) Berry, R. S. *J. Chem. Phys.* **1960**, *32*, 933.

- (13) Holmes, R. R. *Prog. Inorg. Chem.* **1984**, *32*, 119 and references cited therein.
- (14) (a) Holmes, R. R. *Pentacoordinated Phosphorus: Spectroscopy and Structure*; ACS Monograph 175; American Chemical Society: Washington, DC, 1980; Chapter 3. (b) Holmes, R. R., *Penta-coordinated Phosphorus: Reaction Mechanisms*; ACS Monograph 176; American Chemical Society: Washington, DC, 1980; Chapter 1.
- (15) Cited in ref 14a, Chapter 2.
- (16) Holmes, R. R.; Deiters, J. A. *J. Am. Chem. Soc.* **1977**, *99*, 3318.
- (17) Holmes, R. R.; Day, R. O.; Chandrasekhar, V.; Holmes, J. M. *Inorg. Chem.* **1985**, *24*, 2009.
- (18) Holmes, R. R.; Day, R. O.; Chandrasekhar, V.; Harland, J. J.; Holmes, J. M. *Inorg. Chem.* **1985**, *24*, 2016.
- (19) Holmes, R. R.; Day, R. O.; Harland, J. J.; Holmes, J. M. *Organometallics* **1984**, *3*, 347.
- (20) Holmes, R. R.; Day, R. O.; Harland, J. J.; Sau, A. C.; Holmes, J. M. *Organometallics* **1984**, *3*, 341.
- (21) Harland, J. J.; Day, R. O.; Vollano, J. F.; Sau, A. C.; Holmes, R. R. *J. Am. Chem. Soc.* **1981**, *103*, 5269.
- (22) Blinka, T.; Helmer, B.; West, R. *Adv. Organomet. Chem.* **1984**, *23*, 193.
- (23) Bushweller, H. C.; Letendre, L. J.; Brunelle, J. A.; Bilofsky, H. S.; Whalon, M. R.; Fleischman, S. H. *QCPE* **1978**, Program 466. See also Binsch, G. *J. Magn. Reson.* **1978**, *30*, 625. DNMR-4 is a modification of DNMR-3; Kleier, D. A.; Binsch, G. *QCPE* **1969**, Program 165.
- (24) Ragle, J. R. Personal communication.
- (25) Nakadaira, Y.; Kazuya, O.; Sakurai, H. *J. Organomet. Chem.* **1985**, *309*, 247.
- (26) Weidenbruch, M.; Kramer, K. *J. Organomet. Chem.* **1985**, *291*, 159.
- (27) Pearson, D. E.; Frazer, M. G.; Frazer, V. S.; Washburn, L. C. *Synthesis* **1976**, 621.
- (28) Fink, M. J.; Michalczyk, M. J.; Haller, K. J.; West, R.; Michl, J. *Organometallics* **1984**, *3*, 793.
- (29) (a) Auner, N.; Grobe, J. *Z. Anorg. Allg. Chem.* **1982**, *485*, 53. (b) Dabdit, O.; Sutcliffe, L. H. *Org. Magn. Reson.* **1983**, *21*, 412.
- (30) Anderson, H. H. *J. Am. Chem. Soc.* **1959**, *81*, 4785.
- (31) Voronkov, M. G.; Iskorik, Yu. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1964**, *7*, 1215.
- (32) Vcelák, J.; Chvalovsky, V. *J. Organomet. Chem.* **1970**, *23*, 47.
- (33) Sternbach, B.; MacDiarmid, A. J. *J. Am. Chem. Soc.* **1961**, *83*, 3384.
- (34) Ernst, C. R.; Spialter, L.; Buel, G. R.; Wilhit, D. L. *J. Am. Chem. Soc.* **1974**, *96*, 5375.
- (35) Ishikawa, N.; Kurodo, K. *Nippon Kagaku Zasshi* **1968**, *89*, 421.
- (36) Hradil, J.; Chvalovsky, V. *Collect. Czech. Chem. Commun.* **1967**, *32*, 171.
- (37) Frost, L. W. *J. Am. Chem. Soc.* **1956**, *78*, 3855.
- (38) Emeleus, H. S.; Wilkins, C. S. *J. Chem. Soc.* **1944**, 454.
- (39) Helmer, B. J.; West, R.; Corriu, R. J. P.; Poirer, M.; Royo, G.; DeSaxce, A. *J. Organomet. Chem.* **1983**, *251*, 295.
- (40) Sau, A. C.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1981**, *20*, 3076.
- (41) The function minimized was $\Sigma w(|F_o| - |F_c|)^2$, where $w^{1/2} = 2F_0L_p/\sigma_f$.
- (42) *International Tables for X-ray Crystallography*; Kynoch: Birmingham, England, 1969; Vol. I, p. 105.
- (43) $R = \Sigma |F_o| - |F_c|/\Sigma |F_o|$ and $R_w = \langle \Sigma w(|F_o| - |F_c|)^2/\Sigma w|F_o|^2 \rangle^{1/2}$. For **1**, these values are for the configuration having the lowest R_w .
- (44) Reference 42, p. 93.
- (45) Damrauer, R.; Danahey, S. E. *Organometallics* **1986**, *5*, 1490.
- (46) Holmes, R. R.; Day, R. O.; Payne, J. S. *Phosphorus, Sulfur, Silicon* **1989**, *42*, 1.
- (47) Related Si-C bond cleavage reactions induced by steric effects occur on heating K-18-crown-6⁺ salts of $\text{Mes}_2\text{SiF}_5^-$ and $\text{Xyl}_2\text{SiF}_5^-$. See respectively ref 1b and: Damrauer, R.; O'Connell, B.; Danahey, S. E.; Simon, R. *Organometallics* **1989**, *8*, 1167.
- (48) Appel, R.; Krieger, L. *J. Fluorine Chem.* **1984**, *26*, 445.
- (49) Eisenhut, H.; Mitchell, H. L.; Traficante, D. D.; Kaufman, R. J.; Deutch, J. M.; Whitesides, G. M. *J. Am. Chem. Soc.* **1974**, *96*, 5385.
- (50) Cowley, A. H.; Braun, R. W.; Gilje, J. W. *J. Am. Chem. Soc.* **1975**, *97*, 434.
- (51) Peake, S. C.; Schmutzler, R. *J. Chem. Soc. A* **1970**, 1049.
- (52) Cited in ref 14a, p. 147.
- (53) Shannon, R. D. *Acta Crystallogr.* **1976**, *A32*, 751.
- (54) Johnson, S. E.; Day, R. O.; Holmes, J. M.; Holmes, R. R. *Inorg. Chem.*, following paper in this issue.
- (55) Harland, J. J.; Payne, J. S.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1987**, *26*, 760.
- (56) Janes, N.; Oldfield, E. *J. Am. Chem. Soc.* **1985**, *107*, 6769.