

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

1,2,4-SELENADIPHOSPHOLES—NOVEL HETEROCYCLIC COMPOUNDS CONTAINING LOW-COORDINATED PHOSPHORUS ATOMS¹

M. Regitz^a; S. Krill^a

^a Fachbereich Chemie der Universität Kaiserslautern, Kaiserslautern, Germany

To cite this Article Regitz, M. and Krill, S.(1996) '1,2,4-SELENADIPHOSPHOLES—NOVEL HETEROCYCLIC COMPOUNDS CONTAINING LOW-COORDINATED PHOSPHORUS ATOMS¹', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 115: 1, 99 — 103

To link to this Article: DOI: 10.1080/10426509608037957

URL: <http://dx.doi.org/10.1080/10426509608037957>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

1,2,4-SELENADIPHOSPHOLES—NOVEL HETEROCYCLIC COMPOUNDS CONTAINING LOW-COORDINATED PHOSPHORUS ATOMS¹

M. REGITZ and S. KRILL

*Fachbereich Chemie der Universität Kaiserslautern, Erwin-Schrödinger-Strasse,
D-67663 Kaiserslautern, Germany*

Dedicated to Professor J. G. Verkade on the occasion of his 60th birthday

(Received January 25, 1996; in final form February 27, 1996)




The thermal reactions (at 120°C) of the 1,2,3-selenadiazole **8** with the phosphalkynes **6a** and **6b** gave rise to the previously unknown title compounds in 17% and 16% yield, respectively. These compounds are formed by a sequence of [3 + 2]-cycloreversion and cycloaddition reactions.

Key words: Low-coordinated phosphorus, 1,2,4-selenadiphospholes, phosphalkyne selenides, selenoxophosphinidines, phosphalkynes.

INTRODUCTION


Heterophospholes play a major role in the development of the chemistry of low-coordinated phosphorus. Condensation reactions^{2,3} and [3 + 2]-cycloaddition processes, in particular with phosphalkynes,^{4,5} have provided accesses to a wide range of phospholes containing one or more additional heteroatoms in the ring.

In this short paper, we report on the discovery of a previously unknown class of compounds, the 1,2,4-selenadiphospholes **3**.



A 1,2,4-azadiphosphole **1** had been obtained previously by the spontaneous dimerization of *tert*-butylaminophosphaacetylene.⁶ Furthermore, some 1,2,4-thiadiphospholes **2** carrying both identical and different C-substituents at the 3 and 5 positions have been prepared in the last ten years.^{7–11}

The starting point for our investigations in this field was the [3 + 2]-cycloaddition of the selenoxocarbenes **5A** \rightleftharpoons **5B** (or the respective 1,3-diradicals) to phosphalkynes **6**.¹²



The dipoles were generated photolytically or thermally from the 1,2,3-selenadiazoles **4** by cleavage of nitrogen and then underwent regiospecific cycloaddition reactions with the P/C triple bonds of **6** to furnish the 1,3-selenaphospholes **7**.¹² However, when we subjected the parent, unsubstituted 1,2,3-selenadiazole **8** to reaction with the phosphalkynes **6** under comparable conditions, we were confronted with the unexpected and unforeseeable formation of the title compounds **15a** and **b**.

RESULTS

The reactions of **8** with a threefold excess of **6a** or **b** were performed in the absence of a solvent and, after column chromatographic work-up on silica gel, furnished the 1,2,4-selenadiphospholes **15a** and **b** in the form of pale yellow oils. The compositions of the products were established by elemental analysis and mass spectroscopy. Volumetric determination of the amount of gas liberated during the reactions revealed that, in addition to nitrogen, a second equivalent of a gaseous compound (in these cases, acetylene) was released during the reactions. Nevertheless, the yields of the product isolated were unexpectedly low.

The structures of the 1,2,4-selenadiphospholes **15a** and **b** are unequivocally substantiated by the NMR spectroscopic data; this will be shown below for example of the di-*tert*-butyl derivative **15a**. In the ^{77}Se NMR spectrum, the heteroatom gives rise to a double doublet signal at $\delta = 723.0$ with a $^1J_{\text{Se,P}}$ coupling constant of 501.7 Hz. This J value is in good agreement with those of other compounds containing a comparable structural unit.^{13,14} The $^2J_{\text{Se,P}}$ value of 62.3 Hz is also of the expected magnitude.¹⁴ In the ^{31}P NMR spectrum the $\lambda^3\sigma^2$ -phosphorus nuclei give rise to signals at very low field,¹³ namely $\delta = 291.2$ (P-2) and 260.4 (P-4). Both signals are split

Finally, the ^{13}C NMR spectrum of **15a** confirmed the sequence of atoms deduced above. The splitting of the signal for C-3 into a double doublet ($\delta = 222.0$) with $^1J_{\text{C,P}}$ couplings of 82.2 and 60.4 Hz as well as that of the C-5 signal ($\delta = 215.6$)

with $^1J_{\text{C,P}} = 67.2$ Hz and $^2J_{\text{C,P}} = 6.1$ Hz are in complete harmony with the proposed structure. The remaining ^{13}C NMR data are directly related to the structure and are given in the experimental section.

It can safely be assumed that, in the primary step of the thermal reaction, the selenadiazole **8** decomposes to nitrogen and the selenoxocarbene $\mathbf{9A} \rightleftharpoons \mathbf{9B}$; this is followed by the [3 + 2]-cycloaddition of the sextet dipole to the first equivalent of the phosphalkyne **6** furnishing the 1,3-selenaphosphole **11**. Evidence in support of the intermediate formation of **11** is provided by a ^{31}P NMR spectrum of the mixture of **8** + **6a** recorded after a reaction time of 1 h: in addition to the phosphorus signals of **15a**, this spectrum also contains a signal at $\delta = 206.6$ which is in the same region as the signals of previously known compounds of this type (such as, e.g. **7**).¹² The thermal fragmentation of **11** into acetylene and the selenoxophosphinidine $\mathbf{13A} \rightleftharpoons \mathbf{13B}$ is followed by the 1,3-dipolar cycloaddition of the latter species to the second equivalent of the phosphalkyne **6** which furnishes the 1,2,4-selenadiphosphole **15**. Of course, it cannot be excluded that the reaction sequence $\mathbf{9} + \mathbf{6} \rightarrow \mathbf{10} \rightarrow \mathbf{12} (+\mathbf{6}) \rightarrow \mathbf{15}$ may also, to a minor extent, contribute to the formation of the product. The [3 + 2]-cycloaddition of the thiobenzoylphosphinidine (**13**, R = Ph, S in place of Se), which was generated by a completely independent route, with phenylphosphacetylene (**6**, R = Ph) has been reported.^{8,9}

EXPERIMENTAL

The reactions were carried out in an argon (purity: $\geq 99.998\%$) atmosphere in previously evacuated and baked-out Schlenk pressure tubes. The bulb-to-bulb distillations were performed with a Büchi 6KR50 apparatus, the temperatures given are oven temperatures. Microanalyses were determined using a Perkin Elmer 240 apparatus. The ^1H NMR spectra were obtained on a Varian 390 (90 MHz) spectrometer with TMS as internal standard, the ^{13}C NMR spectra on a Bruker AMX 400 (100.67 MHz) spectrometer with TMS as internal standard, the ^{31}P NMR spectra on a Bruker WP 200 (80.8 MHz) spectrometer with 85% H_3PO_4 as external standard, and the ^{77}Se NMR spectra on a Bruker AMX 400 (76.36 MHz) spectrometer with $\text{Se}(\text{CH}_3)_2$ as external standard. The mass spectra were recorded on a Finnigan MAT 90 spectrometer.

3,5-Di-tert-butyl-1,2,4-selenadiphosphole (15a). 1,2,3-Selenadiazole (**8**; 0.80 g, 6.00 mmol) and *tert*-butylphosphacetylene¹⁵ (**6a**; 2.38 ml, 18.2 mmol, threefold excess) were heated at 120°C for 5 h in a Schlenk pressure tube. After cooling of the reaction mixture to 25°C , the evolution of two equivalents of nitrogen plus acetylene can be determined by gas volumetric analysis. After separation of the excess phosphalkyne at $30^\circ\text{C}/10^{-3}$ mbar, the residue was taken up in *n*-hexane/diethyl ether (1:1) and subjected to column chromatography on silica gel (0.063–0.2 mm) with 700 ml of the same solvent mixture. After elution of polymeric components, a pale yellow fraction was obtained which, apart from **15a**, did not contain any other phosphorus-containing compounds (^{31}P NMR monitoring). Similarly, the subsequent fractions also did not contain any phosphorus compounds. After evaporation of the eluted fraction at $30^\circ\text{C}/10^{-3}$ mbar and bulb-to-bulb distillation of the residue at $150^\circ\text{C}/10^{-3}$ mbar, 0.29 g (17%) of product **15a** were obtained as a pale yellow oil. Found (%): C 42.9, H 6.4; calcd. for $\text{C}_{10}\text{H}_{18}\text{P}_2\text{Se}$ (%): C 43.0, H 6.5. ^1H NMR (CDCl_3): $\delta = 1.70$ d (9H, 5-*t*-Bu, $^4J_{\text{H,P}} = 1.8$ Hz), 1.73 dd (9H, 3-*t*-Bu, $^4J_{\text{H,P}} = 0.7$ and 2.6 Hz, respectively). ^{13}C NMR (CDCl_3): $\delta = 35.5$ dd [$3-\text{C}(\text{CH}_3)_3$, $^3J_{\text{C,P}} = 1.7$ and 9.7 Hz, respectively], 35.8 d [$5-\text{C}(\text{CH}_3)_3$, $^3J_{\text{C,P}} = 11.8$ Hz], 42.9 pseudo-t [$3-\text{C}(\text{CH}_3)_3$, $^2J_{\text{C,P-2}} = ^2J_{\text{C,P-4}} = 19.8$ Hz], 43.4 dd [$5-\text{C}(\text{CH}_3)_3$, $^2J_{\text{C,P}} = 16.7$ Hz, $^3J_{\text{C,P}} = 6.3$ Hz], 215.6 dd (C-5, $^1J_{\text{C,P}} = 67.2$ Hz, $^2J_{\text{C,P}} = 6.1$ Hz), 222.0 dd (C-3, $^1J_{\text{C,P}} = 82.2$ Hz, $^1J_{\text{C,P}} = 60.4$ Hz). ^{31}P NMR (CDCl_3): $\delta = 260.4$ d (P-4, $^2J_{\text{P,P}} = 49.2$ Hz), 291.2 dd (P-2, $^2J_{\text{P,P}} = 49.2$ Hz, $^1J_{\text{P,Se}} = 441$ Hz). ^{77}Se NMR (CDCl_3): $\delta = 773.0$ dd (Se-1, $^1J_{\text{Se,P}} = 501.7$ Hz, $^2J_{\text{Se,P}} = 62.3$ Hz). MS (70 eV): $m/z = 279.8$ (61%, M^+).

3,5-Di-tert-pentyl-1,2,4-selenadiphosphole (15b). In analogy to the above procedure starting from 1,2,3-selenadiazole (**8**, 0.80 g, 6.00 mmol) and *tert*-pentylphosphacetylene¹⁶ (**6b**; 7.1 g, 17.5 mmol, as a 45% solution in hexamethyldisiloxane) 0.29 g (16%) of product **15b** were obtained as a pale yellow oil. Found (%): C 46.9, H 7.3; calcd. for $\text{C}_{12}\text{H}_{22}\text{P}_2\text{Se}$ (%): C 46.8, H 7.2. ^1H NMR (CDCl_3): $\delta = 0.75$, 0.80 each t (each 3H, CH_2-CH_3 , $^3J_{\text{H,H}} = 7.5$ Hz), 1.50 d [6H, 5- $\text{C}(\text{CH}_3)_2$, $^4J_{\text{H,P}} = 2.0$ Hz], 1.55 dd [6H,

3-C(CH₃)₂, ⁴J_{HP} = 0.5 and 2.7 Hz, respectively], 1.85, 1.95 each q (each 2H, CH₂—CH₃, ³J_{H,H} = 7.5 Hz). ¹³C NMR (CDCl₃): δ = 9.5 (CH₂—C(CH₃)₂), 32.7 dd [3-C(C(CH₃)₂)₂, ³J_{C,P} = 9.2 and 14.3 Hz, respectively], 33.0 d [5-C(C(CH₃)₂)₂, ³J_{C,P} = 13.1 Hz], 40.6 pseudo-t [3-C—C(CH₃)₂—CH₃, ³J_{C,P-2} = ³J_{C,P-4} = 8.3 Hz], 40.9 d [5-C—C(CH₃)₂—CH₃, ³J_{C,P} = 6.8 Hz], 46.5 pseudo-t [3-C(C(CH₃)₂)₂, ²J_{C,P-2} = ²J_{C,P-4} = 18.6 Hz], 46.7 dd [5-C(CH₃)₂C₂H₅, ²J_{C,P} = 17.5 Hz, ³J_{C,P} = 7.0 Hz], 214.1 dd (C-5, ¹J_{C,P} = 73.4 Hz, ²J_{C,P} = 5.9 Hz), 220.4 d (C-3, ¹J_{C,P} = 72.7 and 80.5 Hz, respectively). ³¹P NMR (CDCl₃): δ = 261.0 d (P-4, ²J_{PP} = 48.1 Hz), 295.6 dd (P-2, ²J_{PP} = 48.1 Hz, ¹J_{P,Se} = 439.5 Hz). ⁷⁷Se NMR (CDCl₃): δ = 723.0 dd (¹J_{Se,P} = 439.5 Hz, ²J_{Se,P} = 62.3 Hz). MS (70 eV): m/z = 293.8 (70%, M⁺).

ACKNOWLEDGEMENTS

We thank the Fonds der Chemischen Industrie for generous financial support of our work and the Landesregierung Rheinland-Pfalz for a post-graduate grant to S.K.

REFERENCES

1. Organophosphorus Compounds 113; for part 112, see: B. Breit and M. Regitz, *Chem. Ber.*, **129** (1996), in press.
2. A. Schmidpeter and K. Karaghiosoff, in "Multiple Bonds and Low Coordination in Phosphorus Chemistry," M. Regitz and O. J. Scherer, eds., Thieme, Stuttgart, 1990, p. 258 ff.
3. R. K. Bansal, K. Karaghiosoff, N. Gandhi and A. Schmidpeter, *Synthesis*, 361 (1995).
4. M. Regitz, in "Multiple Bonds and Low Coordination in Phosphorus Chemistry," M. Regitz and O. J. Scherer, eds., Thieme, Stuttgart, 1990, p. 58 ff.
5. M. Regitz, *Chem. Rev.*, **90**, 191 (1990).
6. A. S. Ionkin, S. N. Ignat'eva, I. A. Litvinov, V. A. Naumov and B. A. Arbusov, *Heteroatom Chemistry*, **2**, 577 (1991).
7. R. Appel and R. Moors, *Angew. Chem.*, **98**, 570 (1986); *Angew. Chem., Int. Ed. Engl.*, **25**, 567 (1986).
8. G. Märkl and W. Hölzl, *Tetrahedron Lett.*, **30**, 4501 (1989).
9. G. Märkl, W. Hölzl, A. Kallmünzer, M. L. Ziegler and B. Nuber, *Tetrahedron Lett.*, **33**, 4421 (1992).
10. E. Lindner, C. Haase, H. A. Mayer, M. Kemmler, R. Fawzi and M. Steinmann, *Angew. Chem.*, **105**, 1521 (1993); *Angew. Chem., Int. Ed. Engl.*, **32**, 1424 (1993).
11. E. Lindner, T. Schlenker and C. Haase, *J. Organomet. Chem.*, **464**, C31 (1994).
12. B. Burkhart, S. Krill, Y. Okano, W. Ando and M. Regitz, *Synlett*, 356 (1991).
13. K. Karaghiosoff and A. Schmidpeter, *Phosphorus and Sulfur*, **36**, 217 (1988).
14. L. Quin, "31P-NMR Spectroscopy," Wiley, New York, 1988.
15. G. Becker, G. Gresser and W. Uhl, *Z. Naturforsch.*, **36b**, 16 (1981). Optimized procedure: W. Rösch, U. Hees and M. Regitz, *Chem. Ber.*, **120**, 1645 (1987).
16. T. Allspach, Dissertation, University of Kaiserslautern, 1986.