This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-

41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t713618290

Synthesis of Thioacrolein S-Oxide and Thioketene S-Oxide using FVT

Frederic Minassian; Jocelyne Levillain; Nadia Pelloux-Leon; Jean-Louis Ripoll; Yannick Vallee

To cite this Article Minassian, Frederic , Levillain, Jocelyne , Pelloux-Leon, Nadia , Ripoll, Jean-Louis and Vallee, Yannick(1999) 'Synthesis of Thioacrolein S-Oxide and Thioketene S-Oxide using FVT', Phosphorus, Sulfur, and Silicon and the Related Elements, 153: 1, 335 - 336

To link to this Article: DOI: 10.1080/10426509908546458 URL: http://dx.doi.org/10.1080/10426509908546458

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of Thioacrolein S-Oxide and Thioketene S-Oxide using FVT

FREDERIC MINASSIAN^a, JOCELYNE LEVILLAIN^b, NADIA PELLOUX-LEON^a, JEAN-LOUIS RIPOLL^b and YANNICK VALLEE^a

^aL.E.D.S.S., Université Joseph Fourier-CNRS, B.P. 53X, 38041 Grenoble, France and ^bL.C.M. T., ISMRA-CNRS, Université de Caen, 14050 Caen, France

FVT was used to synthesize thioacrolein S-oxide and thioketene S-oxide. These reactive species were characterized at low temperature by IR and/or NMR spectroscopies.

INTRODUCTION

In order to synthesize thioacrolein S-oxide 1 and thioketene S-oxide 2, we prepared the retro Diels-Alder precursors 3, 4 and 5.

THIOACROLEIN 5-OXIDE1:

FVT of the vinylic sulfoxide 3 at 600°C quantitatively gave 9,10-dimethylanthracene. Volatile products were trapped at -196°C and analyzed by NMR spectroscopy. The ¹H NMR and ¹³C NMR spectra were in good agreement with the structure of thioacrolein S-oxide. They revealed the formation of two isomeric sulfines. The ratio 12/1E was 78/22. When the sample was allowed to stand at room temperature, the sulfine signals slowly disappeared and a new product was mainly formed. Signals observed by ¹H NMR and ¹³C NMR were tentatively assigned to the dimeric structure 6.

THIOKETENE S-OXIDE2:

The major product obtained by FVT of precursor 4 was the corresponding ketone (60% yield estimated by NMR) resulting from sulfur extrusion. However some anthracene was also obtained and bands at 1700 and 1040 cm⁻¹ were observed in the IR spectrum of the volatile products. These bands were tentatively assigned to thioketene S-oxide. Furthermore the same IR bands were observed after FVT of the second precursor 5 at 650°C. Interestingly when this precursor was thermolyzed at lower temperature (500°C) formation of the sulfine 4 was noticed. We propose that this rearrangement occurs via homolytic cleavage of the dibenzylic C-S bond.

References

- N. Pelloux-Léon, R. Arnaud, J.-L. Ripoll, P. Beslin, Y. Vallée, Tetrahedron Lett., 38, 1385-1388 (1997).
- [2] N. Pelloux-Léon, F. Minassian, J. Levillain, J.-L Ripoll, Y. Vallée, Tetrahedron Lett., 38, 4813–1816 (1998).