

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Deuterium Studies in the Cyclization and Isomerization of 3-Propargylmercapto-1,2,4-Triazines to Thiazolo[3,2-B]-[1,2,4] Triazines

M. M. Heravi^a; M. Shafaie^b; P. Khosrofar^b; M. Ghassemzdeh^c

^a Department of Chemistry, School of Sciences, Azzahra University Vanak, Tehran, Iran ^b Department of Chemistry School of Sciences, Ferdowsi University of Mashhad, Iran ^c Chemistry & Chemical, Engineering Research Center of Iran, Tehran Iran

To cite this Article Heravi, M. M. , Shafaie, M. , Khosrofar, P. and Ghassemzdeh, M.(2000) 'Deuterium Studies in the Cyclization and Isomerization of 3-Propargylmercapto-1,2,4-Triazines to Thiazolo[3,2-B]-[1,2,4] Triazines', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 167: 1, 21 – 27

To link to this Article: DOI: 10.1080/10426500008082384

URL: <http://dx.doi.org/10.1080/10426500008082384>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

DEUTERIUM STUDIES IN THE CYCLIZATION AND ISOMERIZATION OF 3-PROPARGYLMERCAPTO-1,2,4-TRIAZINES TO THIAZOLO[3,2-b]-[1,2,4] TRIAZINES

M.M. HERAVI^{a*}, M. SHAFAYE^b, P. KHOSROFAR^b and
M. GHASSEMZDEH^c

^aDepartment of Chemistry, School of Sciences, Azahra University, Vanak, Tehran, Iran, ^bDepartment of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Iran and ^cChemistry & Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran Iran

(Received April 04, 2000)

Transformation of 3-propargylmercapto-1,2,4-triazin-5(2H)-one (2; R=H or Me) to the corresponding 2-dihydro-3-methylenethiazolo[3,2-b][1,2,4]triazin-7-one (3; R=H or Me) is performed in a dilute solution of triethylamine in MeOH. Compound (3; R=Me) was isomerized to 3,6-dimethyl-thiazolo[3,2-b][1,2,4]-triazin-5-one (4; R=Me) in 5% aqueous sodium hydroxide. Treatment of (3; R=Me) with 5% sodium deuteroxide gave 6-methyl-3-monodeuterated methyl-2-D-thiazolo[3,2-b][1,2,4]-triazin-7-one (5, R=Me).

Keywords: Deuterium studies; cyclization; Propargylmercapto-triazines; Thiazolo-triazines

INTRODUCTION







Synthesis of a variety of bicyclic compounds derived from 1,2,4-triazin has been reported from our laboratory.¹⁻⁶ In continuation of our interest in this field, we now wish to report the base catalyzed cyclization and isomerization of 3-propargyl-mercapto-1,2,4-triazin-5-one (2; R=H or Me) to (3; R=H or Me) and (4; R=Me) respectively. To prove the suggested mechanism, deuterium studies have been implemented. Due to the wide

* To whom correspondence should be addressed.


ranging physiological activities of bicyclic compounds derived from 1,2,4-triazine, this attempt was thought to be worthwhile.

1,2,4-Triazin-3(2H)-thion-5(4H)-one (1; R=H or Me)⁷ was condensed with propargylbromide in presence of sodium methoxide to afford the corresponding 3-propargylmercapto-1,2,4-triazin-5(2H)-one (2; R=H or Me). When (2; R=Me) was refluxed in aqueous sodium hydroxide for 4 hrs, a solid was isolated, that was crystallized from ethanol to give a single white crystalline compound. The compound was identified as 3,6-dimethyl-7H-thiazolo [3,2-b][1,2,4]- triazin-7-one (4; R=Me).

This compound has already been synthesized through the reaction of (1; R=Me) with bromoacetone in the presence of sodium carbonate and subsequent dehydration of the hydroxy compound (6).



By elucidation of the structure, it can be assumed that the base abstracts a hydrogen from the nitrogen 2 of (2). The nitrogen then attacks the acetylenic group to give the cyclized intermediate (3; R=Me). Isomerization and aromatization of the intermediate produces compound (4; R=Me) (Scheme 1).



SCHEME 1

For isolation of (3; R=H and Me) time and concentration of the base were decreased but it could not be isolated. Compound (3; R=H and Me) was obtained when triethylamine in ethanol was used as a base and stopped at an appropriate conversion. Exclusive cyclization on the N-2

atom is reminiscent of the alkylation of 3-methylmercapto-1,2,4-triazin-5(2H)-ones with alkyl halides under basic condition.⁸ The assignment of Structure (3; R=H and Me) was based on their physical and spectral data. The exo methylene protons in (3) separated by 0.5 ppm and appeared as two doublets. When (3; R=Me) was refluxed in 5% aqueous sodium hydroxide compound (4; R=Me) was quantitatively obtained.


To establish the mechanisms deuterium studies were contemplated. Compound (3; R=Me) was refluxed in 5% sodium deuterioxide in D₂O. The isolated product showed m/e at 183 corresponding to exchange of two ¹H with ²H. In the ¹H nmr of this compound the two exo methylene signals and a signal at δ 4.1 for CH₂ in the five membered heterocycle disappeared and there was no sign for any aromatic proton. Based on these spectroscopic data we could assign structure (5; R=Me) for the obtained compound. The suggested mechanism is as follows (Scheme 2).

EXPERIMENTAL

The melting points are uncorrected and were obtained by a Kofler Heizbank Richert type 7841, melting point apparatus. Ir spectra were obtained on a Pye Unicam SP 1100 and 4300 Shimadzu spectrometer. The ¹H nmr spectra were recorded on a Varian 50 A Spectrometer using TMS as internal reference and mass spectra were scanned on a Varian Mat CH-7 instrument at 70 eV.

3-Propargylmercapto-1,2,4-triazin-5(2H)-one (2; R=H)

Sodium (1 g) was dissolved in MeOH (100 mL). To this solution, the compound (1; R=H) (4.5 g, 0.035 mol) was added. To this mixture propargylbromide (4.65 mL, of 80% solution in toluene) was added dropwise at room temperature. The reaction mixture was stirred for 4 hrs at ambient temperature. The solid was filtered, washed with water and crystallized from methanol to afford the title compound. Yield: 2.2 g (38%), mp.: 171 °C (from methanol). ¹H nmr δ (d₆-DMSO), 3.1 (s, 1H, CH), 3.9 (s, 2H, CH₂), 7.5 (s, 1H, H of triazine ring). M+. m/e: 167 (rel. intensity), 167 (2), 164 (18), 137 (13), 110 (38), 106 (100), 82 (20), 78 (23), 68 (41), 67 (90). IR (KBr disc, cm⁻¹) 3230, 2150, 1620, 1425, 1365.



SCHEME 2

6-Methyl-3-propargylmercapto-1,2,4-triazin-5(2H)-one (2; R=Me)

Sodium (1 g) was dissolved in MeOH (100 mL). To this solution, the compound (1; R=Me) (5 g, 0.035 mol) was added. To this mixture propargylbromide (4.65 mL, of 80% solution in toluene) was added dropwise at room temperature. The reaction mixture was stirred for 4 hrs at ambient temperature. The solid was filtered, washed with water and crystallized from methanol to afford the title compound. Yield: 5.5 g (87%), mp.: 180–1°C (from ethanol). ¹H nmr δ (d₆-DMSO), 2 (s, 3H, Me), 3.1 (s, 1H, CH), 3.9 (s, 2H, CH₂). M+. m/e: 181 (rel. intensity), 181 (M+, 100), 151 (4), 141 (3), 140 (25), 75 (8), 71 (8), 71 (6). IR (KBr disc, cm⁻¹) 3456, 2262, 1625, 1512, 1465, 1361, 1271.

3,6-Dimethyl-7H-thiazolo[3,2-b][1,2,4]-triazin-7-one (4; R=Me)

Compound (2; R=Me) (1 g, 0.0055 mol) was added to a 5% sodium hydroxide solution (20 mL) and the reaction mixture was refluxed for 4 hrs. The solution was cooled to room temperature and the solid was filtered, washed with water and crystallized from MeOH. Yield: 0.7 g (70%), mp.: 226°C. ^1H nmr δ (CDCl_3): 2.2 (s, 1H, 2Me), 6.5 (s, 1H, CH). M+ m/e 181 (rel. intensity) 181 (2), 174(36), 178 (73), 150 (9), 137 (100), 106 (9), 68 (40), 67 (40), 43 (9).

2-Dihydro-3-methylene-thiazolo-[3,2-b][1,2,4]-triazin-7-one (3; R=H)

Compound (2; R=H) (1.67 g, 0.001 mol) was dissolved in a mixture of NaOH (0.2 g) in MeOH (100 mL). The reaction mixture was refluxed for 4 hrs. The solvent was evaporated and the crude was directly subjected to column chromatography on silicagel using CHCl_3 :MeOH. 90:10 as eluent. Yield: 0.6 g (32%), mp. 154–5 °C. ^1H nmr δ (CDCl_3): 4.2 (s, 2H, CH_2), 4.9 (d, 1H, exo methylene), 5.5 (d, 1H, exo methylene), 7.5 (s, 1H, H of triazine ring). M+: m/e (rel. intensity): 167 (8), 165 (100), 142 (90), 12? (24), 72 (92), 71 (56), 67 (40). IR (KBr disc, cm^{-1}): 1650, 1470, 1370

2-Dihydro-3-methylene-6-methyl-thiazolo-[3,2-b][1,2,4]-triazin-7-one (3; R=Me)

Compound (2; R=Me) (5 g, 0.027 mol) was dissolved in a mixture of triethylamine (50 mL) and ethanol (250 mL). The reaction mixture was refluxed for 2 hrs. The solvent was evaporated and the crude was directly subjected to column chromatography on silicagel. a mixture of CHCl_3 :MeOH 90:10 was used as an eluent.

Yield: 2.9 g (58%). Mp. 193°C, ^1H nmr δ (CDCl_3) 2.2 (s, 3H, Me), 4.1 (s, 2H, CH_2), 4.8 (d, 1H, exo methylene), 5.5 (d, 1H, exo methylene). M+ m/e (rel. Intensity) 181 (5), 179 (5), 178 (50), 137 (64), 68 (100), 67 (36). IR (KBr disc, cm^{-1}): 2260, 1620, 1460, 1340.

6-Methyl-3-monodeuterated methyl-2D-thiazolo[3,2-b][1,2,4]-triazin-5-one (5; R=Me)

Compound (3; R=Me) (0.25 g, 0.0013 mol) was dissolved in 5% sodium deuteroxide in D_2O (10 mL). The reaction mixture was refluxed for 2 hrs

and chilled in an ice bath. The precipitated solid was filtered and washed with water to afford the title compound. Yield: 0.15 g (60%). Mp: 224 °C. ^1H nmr δ (CDCl_3) 2.2 (s, 3H, Me), 2.3 (s, 2H, CH_2D). M^+ m/e 183.

References

1. M.M. Heravi, M. Bakavoli, J. Chem. Res. 480 (1995).
2. M.M. Heravi, M. Bakavoli, M. Shafaie, M.M. Sadeghi, A.R. Khoshdast, Indian J. Chem. 35, 1260 (1996).
3. M.M. Heravi, K. Aghapoor, M.A. Nooshabadi, M.M. Mojtahedi, Monatsh. Chem. 128, 1143 (1997).
4. M.M. Heravi, K. Aghapoor, M.A. Nooshabadi, Synth. Commun. 28, 233 (1998).
5. M.M. Heravi, M. Rahimizadeh, E. Iravani, M. Ghassemzadeh, Indian J. Chem. 8, 75 (1999).
6. M.M. Heravi, M.M. Sadeghi, M. Hazrati, M. Ghassemzadeh, Indian J. Heterocyclic Chem. 8, 339 (1999).
7. J. Gut, Collection Czechoslov. Chem. Commun. 23, 1588 (1958).
8. G. Doledchall, G. Harvyk, M. Hamori, K. Lenyert, A. Wolfner, Acta Chem. Acad. Sci. Hung. 53, 385 (1967).