

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### Substitution Reactions of Mono- and Trichloro Acetic Acids with Ammonium Dialkyl (Alkylene) Dithiophosphates

Neelam Harkut<sup>a</sup>; Padam N. Nagar<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of Rajasthan, Jaipur, India

**To cite this Article** Harkut, Neelam and Nagar, Padam N.(2007) 'Substitution Reactions of Mono- and Trichloro Acetic Acids with Ammonium Dialkyl (Alkylene) Dithiophosphates', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 182: 1, 219 – 226

**To link to this Article:** DOI: 10.1080/10426500600892693

**URL:** <http://dx.doi.org/10.1080/10426500600892693>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Substitution Reactions of Mono- and Trichloro Acetic Acids with Ammonium Dialkyl (Alkylene) Dithiophosphates

Neelam Harkut  
Padam N. Nagar

Department of Chemistry, University of Rajasthan, Jaipur, India

*A reaction of mono- and trichloro acetic acid with ammonium salts of alkylene (dialkyl) dithiophosphate;  $[\text{OGOPS}_2\text{NH}_4]$ ;  $G = -\text{CH}_2\text{CH}_2\text{CHMe}-$ ,  $-\text{C}(\text{Me})_2\text{C}(\text{Me})_2-$ ,  $-\text{CH}_2\text{C}(\text{Me})_2\text{CH}_2-$ , and  $-\text{C}(\text{Me})_2\text{CH}_2\text{CHMe}-$ ;  $(\text{RO})_2\text{PS}_2\text{-NH}_4$ ; and  $\text{R} = \text{C}_2\text{H}_5$ ,  $\text{C}_3\text{H}_7$ ,  $i\text{-C}_3\text{H}_7$ ] in a 1:1 molar ratio in refluxing benzene solution yields low melting solids and light-yellow oily liquids of the type  $[(\text{RO})_2\text{PS}_2\text{R}'$  and  $\text{OGOPS}_2\text{R}'$ , where  $\text{R}' = \text{CH}_2\text{COOH}$  and  $\text{Cl}_2\text{CCOOH}]$ , which are hygroscopic in nature. These newly synthesized complexes have been characterized by physicochemical and spectroscopic techniques (MW, IR,  $^1\text{H}$ , and  $^{31}\text{P}$ NMR) On the basis of the previously discussed studies, the formation of a  $\text{P}-\text{S}-\text{C}$  chemical bond has been established.*

**Keywords** [(Dialkoxyposphorothioyl)thio]acetic acid; Dichloro alkylene dithiophosphato acetic acid; IR spectra; NMR spectra

## INTRODUCTION

Earlier investigations on a variety of metal, organometal, and organic derivatives of alkylene(dialkyl)dithiophosphates<sup>1–17</sup> yielded interesting chemical-bonding patterns as well as biological applications. It has been observed that biological activity is markedly governed by the nature of alkyl/aryl(substituted) groups substituted on an alkylene(alkyl) dithiophosphato moiety.


In view of this, it was considered of interest to extend the present course of investigations on the syntheses of mono-, chloro-, and trichloro acetic acid derivatives of ammonium alkylene(dialkyl) dithiophosphates.

Received March 31, 2006; accepted June 26, 2006.


Address correspondence to Neelam Harkut, Department of Chemistry, University of Rajasthan, Jaipur, 302004 India. E-mail: neelam.gats2005@rediffmail.com

## RESULTS AND DISCUSSION

Dialkyl dithiophosphato acetic acid and alkylene dithiophosphato acetic acid have been synthesized by reacting monochloroacetic acid with ammonium dialkyl (alkylene) dithiophosphates in refluxing benzene (8–6 h). These displacement reactions appear to be slow due to the high acidic strength of monochloroacetic acid, which generates chloride ions very slowly. No change in the reactivity has been observed even when using acetonitrile as a solvent.



These new derivatives are white-colored low-melting solids that are hygroscopic in nature and are soluble in common organic solvents. Similarly, compounds of the type dichloro dialkyl dithiophosphato acetic acid and dichloro dialkylene dithiophosphato acetic acid have been synthesized by reacting trichloroacetic acid with ammonium dialkyl/alkylene dithiophosphates in refluxing benzene for 20–22 h. Due to the high acidic strength of trichloroacetic acid, the substitution of a chloride ion is difficult and thus requires a longer time refluxing for completion.



An attempt has been made to synthesize 1:2 and 1:3 derivatives of trichloroacetic acid, but it has been observed that in this reaction, always and only 1:1 product has been isolated. The products formed are pale-yellow oily liquids that are hygroscopic in nature and are soluble in common organic solvents (tetrahydrofurane, ethanol, acetone etc.).

## IR Spectra

IR spectra of the newly synthesized derivatives show the following characteristic changes (Table I).

1. A broad and intense  $\nu\text{OH}$  absorption band has been present at  $3520\text{ cm}^{-1}$  for the  $-\text{COOH}$  group of monochloroacetic acid derivatives, and a broad and intense absorption band for the  $\nu\text{OH}$  group has been observed in the region  $3300\text{--}3000\text{ cm}^{-1}$  for trichloroacetic acid derivatives.

TABLE I IR Spectral Data of Mono and Tri Chloro Acetic Acid Derivatives of Ammonium Dialky/alkylene Dithiophosphate

| S. No. | Compounds   | $\nu(\text{P}=\text{O}-\text{C})$ | $\nu\text{P}-\text{O}(\text{C})$ | Ring vibrations | $\nu\text{P}=\text{S}$ | $\nu\text{P}-\text{S}$ | $\nu\text{C}=\text{O}$ | $\nu\text{C}-\text{S}$ | $\nu\text{C}-\text{Cl}$ |
|--------|---|-----------------------------------|----------------------------------|-----------------|------------------------|------------------------|------------------------|------------------------|-------------------------|
| 1      | $\text{OCH}_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCH}_2\text{COOH}$                   | 1065                              | 820                              | 970             | 650                    | 580                    | 1716                   | 625                    |                         |
| 2      | $\text{OCH}_2\text{C}(\text{Me})_2\text{CH}_2\text{OP}(\text{S})\text{SCH}_2\text{COOH}$  | 1070                              | 830                              | 960             | 655                    | 535                    | 1716                   | 610                    |                         |
| 3      | $\text{OC}(\text{Me})_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCH}_2\text{COOH}$         | 1025                              | 830                              | 915             | 665                    | 570                    | 1716                   | 610                    |                         |
| 4      | $\text{OC}(\text{Me})_2\text{C}(\text{Me})_2\text{OP}(\text{S})\text{SCH}_2\text{COOH}$   | 1075                              | 820                              | 955             | 665                    | 580                    | 1720                   | 620                    |                         |
| 5      | $(i\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$             | 1025                              | 770                              | —               | 655                    | 560                    | 1726                   | 562                    |                         |
| 6      | $(\text{C}_2\text{H}_5\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$               | 1035                              | 815                              | —               | 655                    | 550                    | 1726                   | 555                    |                         |
| 7      | $(n\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$             | 1025                              | 810                              | —               | 660                    | 530                    | 1726                   | 560                    |                         |
| 8      | $\text{OCH}_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCCl}_2\text{COOH}$                  | 1070                              | 850                              | 990             | 660                    | 590                    | 1720                   | 700                    | 770                     |
| 9      | $\text{OCH}_2\text{C}(\text{Me})_2\text{CH}_2\text{OP}(\text{S})\text{SCCl}_2\text{COOH}$ | 1060                              | 840                              | 960             | 675                    | 540                    | 1720                   | 680                    | 770                     |
| 10     | $\text{OC}(\text{Me})_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCCl}_2\text{COOH}$        | 1060                              | 840                              | 920             | 675                    | 540                    | 1716                   | 690                    | 770                     |
| 11     | $\text{OC}(\text{Me})_2\text{C}(\text{Me})_2\text{OP}(\text{S})\text{SCCl}_2\text{COOH}$  | 1070                              | 845                              | 950             | 675                    | 590                    | 1730                   | 685                    | 780                     |
| 12     | $(i\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$            | 1040                              | 810                              | —               | 645                    | 545                    | 1730                   | 675                    | 750                     |
| 13     | $(\text{C}_2\text{H}_5\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$              | 1045                              | 815                              | —               | 640                    | 550                    | 1725                   | 675                    | 755                     |
| 14     | $(n\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$            | 1040                              | 815                              | —               | 650                    | 555                    | 1725                   | 675                    | 750                     |

2. A comparison of the  $\nu\text{C}=\text{O}$  absorption band present in monochloro acetic acid at  $1736\text{ cm}^{-1}$  indicates a slight shift toward lower wave numbers ( $10\text{--}20\text{ cm}^{-1}$ ). Similarly, a  $\nu\text{C}=\text{O}$  absorption band of trichloroacetic acid present at  $1740\text{ cm}^{-1}$  shows a slight shift toward lower wave numbers ( $10\text{--}20\text{ cm}^{-1}$ ) due to a lower electrical effect of the chlorine atom.
3. The  $\nu(\text{P})\text{--OC}$  and  $\nu\text{P}\text{--O}(\text{C})$  absorption band appears in the regions  $1075\text{--}1025\text{ cm}^{-1}$  and  $830\text{--}770\text{ cm}^{-1}$ , respectively, while a broad absorption band around  $915\text{--}970\text{ cm}^{-1}$  is due to ring vibrations in monochloro acetic acid derivatives. For trichloro acetic acid derivatives, the absorption band appears in the regions  $1070\text{--}1040\text{ cm}^{-1}$  and  $850\text{--}810\text{ cm}^{-1}$ , respectively.
4. The  $\nu\text{C}\text{--Cl}$  absorption band present in the region  $720\text{--}715\text{ cm}^{-1}$  has disappeared in monochloro acetic acid derivatives, and in trichloro acetic acid derivatives, the  $\nu\text{C}\text{--Cl}$  absorption band present in the region  $750\text{--}780\text{ cm}^{-1}$  has shifted toward lower wave numbers ( $10\text{--}15\text{ cm}^{-1}$ ) in comparison to its position in trichloroacetic acid.
5.  $\nu\text{P}=\text{S}$  and  $\nu\text{P}\text{--S}$  absorption bands have been observed at  $665\text{--}650\text{ cm}^{-1}$  and  $580\text{--}530\text{ cm}^{-1}$ , respectively, in monochloro acetic acid derivatives. The  $\nu\text{P}=\text{S}$  and  $\nu\text{P}\text{--S}$  absorption bands have been observed in the region  $675\text{--}640\text{ cm}^{-1}$  and  $590\text{--}540\text{ cm}^{-1}$ , respectively, in trichloro acetic acid derivatives.
6. A new medium intensity absorption band has been observed in the region  $560\text{--}675\text{ cm}^{-1}$ , which was tentatively assigned to phosphorus-sulfur-carbon( $\text{S}=\text{P}\text{--S}\text{--C}$ ) chemical linkage in these derivatives.

## PMR Spectra

PMR spectra of these derivatives have been recorded in  $\text{CDCl}_3$ , and these are tabulated in Table II. The PMR spectra show a multiplet for  $\text{OCH}_2$  and  $\text{OCH}$  protons due to long range coupling with the magnetically active phosphorus atom. In addition to this, the PMR signal presented at  $\delta\ 12\text{--}11.5\text{ ppm}$  is due to an acidic proton of substituted acetic acid. Deammonization has been observed in the PMR spectra of these derivatives, thus showing the formation of the ( $\text{S}=\text{P}\text{--S}\text{--C}$ ) chemical bond.

## $^{31}\text{P}$ NMR Spectra

A  $^{31}\text{P}$  NMR resonance signal in dialkyl (alkylene) dithiophosphato acetic acid has been observed in the range  $50.4\text{ to }62.4\text{ ppm}$ , which exhibits deshielding ( $\delta\ 13\text{--}19\text{ ppm}$ ) in the  $^{31}\text{P}$  chemical shift value in comparison to dithiophosphates ( $70\text{--}104\text{ ppm}$ ).

**TABLE II NMR  $^1\text{H}$  and  $^{31}\text{P}$  Spectral Data of Mono and Tri Chloro Acetic Acid Derivatives of Ammonium Dialkyl/alkylene Dithiophosphate**

| S. No. | Compounds   | $^1\text{H}$ ( $\delta$ ppm)   | $^{31}\text{P}$ ( $\delta$ ppm) |
|--------|---|--|---------------------------------|
| 1      | $\text{OCH}_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCH}_2\text{COOH}$                   | 1.35, d, 3H ( $\text{CH}_3$ ); 2.75–2.98, m, 2H( $\text{CH}_2$ ); 4.2–4.5, m, 3H ( $\text{OCH}$ , $\text{OCH}_2$ ); 12.5, S, 1H( $\text{COOH}$ ) | 62.31                           |
| 2      | $\text{OCH}_2\text{C}(\text{Me})_2\text{CHOP}(\text{S})\text{SCH}_2\text{COOH}$           | 1.5–1.6, S, 6H( $\text{CH}_3$ ); 4.1–4.2, d, 4H( $\text{CH}_2\text{O}$ ); 11.3, S, 1H ( $\text{COOH}$ )  | 60.40                           |
| 3      | $\text{OC}(\text{Me})_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCH}_2\text{COOH}$         | 1.99–2.2, m, 11H ( $\text{CH}_3$ , $\text{CH}_2$ ); 4.0–4.5, m, 1H( $\text{CHO}$ ); 12.5, S, 1H ( $\text{COOH}$ )                                | 62.45                           |
| 4      | $\text{OC}(\text{Me})_2\text{C}(\text{Me})_2\text{OP}(\text{S})\text{SCH}_2\text{COOH}$   | 1.80, S, 12H ( $\text{CH}_3$ ); 12.5, S, 1H ( $\text{COOH}$ )  | 61.35                           |
| 5      | $(i\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$             | 1.48–2.1, d, 12H ( $\text{CH}_3$ ); 11.9, S, 1H ( $\text{COOH}$ )  | 56.49                           |
| 6      | $(\text{C}_2\text{H}_5\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$               | 2.28–3.3, t, 6H ( $\text{CH}_3$ ); 3.7–4.5, m, 4H( $\text{OCH}_2$ ); 12.3, S, 1H ( $\text{COOH}$ )   | 55.70                           |
| 7      | $(n\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCH}_2\text{COOH}$             | 1.52–1.58, t, 6H ( $\text{CH}_3$ ); 2.27–3.1, m, 4H( $\text{CH}_2$ ); 3.5–4.2, m, 4H( $\text{OCH}_2$ ); 12.3, S, 1H( $\text{COOH}$ )             | 54.51                           |
| 8      | $\text{OCH}_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCCl}_2\text{COOH}$                  | 1.4, d, 3H( $\text{CH}_3$ ); 3.0–2.8, m, 2H ( $\text{CH}_2$ ); 4.5–4.9, m, 3H( $\text{OCH}$ , $\text{OCH}_2$ ); 12.1, S, 1H ( $\text{COOH}$ )    | 56.59                           |
| 9      | $\text{OCH}_2\text{C}(\text{Me})_2\text{CH}_2\text{OP}(\text{S})\text{SCCl}_2\text{COOH}$ | 1.24, s, 6H( $\text{CH}_3$ ); 4.2–4.5, d, 4H( $\text{CH}_2\text{O}$ ); 11.9, S; 1H( $\text{COOH}$ )  | 60.21                           |
| 10     | $\text{OC}(\text{Me})_2\text{CH}_2\text{CHMeOP}(\text{S})\text{SCCl}_2\text{COOH}$        | 2.0–2.5, m, 11H ( $\text{CH}_3$ , $\text{CH}_2$ ); 5.0–5.2, m, 1H( $\text{CHO}$ ); 12.1, S, 1H ( $\text{COOH}$ )                                 | 59.41                           |
| 11     | $\text{OC}(\text{Me})_2\text{C}(\text{Me})_2\text{OP}(\text{S})\text{SCCl}_2\text{COOH}$  | 1.52, S, 12H ( $\text{CH}_3$ ); 4.6, m, 2H( $\text{CHOP}$ ); 11.9, S, 1H ( $\text{COOH}$ )   | 58.51                           |
| 12     | $(i\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$            | 1.46–1.5, d, 12H ( $\text{CH}_3$ ); 11.5, S, 1H( $\text{COOH}$ )   | 50.41                           |
| 13     | $(\text{C}_2\text{H}_5\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$              | 1.67–1.98, t, 6H ( $\text{CH}_3$ ); 5.01–5.2, m, 4H ( $\text{OCH}_2$ ); 11.9–12.0, s, 1H ( $\text{COOH}$ )                                       | 55.52                           |
| 14     | $(n\text{-C}_3\text{H}_7\text{O})_2\text{P}(\text{S})\text{SCCl}_2\text{COOH}$            | 1.57–1.70, t, 6H ( $\text{CH}_3$ ); 2.75–3.01, m, 4H ( $\text{CH}_2$ ); 5.5–5.7, m, 4H ( $\text{OCH}_2$ ); 11.9, S, 1H ( $\text{COOH}$ )         | 50.61                           |

$^{31}\text{P}$  NMR spectra of dichloro dialkyl (alkylene) dithiophosphato acetic acid have been recorded in benzene. In a proton-decoupled  $^{31}\text{P}$  spectra, one sharp signal has been observed at  $\delta$  50–60 ppm, which indicates a presence of one type of phosphorus atom (Table II). The  $^{31}\text{P}$  NMR chemical shift values observed in ammonium dialkyl (alkylene) dithiophosphate (77–102 ppm) was shifted upfield ( $\delta$  = 18–28 ppm) in the corresponding trichloro acetic acid derivatives, which indicates the covalent character of a sulphur-carbon linkage as well as an absence of coordinating tendency in these derivatives.

On the basis of the previously discussed studies, the formation of a phosphorus-sulfur-carbon ( $\text{S}=\text{P}-\text{S}-\text{C}$ ) chemical linkage with a free thiophosphoryl group has been tentatively proposed.

## EXPERIMENTAL

Solvents were dried by standard methods. Ammonium salt of dialkyl/alkylene dithiophosphates have been prepared by the method reported in the literature.<sup>17</sup> Sulphur was estimated gravimetrically as barium sulphate (messenger method)<sup>17</sup> and has been purified by vacuum distillation. Molecular weights were determined by the Knauer Vapour Pressure Osmometer using a chloroform solution at 45°C. IR spectra were recorded in Nujol mull ( $4000\text{--}200\text{ cm}^{-1}$ ) on an FTIR spectrophotometer model Megna-IR-550 MICOLAC-USA. Carbon and hydrogen analyses were performed on a Perkin Elemer CHN/O analyzer.  $^1\text{H}$  NMR spectra were recorded in  $\text{CDCl}_3$  solution on a 90 MHz JEOL FX 300 Mhz FT NMR spectrometer using TMS as an internal reference.  $^{31}\text{P}$  NMR were recorded in  $\text{C}_6\text{H}_6$  using  $\text{H}_3\text{PO}_4$  as an external reference. The experimental details of representative compounds are described in the following sections. Analytical results are summarized in Table III.

### Preparation of Ammonium Dialkyl (Alkylene) Dithiophosphate Ligands<sup>17</sup>

These ligands can be prepared by the reactions of  $\text{P}_2\text{S}_5$  with the corresponding alcohol/1,2 or 1,3 glycols in anhydrous benzene by passing dry ammonia gas.

These may be purified by washing with benzene or ether or may be crystallized from a benzene parent alcohol mixture.

### Preparation of [(Diisopropoxyphosphorothioyl) thio] Acetic Acid

A mixture of ammonium diisopropyl dithiophosphate (2.42 g, 10.47 mmole) and monochloro acetic acid (0.98 g, 10.37 mmole) in

TABLE III Synthetic and Analytical Data of Mono and Tri Chloro Acetic Acid Derivatives of Ammonium Diaryl/alkylene Dithiophosphate

| S.No. | Reactant g <sub>1</sub> (mMole)                                       |                                      | R= | ClCH <sub>2</sub> COOH | Product g <sub>2</sub>  | Found (Calculated) |                |                  | M. Wt. Found (Calculated) |
|-------|---|--------------------------------------|----|------------------------|---|--------------------|----------------|------------------|---------------------------|
|       | G=  | RO <sub>2</sub> P(S)SNH <sub>4</sub> |    |                        |   | C                  | H              | S                |                           |
| 1     | —CH <sub>2</sub> CH <sub>2</sub> CHMe—<br>2.20 (10.94)                |                                      |    | 1.02<br>(1079)         | —<br>OCH <sub>2</sub> CH <sub>2</sub> CHMeOP(S)SCH <sub>2</sub> COOH<br>2.22, 85.12                 | 30.15<br>(29.75)   | 5.12<br>(4.54) | 25.76<br>(26.44) | —<br>(242)                |
| 2     | —CH <sub>2</sub> C(Me) <sub>2</sub> CH <sub>2</sub> —<br>5.20 (24.18) |                                      |    | 2.31<br>(24.44)        | —<br>OCH <sub>2</sub> C(Me) <sub>2</sub> CH <sub>2</sub> OP(S)SCH <sub>2</sub> COOH<br>4.94, 79.16  | 33.35<br>(32.81)   | 4.52<br>(5.07) | 24.61<br>(25.00) | 246<br>(256)              |
| 3     | —C(Me) <sub>2</sub> CH <sub>2</sub> CHMe—<br>2.60 (11.35)             |                                      |    | 1.09<br>(11.53)        | —<br>OC(Me) <sub>2</sub> CH <sub>2</sub> CHMeOP(S)SCH <sub>2</sub> COOH<br>2.24, 72.12              | 36.12<br>(35.55)   | 6.01<br>(5.55) | 24.12<br>(23.70) | —<br>(270)                |
| 4     | —C(Me) <sub>2</sub> C(Me) <sub>2</sub> —<br>5.19 (22.66)              |                                      |    | 2.15<br>(22.75)        | —<br>OC(Me) <sub>2</sub> C(Me) <sub>2</sub> OP(S)SCH <sub>2</sub> COOH<br>5.46, 89.01               | 34.92<br>(35.55)   | 4.88<br>(5.55) | 24.32<br>(23.70) | 255<br>(270)              |
| 5     | i-C <sub>3</sub> H <sub>7</sub> —<br>2.42 (10.47)                     |                                      |    | 0.98<br>(10.37)        | —<br>(i-C <sub>3</sub> H <sub>7</sub> O) <sub>2</sub> P(S)SCH <sub>2</sub> COOH<br>1.70, 78.01      | 34.81<br>(35.42)   | 7.02<br>(6.64) | 24.10<br>(23.61) | —<br>(271)                |
| 6     | C <sub>2</sub> H <sub>5</sub> —<br>2.91 (14.33)                       |                                      |    | 1.33<br>(14.07)        | —<br>(C <sub>2</sub> H <sub>5</sub> O) <sub>2</sub> P(S)SCH <sub>2</sub> COOH<br>2.51, 73.34        | 30.01<br>(29.52)   | 4.91<br>(5.33) | 25.79<br>(26.24) | 230<br>(243)              |
| 7     | n-C <sub>3</sub> H <sub>7</sub> —<br>3.51 (15.91)                     |                                      |    | 1.40<br>(14.81)        | —<br>(n-C <sub>3</sub> H <sub>7</sub> O) <sub>2</sub> P(S)SCH <sub>2</sub> COOH<br>3.03, 75.54      | 34.61<br>(35.42)   | 5.69<br>(6.64) | 24.12<br>(23.61) | —<br>(271)                |
| 8     | —CH <sub>2</sub> CH <sub>2</sub> CHMe—<br>1.36 (6.76)                 |                                      |    | 1.1<br>(6.72)          | —<br>OCH <sub>2</sub> CH <sub>2</sub> CHMeOP(S)SCCl <sub>2</sub> COOH<br>1.56, 75.00                | 22.61<br>(23.15)   | 3.10<br>(2.89) | 21.10<br>(20.57) | —<br>(311)                |
| 9     | —CH <sub>2</sub> C(Me) <sub>2</sub> CH <sub>2</sub> —<br>1.30 (6.04)  |                                      |    | 0.98<br>(5.99)         | —<br>OCH <sub>2</sub> C(Me) <sub>2</sub> CH <sub>2</sub> OP(S)SCCl <sub>2</sub> COOH<br>1.64, 85.01 | 26.15<br>(25.84)   | 2.79<br>(3.38) | 20.13<br>(19.69) | 308<br>(325)              |
| 10    | —C(Me) <sub>2</sub> CH <sub>2</sub> CHMe—<br>1.70 (7.42)              |                                      |    | 1.21<br>(7.33)         | —<br>OC(Me) <sub>2</sub> CH <sub>2</sub> CHMeOP(S)SCCl <sub>2</sub> COOH<br>2.23, 89.22             | 27.61<br>(28.31)   | 4.20<br>(3.83) | 19.20<br>(18.87) | —<br>(339)                |
| 11    | —C(Me) <sub>2</sub> C(Me) <sub>2</sub> —<br>1.60 (6.98)               |                                      |    | 1.15<br>(7.03)         | —<br>OC(Me) <sub>2</sub> C(Me) <sub>2</sub> OP(S)SCCl <sub>2</sub> COOH<br>1.90, 79.92              | 27.99<br>(28.31)   | 4.02<br>(3.83) | 19.30<br>(18.87) | 303<br>(339)              |
| 12    | i-C <sub>3</sub> H <sub>7</sub> —<br>1.40 (6.06)                      |                                      |    | 0.99<br>(6.05)         | —<br>(i-C <sub>3</sub> H <sub>7</sub> O) <sub>2</sub> P(S)SCCl <sub>2</sub> COOH<br>2.23, 90.12     | 27.65<br>(28.15)   | 5.11<br>(4.69) | 19.20<br>(18.76) | —<br>(341)                |
| 13    | C <sub>2</sub> H <sub>5</sub> —<br>1.60 (7.88)                        |                                      |    | 1.30<br>(7.95)         | —<br>(C <sub>2</sub> H <sub>5</sub> O) <sub>2</sub> P(S)SCCl <sub>2</sub> COOH<br>1.79, 72.45       | 22.79<br>(23.00)   | 4.02<br>(3.51) | 21.13<br>(20.44) | —<br>(313)                |
| 14    | n-C <sub>3</sub> H <sub>7</sub> —<br>1.77 (7.66)                      |                                      |    | 1.25<br>(7.64)         | —<br>(n-C <sub>3</sub> H <sub>7</sub> O) <sub>2</sub> P(S)SCCl <sub>2</sub> COOH<br>2.08, 80.12     | 27.61<br>(28.15)   | 5.15<br>(4.69) | 19.21<br>(18.76) | 328<br>(341)              |



benzene (50–60 mL) were refluxed for 6–8 h. Ammonium chloride that precipitated was filtered off, and the product was obtained (1.70 g, 78.01%) after evaporating the solvent under reduced pressure; a white low-melting solid was obtained. Relevant data are tabulated in Table III.

### Preparation of Dichloro [(4-Methyl-2-sulfido-1,3,2-dioxaphosphinane-2-yl)thio] Acetic Acid

A benzene solution of trichloro acetic acid (1.21 g, 7.33 mmole) was added into an ammonium salt of hexylene dithiophosphate (1.70 g, 7.42 mmole) and refluxed for 20–22 h. The ammonium chloride precipitated during the course of reaction was filtered off under anhydrous reaction conditions, and the product was isolated after evaporating the solvent under reduced pressure. A pale-yellow oily liquid was obtained (2.23 g, 89.22%). Relevant data are tabulated in Table III.

### REFERENCES

- [1] H. P. S. Chauhan, C. P. Bhasin, G. Srivastava, and R. C. Mehrotra, *Phosphorus, Sulfur, and Silicon*, **15**, 99 (1983).
- [2] K. C. Molloy, M. B. Hossain, D. V. Helim, J. J. Zuckerman, and I. Haidue, *Inorg. Chem.*, **18**, 3507 (1979).
- [3] K. C. Molloy, M. B. Hossain, D. V. Helim, J. J. Zuckerman, and I. Haidue, *Inorg. Chem.*, **19**, 3507 (1980).
- [4] R. C. Mehrotra, G. Srivastava, and B. P. S. Chauhan, *Coord. Chem. Revs.*, **55**, 207 (1984).
- [5] A. Chaturvedi, P. N. Nagar, and G. Srivastava, *Phosphorus, Sulfur, and Silicon*, **80**, 141 (1993).
- [6] P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **79**, 207 (1993).
- [7] A. Chaturvedi, P. N. Nagar, and G. Srivastava, *Main Group Met. Chem.*, **16**, 45 (1993).
- [8] A. Chaturvedi, P. N. Nagar, and A. K. Rai, *Synth. React., Inorg. Met. Chem.*, **26**, 1025 (1996).
- [9] A. Chaturvedi, R. K. Sharma, P. N. Nagar, and A. K. Rai, *Phosphorus, Sulfur, and Silicon*, **112**, 179 (1996).
- [10] R. Purwar and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **86**, 211 (1994).
- [11] R. Purwar, M. K. Sharma, R. K. Sharma, and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **174**, 15 (2001).
- [12] N. Harkut and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **180**, 2517 (2005).
- [13] N. Harkut, A. Keshawat, and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **181**, 2177–2185 (2006).
- [14] C. S. Sharma, M. K. Sharma, and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **177**, 981 (2002).
- [15] C. S. Sharma and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **179**, 1793 (2004).
- [16] C. S. Sharma and P. N. Nagar, *Phosphorus, Sulfur, and Silicon*, **181**, 1 (2006).
- [17] A. I. Vogel, *A Textbook of Quantitative Inorganic Analysis*, 4th Ed. (ELBS, London 1973).