

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Chiral P-Heterocycles: Efficient Method for the Resolution of 3-Methyl-3-phospholene 1-Oxides

Tibor Novak^a; József Schindler^a; Viktória Ujj^b; Mátyás Czugler^c; Elemér Fogassy^b; György Keglevich^b

^a Research Group of the Hungarian Academy of Sciences at the Department of Organic Chemical Technology, Budapest University of Technology and Economics, Budapest, Hungary ^b Department of Organic Chemical Technology, Budapest University of Technology and Economics, Budapest, Hungary ^c Hungarian Academy of Sciences, Chemical Research Center, Budapest, Hungary

To cite this Article Novak, Tibor , Schindler, József , Ujj, Viktória , Czugler, Mátyás , Fogassy, Elemér and Keglevich, György(2008) 'Chiral P-Heterocycles: Efficient Method for the Resolution of 3-Methyl-3-phospholene 1-Oxides', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 183: 2, 543 — 546

To link to this Article: DOI: 10.1080/10426500701764890

URL: <http://dx.doi.org/10.1080/10426500701764890>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Chiral P-Heterocycles: Efficient Method for the Resolution of 3-Methyl-3-phospholene 1-Oxides

Tibor Novak,¹ József Schindler,¹ Viktória Ujj,²
Mátyás Czugler,³ Elemér Fogassy,²
and György Keglevich²

¹Research Group of the Hungarian Academy of Sciences at the
Department of Organic Chemical Technology, Budapest University of
Technology and Economics, Budapest, Hungary

²Department of Organic Chemical Technology, Budapest University of
Technology and Economics, Budapest, Hungary

³Hungarian Academy of Sciences, Chemical Research Center, Budapest,
Hungary

The enantiomers of 1-aryl-, 1-alkyl- and 1-alkoxy-3-methyl-3-phospholene 1-oxides were separated in good yields and in high enantiomeric excesses (up to >99% ee) by resolution via formation of diastereomeric complexes with (–)-(4R,5R)-4,5-bis(diphenylhydroxymethyl)-2,2-dimethyldioxolane (TADDOL) or (–)-(2R,3R)- $\alpha,\alpha,\alpha',\alpha'$ -tetraphenyl-1,4-dioxaspiro[4.5]decan-2,3-dimethanol.

Keywords Chiral P-heterocycles; diastereomeric complex formation; resolution; TADDOL

INTRODUCTION

Phosphine oxides form an important class of phosphorus compounds, since they are precursors of the corresponding phosphines which, in turn, may serve as ligands in transition metal complexes that can be applied in several highly efficient homogenous catalytic processes.^{1,2} Five-membered P-heterocycles, such as 1-substituted-3-phospholene 1-oxides (e.g., **1**) are of synthetic importance, as they can be used as starting materials in the preparation of a variety

The authors are grateful for the financial support from the Hungarian Scientific Research Fund (OTKA, Grants No. T 067679 and T 042725). TN is a recipient of a Bolyai fellowship.

Address correspondence to György Keglevich, Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, H-1521 Budapest, Hungary. E-mail: keglevich@mail.bme.hu




FIGURE 1 Racemic phospholene oxides (**1**) and the resolving agents (**2** and **3**).

of five-, six-, seven-, and eight-membered P-heterocycles including bridged derivatives.^{3–6} Pietrusiewicz and coworkers reported several methods for the preparation of optically pure 1-phenyl-2- or 1-phenyl-3-phospholene 1-oxides and their epoxide derivatives based on dipolar cycloaddition,⁷ enantioselective desymmetrization,^{8–10} and quaternization of their deoxygenated derivatives with a chiral reactant.¹⁰ A simple procedure for the resolution of 1-substituted-3-methyl-3-phospholene 1-oxides (**1**) by separation of the diastereomeric complexes formed by the interaction of (*R*)-**1** or (*S*)-**1** with chiral hosts, such as TADDOL **2** or TADDOL analogue **3**¹¹ is described (Figure 1).¹²


Racemic 1-substituted-3-methyl-3-phospholene 1-oxides (**1**) were resolved by adding half equivalent of (–)-TADDOL (**2**) or its derivatives (**3**) to their hot ethyl acetate solution and precipitating an 1:1 crystalline complexes (+ or –)-**1**·(–)-**2** by the addition of hexane. After two recrystallizations from a mixture of ethyl acetate–hexane and flash chromatography on a silica gel, complexes (+ or –)-**1a–h** were regenerated and analyzed on a chiral HPLC column (Chiralpack AD) or on a chiral GC column (BETA DECTM). The optical purities of (+ or –)-**1a–h** are shown in Table I. As it can be seen, in all cases resolution took place and quite often we could separate the optically pure compound.

The complete resolution procedure is shown for **1a** with TADDOL **2** in Figure 2.

TABLE I Optical Purities of P-Heterocycles (**1a–h**) After Resolution with **2** or **3**


Resolving agent	Optical purity of the products [ee %]							
	1a	1b	1c	1d	1e	1f	1g	1h
2	97	57	69	70	24	95	44	>99
3	>99	>99	>99	>99	58	89*	95	>99

*for the 1:2 complex formed by the use of one equivalent of **3**.



Downloaded At: 17:45 27 January 2011

Downloaded At: 17:45 27 January 2011



Downloaded At: 17:45 27 January 2011

The resolution procedure described above is the first example for the resolution of a cyclic phosphine oxide or a phosphinate via inclusion complex formation. The method comprises crystallization with TADDOL **2** or its analogue **3**, recrystallization of the supramolecular formation and the regeneration of **1**. Our procedure form a simple and convenient method for the preparation of both enantiomers with high enantiomeric purity. The method introduced may be suitable for the resolution of other P-heterocycles or acyclic P-compounds with P = O function, as well.

REFERENCES

- [1] K. M. Pietrusiewicz and M. Zabocka, *Chem. Rev.*, **93**, 1375 (1994).
- [2] R. Noyori, *Asymmetric Catalysis in Organic Synthesis* (John Wiley & Sons, New York, 1994).
- [3] F. Mathey, Ed., *Phosphorus-Carbon Heterocyclic Chemistry: The Rise of a New Domain* (Pergamon/Elsevier, Amsterdam, 2001).
- [4] Gy. Keglevich, *Synthesis*, 931 (1993).
- [5] Gy. Keglevich, *Rev. Heteroatom Chem.*, **14**, 119 (1996).
- [6] Gy. Keglevich, *Curr. Org. Chem.*, **10**, 93 (2006).
- [7] K. M. Pietrusiewicz, W. Hoody, M. Koprowski, S. Cicchi, A. Goti, and A. Brandi, *Phosphorus, Sulfur and Silicon*, **144–146**, 389 (1999).
- [8] K. M. Pietrusiewicz, M. Koprowski, Z. Pakulski, *Tetrahedron: Asymmetry*, **13**, 1017 (2002).
- [9] Z. Pakulski and K. M. Pietrusiewicz, *Tetrahedron: Asymmetry*, **15**, 41 (2004).
- [10] K. M. Pietrusiewicz, Phosphorus, Sulfur and Silicon, **109–110**, 573 (1996).
- [11] A. K. Beck, B. Bastani, D. A. Plattner, W. Petter, D. Seebach, H. Braunschweiler, P. Gysi, and L. L. Vecchia, *Chimia*, **45**, 238 (1991).
- [12] T. Novak, J. Schindler, V. Ujj, M. Czugler, E. Fogassy, and Gy. Keglevich, *Tetrahedron: Asymmetry*, **17**, 2599 (2006).