

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

A Convenient One-Pot Synthesis of Some New 3-(2-Phenyl-6-(2-thienyl)-4-pyridyl)hydroquinolin-2-ones Under Microwave Irradiation and Their Antimicrobial Activities

Niraj K. Ladani^a; Manish P. Patel^a; Ranjan G. Patel^a

^a Department of Chemistry, S.P. University, Vallabh Vidyanagar, Gujarat, India

Online publication date: 24 February 2010

To cite this Article Ladani, Niraj K. , Patel, Manish P. and Patel, Ranjan G.(2010) 'A Convenient One-Pot Synthesis of Some New 3-(2-Phenyl-6-(2-thienyl)-4-pyridyl)hydroquinolin-2-ones Under Microwave Irradiation and Their Antimicrobial Activities', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185: 3, 658 – 662

To link to this Article: DOI: 10.1080/10426500902915507

URL: <http://dx.doi.org/10.1080/10426500902915507>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A CONVENIENT ONE-POT SYNTHESIS OF SOME NEW 3-(2-PHENYL-6-(2-THIENYL)-4-PYRIDYL) HYDROQUINOLIN-2-ONES UNDER MICROWAVE IRRADIATION AND THEIR ANTIMICROBIAL ACTIVITIES

Niraj K. Ladani, Manish P. Patel, and Ranjan G. Patel

Department of Chemistry, S.P. University, Vallabh Vidyanagar, Gujarat, India

A series of 3-(2-phenyl-6-(2-thienyl)-4-pyridyl)hydroquinolin-2-ones 4a–o were synthesized in high yields by a one-pot cyclocondensation reaction under Kröhnke's reaction conditions using 2-chloro-3-formyl quinoline 1a–c, 2-acetyl thiophene 2, and various N-phenacylpyridinium bromides 3a–e in a mixture of ammonium acetate and acetic acid by microwave irradiation. All the compounds have been characterized by elemental analysis, FT-IR, ¹H NMR, and ¹³C NMR spectral analysis. These compounds have been screened for their antimicrobial activities.

Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.

Keywords 2-Acetyl thiophene; anti-microbial activity; 2-chloro-3-formyl quinoline; Kröhnke reaction condition; microwave irradiation; various N-phenacylpyridinium bromides

INTRODUCTION

The polysubstituted pyridines are prominent building blocks in supramolecular chemistry with their π -stacking and directional H-bonding ability.^{1,2} The pyridyl heterocyclic core is also a widespread subunit in numerous natural products and pharmaceuticals.^{3,4} These facts provide the basis for the development of new efficient synthetic pathways for pyridines. Polysubstituted pyridines have been synthesized using an enormous number of preparative approaches such as [5+1]-type Hantzsch synthesis from 1,5-diketone and nitrogen derivatives,^{5,6} [3+3]-type cyclization of chalcones and iminophosphoranes,⁷ [4+2] reactions of unsaturated imines with enolates,⁸ and [3+2+1]-type cyclization of α,β -unsaturated compounds with α -substituted ketones and a nitrogen source.⁹ Among these approaches, the latter approach is the most frequently employed. The two-step Kröhnke synthesis^{10–12} via condensation of α,β -unsaturated ketones with pyridinium salts in the presence of a mixture of ammonium acetate and acetic acid gives a variety of polysubstituted pyridines and has distinct advantages over the other routes. The drawback of this

Received 28 November 2008; accepted 23 March 2009.

The authors are thankful to the Department of Chemistry, S.P. University, for providing research facilities and Vaibhav laboratory, Ahmedabad, for providing IR facilities.

Address correspondence to Ranjan G. Patel, Department of Chemistry, S.P. University, Vallabh Vidyanagar-388 120, Gujarat, India. E-mail: patelranjanben@yahoo.com


method is that pyridinium salts and α,β -unsaturated ketones have to be prepared first. To overcome this problem Chao-Guo Yan and his coworkers¹³ have described a simple but effective modification of the Kröhnke's synthesis of pyridines in one-pot reactions of *N*-phenacylpyridinium bromide with aromatic aldehydes and acetophenone in the presence of ammonium acetate and acetic acid under microwave irradiation to give polysubstituted pyridine derivatives. A literature survey reveals that a number of pyridine^{14,15} derivatives have been synthesized by Kröhnke reaction conditions using various aldehydes, but not a single reference has been found where 2-chloro-3-formyl quinoline is used. We report in this article biologically active new pyridine derivatives containing quinoline moiety by Kröhnke reaction conditions utilizing this heterocyclic aldehyde^{16–18} by microwave irradiation.

Recently, the synthesis of organic compounds assisted by microwaves^{19,20} has become an improved technique due to greater selectivity, rapid transfer of energy, significant practical simplicity, and pure products. Compared with classical heating, microwave-assisted organic synthesis is characterized by spectacular acceleration, higher yields, milder reaction conditions, and shorter reaction times as well as allowing synthesis to become environmentally benign, improving many processes.^{21,22} The coupling of microwave technology with one-pot multicomponent condensation reactions has received significant attention,²³ since two or more steps in the synthetic sequence can be carried out without the isolation of intermediates. This leads to reduction of time and energy and constitutes overall an economical way of developing new pharmaceutically important compounds.²⁴

RESULTS AND DISCUSSION

Under microwave irradiation, 2-chloro-3-formyl quinoline **1a–c**, 2-acetyl thiophene **2**, and *N*-phenacylpyridinium bromide **3a–e** in the presence of ammonium acetate and acetic acid reacted smoothly to give **4a–o** in high yield (78–90%). The required 2-chloro-3-formyl quinoline **1a–c** was prepared by Vilsmeier–Haack reaction,²⁵ and *N*-phenacylpyridinium bromide **3a–e** was prepared by the procedure in the literature.²⁶

The formation of compounds **4a–o** involves the Kröhnke mechanism (see Scheme S1, available online in the Supplemental Materials). The reaction is homogeneous and proceeds via 2-chloro-3-formyl quinoline **1a–c**, reacted first with 2-acetyl thiophene **2** to form an α,β -unsaturated ketone (**4I**). **4I** is further reacted with *N*-phenacylpyridinium bromide **3a–e** to give 1,5-diketone derivatives (**4II**). The cyclization of **4II** occurs with ammonia, and finally the pyridine cation is eliminated with the formation of 3-(2-phenyl-6-(2-thienyl)-4-pyridyl) hydroquinolin-2-ones^{27,28} **4a–o**. During the course of the reaction, 2-chloro quinoline is converted into 2-quinolones,²⁹ The structures of compounds **4a–o** were confirmed by IR, ¹H NMR, and ¹³C NMR spectra. The IR spectra of **4a** exhibited absorptions at 3180 for –NH–, 3010 for aromatic C–H stretching, 1655 for (carbonyl group), and 1420–1600 cm^{–1} for C=C aromatic and C=N stretching of pyridine. The ¹H NMR of **4a** showed a singlet at δ 12.12 for –NH– proton and aromatic protons resonate as multiplets at 7.21–8.54. The ¹³C NMR spectrum of **4a** showed signals at δ 128.89 (–CH–C–S–), 131.65 (–CO–C–), 138.84 (–NH–C–), 140.15 (–N=C–C–), 145.32 (=N–C–C–S–), 146.40 (–CO–C–C=), 152.10 (–C–N–), 156.03 (–C=N–), 115.43–139.42 for aromatic carbon, and the carbonyl carbon was observed at 161.10. The structure was further confirmed by its mass spectral studies. It gave a molecular ion peak at *m/z* 381 (M+1) corresponding to the molecular formula C₂₄H₁₆N₂OS (Scheme 1). All the compounds were screened for their antibacterial and antifungal activities using ciprofloxacin, ampicillin, and griseofulvin as standard drugs.



Scheme 1

EVALUATION OF ANTIMICROBIAL ACTIVITY

The in vitro antimicrobial activity of compounds **4a–o** was carried out against 24-h old cultures of three bacteria: *Escherichia coli* as Gram-negative bacteria and *Bacillus subtilis* and *Staphylococcus aureus* as Gram-positive bacteria, and two fungi: *Aspergillus niger* and *Rhizopus oryzae* by the disc-diffusion method.^{30,31} (For full details regarding antimicrobial activity, see the Supplemental Materials online and Table S1.)

EXPERIMENTAL

All melting points were taken in open capillaries and are uncorrected. The monitoring of the progress of all reactions and homogeneity of the synthesized compounds was carried out by TLC. TLC was run using TLC aluminum sheets silica gel 60 F₂₅₄ (Merck). Elemental analysis (% C, H, N) was carried out by a Perkin Elmer 2400 CHN elemental analyzer. IR spectra were recorded on a Shimadzu FTIR 8401 spectrophotometer using KBr pellets. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker Avance 400 MHz spectrometer using solvent peak as the internal standard. Mass spectra were scanned on a Shimadzu LCMS 2010 spectrometer. The microwave oven (700 W) used was a modified microwave oven (RAGA's Electromagnetic Systems).

Preparation of 3-(2-Phenyl-6-(2-thienyl)-4-pyridyl)hydroquinolin-2-ones **4a–o**

2-Chloro-3-formyl quinoline (0.5 g, 0.0026 mol), 2-acetyl thiophene (0.28 mL, 0.0026 mol), *N*-phenacylpyridinium bromide (0.72 g, 0.0026 mol), ammonium acetate (2.0 g, 0.026 mol), and acetic acid (5 mL) were charged in a 25 mL round bottom flask.

Table I Substitutes on compounds 4a–o

Compound	R	R ₁
4a	H	H
4b	CH ₃	H
4c	OCH ₃	H
4d	H	CH ₃
4e	CH ₃	CH ₃
4f	OCH ₃	CH ₃
4g	H	OCH ₃
4h	CH ₃	OCH ₃
4i	OCH ₃	OCH ₃
4j	H	Cl
4k	CH ₃	Cl
4l	OCH ₃	Cl
4m	H	F
4n	CH ₃	F
4o	OCH ₃	F

The flask was heated for 4–6 min at 118°C under microwave irradiation (at 420 W). The reaction was monitored by TLC; after the completion of reaction, the reaction mixture was cooled to room temperature and stirred for 30 min, and the resulting solid was collected by filtration and washed with water. The crude product was purified by leaching in an equimolar mixture of chloroform and methanol (20 mL) to obtain the pure solid sample 4a–o. A sample characterization is given, and the remainder of the data are found in the Supplemental Materials online.

3-(2-Phenyl-6-(2-thienyl)-4-pyridyl)hydroquinolin-2-one 4a

A pale yellow solid, mp 295–296°C, yield: (90%). Anal. Calcd. for C₂₄H₁₆N₂OS (380.47): C, 75.76; H, 4.23; N, 7.36. Found: C, 75.61; H, 4.36; N, 7.54%. IR (KBr): 3180 (N–H stretching), 3010 (aromatic C–H stretching), 1655 (>C=O stretching), 1420–1600 (C=C aromatic and C=N stretching of pyridine), 690 (C–S stretching) cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 7.21–8.54 (15H, m, Ar–H), 12.12 (1H, s, NH) ppm; ¹³C NMR: δ 115.43, 117.20, 118.62, 119.71, 122.65, 125.91, 127.13, 128.96, 129.07, 129.21, 129.27, 129.75, 139.42 (–C–Ar–), 128.89 (–CH–C–S–), 131.65 (–CO–C–), 138.84 (–NH–C–), 140.15 (–N=C–C–), 145.32 (=N–C–C–S–), 146.40 (–CO–C–C=), 152.10 (–C–N–), 156.03 (–C=N–), 161.10 (–CO–NH–) ppm; Mass: *m/z*: 381 (M+1).

CONCLUSION

A simple and efficient, one-pot procedure has been followed for the generation of polysubstituted pyridines with microwave assistance. The method allows the introduction of various substituted alkyl and aryl groups into the 2-, 4-, and 6-positions of pyridine. It can be concluded from Table I that among all the compounds, 4e and 4m showed good antibacterial activity against the tested organisms *E. coli* and *S. aureus*, while compounds 4d and 4m showed good antibacterial activity against the tested organism *B. subtilis*. The compounds 4h, 4l, and 4m against the tested organism *R. oryzae* and compounds 4d and 4n against the tested organism *A. niger* showed good antifungal activity, while the remaining

compounds of the series were moderately active against the tested organisms *E. coli*, *S. aureus*, *B. subtilis*, *A. niger*, and *R. oryzae*.

REFERENCES

1. E. C. Constable, R. M. Maney, A. M. W. C. Thompson, and J. V. Walker, *J. Chem. Soc., Dalton Trans.*, 1585 (1994).
2. I. Eryazici, C. N. Moorefield, S. Drumus, and G. R. Newkome, *J. Org. Chem.*, **71**, 1009 (2006).
3. D. Barton and D. Ollis, *Comprehensive Organic Chemistry* (Pergamon, New York, 1979), vol. 4, p. 468.
4. A. R. Katritzky and C. M. Marson, *Angew. Chem. Int. Ed. Engl.*, **23**, 420 (1984).
5. G. Jones, A. R. Katritzky, and C. W. Rees, *Comprehensive Heterocyclic Chemistry* (Pergamon, New York, 1984), vol. 2, Part 2A, p. 395.
6. A. Sausins and D. G. Durburs, *Heterocycles*, **27**, 291 (1988).
7. T. Kobayashi and M. Nita, *Chem. Lett.*, **15**, 1549 (1986).
8. A. S. Kiselyov, *Tetrahedron Lett.*, **36**, 9297 (1995).
9. M. C. G. Barrio, J. R. Barrio, G. Walker, A. Novelli, and N. J. Leonard, *J. Am. Chem. Soc.*, **95**, 4891 (1973).
10. F. Krohnke, *Angew. Chem. Int. Ed. Engl.*, **2**(5), 225 (1963).
11. F. Krohnke and W. Zecher, *Angew. Chem. Int. Ed. Engl.*, **1**(12), 626 (1962).
12. (a) F. Krohnke, *Synthesis*, 1 (1976); (b) F. Neve, A. Crispini, and S. Campagna, *Inorg. Chem.*, **36**, 6150 (1997); (c) L. R. Macgillivray, P. R. Diamente, J. L. Reid, and J. A. Ipmeester, *Chem. Commun.*, 359 (2000).
13. C. G. Yan, X. M. Cai, Q. F. Wang, T. Y. Wang, and M. Zheng, *Org. Biomol. Chem.*, **5**, 945 (2006).
14. X. M. Cai, Q. F. Wang, and C. G. Yan, *Synth. Commun.*, **37**, 223 (2007).
15. V. V. Mulwad and D. S. Satwe, *Indian J. Chem.*, **43B**, 2727 (2004).
16. B. Kalluraya, J. Nayak, A. Adhikari, K. V. Sujith, N. S. Shetty, and M. Winter, *Phosphorus, Sulfur, and Silicon*, **183**, 1870 (2008).
17. K. Srinivasu, M. Sunilkumar, and A. V. Reddy, *Indian J. Chem.*, **38B**, 553 (1999).
18. R. N. Kumar, T. Suresh, and P. S. Mohan, *Indian J. Chem.*, **42B**, 688 (2003).
19. L. Perreux and A. Loupy, *Tetrahedron*, **57**, 9199 (2001).
20. A. R. Desai and K. R. Desai, *Arkivoc*, **xiii**, 98 (2005).
21. S. S. Laddha and S. P. Bhatnagar, *Phosphorus, Sulfur, and Silicon*, **183**, 2262 (2008).
22. A. D. Ortiz, F. Lang, A. D. L. Hoz, and A. Moreno, *Eur. J. Org. Chem.*, **4**, 3659 (2000).
23. J. H. Clark and D. J. Macquarrie, *Chem. Soc. Rev.*, **25**, 303 (1996).
24. C. S. Miller and C. A. Wagner, *J. Org. Chem.*, **13**, 89 (1948).
25. O. M. Cohn, B. Narine, and B. Tarnowski, *J. Chem. Soc. Perkin Trans I*, 1520 (1981).
26. R. K. Bansal and B. Gope, *Indian J. Chem.*, **18B**, 362 (1979).
27. S. Tu, R. Jia, B. Jiang, J. Zhang, Y. Zhang, C. Yao, and S. Ji, *Tetrahedron*, **63**, 381 (2007).
28. U. Siemeling, J. V. D. Bruggen, U. Vorfeld, A. Stammeler, and H. G. Stammeler, *Z. Naturforsch.*, **58b**, 443 (2003).
29. A. Srivastava and R. M. Singh, *Indian J. Chem.*, **44B**, 1868 (2005).
30. C. H. Collins and P. M. Lyne, *Microbial Methods* (University Park Press, Baltimore, 1970).
31. T. D. Thangadurai and K. Natarajan, *Indian J. Chem.*, **40A**, 573 (2001).