

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

An Efficient Synthesis of New Substituted Spiro Pyrazolethieno, Pyrimidinethieno, and Benzodiazepinethieno Pyridazine Derivatives with Biological Activities

Rasha A. M. Faty^a; Hoda A. R. Hussein^b; Ayman M. S. Youssef^c

^a Cairo University, Faculty of Science, Giza, Egypt ^b Department of Photochemistry (Heterocyclic Unit), National Research Centre, Dokki, Egypt ^c Fayoum University, Faculty of Science, Fayoum, Egypt

Online publication date: 03 July 2010

To cite this Article Faty, Rasha A. M. , Hussein, Hoda A. R. and Youssef, Ayman M. S.(2010) 'An Efficient Synthesis of New Substituted Spiro Pyrazolethieno, Pyrimidinethieno, and Benzodiazepinethieno Pyridazine Derivatives with Biological Activities', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 185: 7, 1484 – 1490

To link to this Article: DOI: 10.1080/10426500903095531

URL: <http://dx.doi.org/10.1080/10426500903095531>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN EFFICIENT SYNTHESIS OF NEW SUBSTITUTED SPIRO PYRAZOLETHIENO, PYRIMIDINETHIENO, AND BENZODIAZEPINETHIENO PYRIDAZINE DERIVATIVES WITH BIOLOGICAL ACTIVITIES

Rasha A. M. Faty,¹ Hoda A. R. Hussein,²
and Ayman M. S. Youssef³

¹Cairo University, Faculty of Science, Giza, Egypt

²Department of Photochemistry (Heterocyclic Unit), National Research Centre, Dokki, Egypt

³Fayoum University, Faculty of Science, Fayoum, Egypt

3-Ethyl 2-amino-4-methyl-5-phenyl thiophene carboxylate 1 was used as a starting material to synthesize 2a,b via coupling with malononitrile or acetyl acetone, respectively. When heated, under reflux in sodium ethoxide solution, 2a,b give 3a,b. On the other hand, when compounds 3a,b were heated under reflux in ethanol with hydrazine hydrate, thiourea, or 1,1-phenylenediamine hydrochloride and a catalytic amount of piperidine 4a,b, 5a,b and 6a,b, were produced, respectively. The new compounds were tested for their antimicrobial activity. Compounds 2a–6b showed antibacterial activities, and 2a,2b and 4b showed antifungal activities.

Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.

Keywords Antibacterial; antifungal; benzodiazepinethieno-pyridazine; biological activity; pyrimidinethieno; spiro pyrazolethieno

INTRODUCTION

From a chemical structure point of view, various natural products or synthetic molecules containing the rigid conformations of spiro cyclic skeletons show a wide range of biological and pharmacological properties.^{1–3} For example, spirohydantoins have been reported as glycogen and phosphorylase inhibitors, herbicides, and anti-inflammatory agents,⁴ and spiro-isoxazolines and spiro-dioxazoles exhibit antitumor and anti-HIV activities.^{5–8} In spite of the immense biological activities of many spiro heterocyclic compounds, no report is yet available on the synthesis of pyrazolothienopyridazine spiro derivatives. Prompted by this observation and in continuation of our earlier interest on chemical synthesis of biodynamic heterocyclic compounds,⁹ we have made an attempt to synthesize novel tricyclic spiro derivatives.

Received 7 January 2009; accepted 3 June 2009.

Address correspondence to Rasha A. M. Faty, Cairo University, Faculty of Science, Giza, Egypt. E-mail: rashafaty@yahoo.com

RESULTS AND DISCUSSION

Chemistry

For synthesis of novel spiro derivatives, at first, we attempted to obtain dinitrile or diacetyl derivatives by oxidative cyclization of diazo thieno derivatives. Thus, by using 3-ethyl 2-amino-4-methyl-5-phenylthiophene carboxylate¹⁰ **1** as a starting material, it was coupled with some active methylene groups namely, malononitrile or acetylacetone, respectively, to obtain 3-Ethoxycarbonyl-4-methyl-5-phenyl(thien-2-ylazo)malononitrile **2a** or 3-ethoxycarbonyl-4-methyl-5-phenyl(thien-2-ylazo)acetylacetone **2b**. Elemental analyses as well as spectral data of compounds **2a,b** are in agreement with the proposed structures (see the Experimental section).

Heating **2a,b** under reflux with ethanolic sodium ethoxide solution for 3 h yielded 3,3-disubstituted thienopyridazine derivatives **3a,b**. Compounds **3a,b** give expected values in elemental analyses and spectral data. The IR spectrum (KBr) cm^{-1} of **3b**, for example, showed absorption bands at: 1719 (2CO), 1673 (CO); ^1H NMR (DMSO- d_6) δ ppm showed signals at: 2.31 (s, 3H, CH_3), 3.80 (s, 6H, 2CH_3), 7.43–7.46 (m, 5H, aromatic protons); MS (m/z): 326 (M^+) 80%.

Compounds **3a,b** underwent oxidative cyclization upon heating under reflux in ethanol with hydrazine hydrate in the presence of a catalytic amount of piperidine to produce 3,5-diamino(diacetyl)-5'-methyl-6'-phenyl-4'H-spiro-(pyrazole-4,3'-thieno[2,3-c]pyridazine)-4'-one **4a,b** with new ring systems (Scheme 1). Compounds **4a,b** gave an expected elemental analyses (see the Experimental section). The IR spectrum (KBr) cm^{-1} of **4a**, for example, showed: 3313, 3180(NH_2), and 1660(CO); ^1H NMR (DMSO- d_6) δ ppm: 2.22 (s, 3H, CH_3), 3.81 (br., 2H, NH_2 , D_2O exchangeable), 7.40–7.57 (m, 5H, aromatic protons), and 9.44 (br., 2H, NH_2 , D_2O exchangeable); ^{13}C NMR (75 MHz, CDCl_3) of **4a** for example showed signals at δ 189 (S) ppm, 160 (S, 2C), 145 (S), 132 (S), 130 (S, 3C), 129 (d, 2C), 127 (d, 2C), 126 (d), 79 (S), 5 (q).

On the other hand, compounds **3a,b** also reacted with thiourea in boiling ethanol containing a catalytic amount of piperidine to produce other novel spiro- compounds 4,6-diamino(diacetyl)-5'-methyl-6'-phenyl-2-thioxo-2H,4'H-spiro-[pyrimidine-5,3'-thieno[2,3-c]pyridazin]-4'-one **5a,b**, which showed expected spectral data and elemental analyses (see the Experimental section and Scheme 1).


Also, heating **3a,b** under reflux with o-phenylenediamine in boiling ethanol containing a catalytic amount of piperidine produced other novel spiro compounds 2,4-diamino(diacetyl)-5'-methyl-6'-phenyl-4'H-spiro[1,5-benzodiazepine-3,3'-thieno[2,3-c]pyridazin]-4'-one **6a,b**. Besides the expected values of elemental analyses, the IR, ^1H NMR, and ^{13}C NMR spectra of **6a,b** are in agreement with the assigned structure (see the Experimental section and Scheme 1).

Biology

All new compounds were evaluated for their antimicrobial properties. Compound **2a** showed inhibition zones and therefore antibacterial activity against *staphylococcus aureus* (G^+) more than the reference compound tetracycline antibacterial agent. (See the Supplemental Materials available online, Table S1.)

CONCLUSION

This work is concerned with the synthesis and reactions of 3-ethyl-2-amino-4-methyl-5-phenyl thiophene carboxylate with functional and bifunctional groups leading to spiro



Downloaded At: 16:31 27 January 2011

Downloaded At: 16:31 27 January 2011

Downloaded At: 16:31 27 January 2011

Downloaded At: 16:31 27 January 2011

Table I Physical data for the products **2a–6b**

Compound No.	Mp °C	Yield %	Molecular formula (M. wt.)	Elemental analyses (calcd./found)		
				%C	%H	%N
2a	151–153	80	C ₁₇ H ₁₄ N ₄ O ₂ S	60.34	4.17	16.55
			338.39	60.30	4.00	16.50
2b	170–172	75	C ₁₉ H ₂₀ N ₂ O ₄ S	61.27	5.41	7.52
			372.44	61.22	5.40	7.50
3a	212–214	80	C ₁₅ H ₈ N ₄ OS	61.63	2.75	19.16
			292.32	61.60	2.74	19.12
3b	180–182	75	C ₁₇ H ₁₄ N ₂ O ₃ S	62.56	4.32	8.58
			326.37	62.54	4.30	8.55
4a	190–192	73	C ₁₅ H ₁₂ N ₆ OS	55.54	3.72	25.90
			324.36	55.50	3.70	25.90
4b	200–202	70	C ₁₇ H ₁₄ N ₄ OS	63.33	4.37	17.37
			322.39	63.30	4.35	17.35
5a	195–197	75	C ₁₆ H ₁₂ N ₆ OS ₂	52.15	3.28	22.80
			368.44	52.10	3.25	22.80
5b	205–207	65	C ₁₈ H ₁₄ N ₄ OS ₂	58.99	3.85	15.28
			366.47	58.97	3.85	15.26
6a	185–187	75	C ₂₁ H ₁₆ N ₆ OS	62.98	4.02	20.98
			400.46	62.97	4.00	20.97
6b	202–204	60	C ₂₃ H ₁₈ N ₄ OS	69.32	4.55	14.06
			398.49	69.32	4.55	14.00

Faculty of Science, Cairo University, Giza, Egypt. All solid compounds were recrystallized to produce constant melting points. See Table I.

3-Ethyl-2-amino-4-methyl-5-phenyl thiophene carboxylate (**1**)

Compound **1** was prepared according to the Gewald method.¹⁰

General Method for Preparation of **2a,b**

To an ice-cold solution of the appropriate amine **1** (0.01 mol), glacial acetic acid (30 mL), and phosphoric acid (10 mL), a solution of sodium nitrite (1.03 g, 0.01 mol) dissolved in the minimum amount of water was added dropwise, in an ice bath at -5°C . This previously prepared diazonium salt was added dropwise to a mixture of active methylene malononitrile or acetyl acetone (0.01 mol) and anhydrous sodium acetate in ethanol. The reaction mixture was allowed to stand overnight at room temperature, then it was poured into water. The formed solid was filtered off, washed with water, dried, and recrystallized from the appropriate solvent to produce **2a,b**.

3-Ethoxycarbonyl-4-methyl-5-phenyl(thien-2-ylazo)malononitrile (**2a**).

Compound **2a** was obtained by the reaction mixture of **1** (2.60 g, 0.01 mol) and malononitrile (0.66 g, 0.01 mol). The compound was recrystallized from ethanol to produce **2a** as deep orange crystals; IR spectrum (KBr) cm^{-1} : 3318 (br, NH), 2232, 2211 (2CN), and 1668 (CO); ^1H NMR (DMSO- d_6) δ ppm (J, Hz): 1.29 (t, 3H, CH₃, J = 4.2), 2.24(s, 3H,

CH₃), 3.12 (s, 1H, NH, D₂O exchangeable), 4.26 (q, 2H, CH₂, J = 4.2) and 7.34–7.47 (m, 5H, aromatic protons); MS (m/z): 338 (M⁺) 80%.

3-Ethoxycarbonyl-4-methyl-5-phenyl(thien-2-ylazo)acetylacetone (2b).

Compound **2b** was obtained by the reaction mixture of **1** (2.60 g, 0.01 mol) and acetyl acetone (1.00 g, 0.01 mol). The compound was recrystallized from ethanol to produce **2b** as orange crystals; IR spectrum (KBr) cm⁻¹: 3324(br, NH), 1671, 1645 (3CO); ¹H NMR (DMSO-d₆) δ ppm (J, Hz): 1.31 (t, 3H, CH₃, J = 4.7), 2.21 (s, 3H, CH₃), 4.00 (s, 6H, 2CH₃), 4.10 (s, 1H, NH, D₂O exchangeable), 4.40 (q, 2H, CH₂, J = 4.7) and 7.34–7.45 (m, 5H, aromatic protons); MS(m/z): 372(M⁺) 100%.

3,3-Dicyano(diacetyl)-5-methyl-6-phenylthieno[2,3-c]pyridazine-4-one (3a,b)

Compound **2a,b** (0.01 mol) was refluxed in sodium ethoxide solution [(prepared by dissolving sodium metal (0.23 g, 0.01 mol) in absolute ethanol (30 mL)] was heated under reflux for 5 h. The reaction mixture was allowed to cool to room temperature, poured into water, and neutralized by dilute acetic acid solution. The solid product that precipitated was filtered off, dried, and recrystallized from the proper solvent to produce **3a,b**.

3,3-Dicyano -5-methyl-6-phenylthieno[2,3-c]pyridazine-4-one (3a). Compound **3a** was obtained by refluxing **2a** (3.38 g, 0.01 mol) in sodium ethoxide solution. The solid product was recrystallized from dioxane to produce **3a** as deep orange crystals; IR spectrum (KBr) cm⁻¹: 2219 (2CN) and 1708 (CO); ¹H NMR (DMSO-d₆) δ ppm: 2.33 (s, 3H, CH₃), 7.34–7.47 (m, 5H, aromatic protons); ¹³C NMR (75 MHz, CDCl₃): δ 190 (s) ppm, 148 (s), 135 (s), 134 (s, 3C), 129 (d, 3C), 128 (d, 2C), 117(s, 2C), 58 (s), 5 (q); MS (m/z): 292(M⁺) 70%.

3,3-Diacetyl-5-methyl-6-phenylthieno[2,3-c]pyridazin-4-one (3b). Compound **3b** was obtained by refluxing **2b** (3.72 g, 0.01 mol) in sodium ethoxide solution. The solid product was recrystallized from dioxane to produce **3b** as orange crystals; IR spectrum (KBr) cm⁻¹: 1719 (2CO), 1673 (CO); ¹H NMR (DMSO-d₆) δ ppm: 2.31(s, 3H, CH₃), 3.80 (s, 6H, 2CH₃), 7.43–7.46 (m, 5H, aromatic protons); ¹³C NMR (75 MHz, CDCl₃): δ 204 (s, 2C) ppm, 192 (s), 147 (s), 135 (s), 133 (s, 3C), 129 (d, 3C), 128 (d, 2C), 103 (s), 24 (q, 2C), 5 (q); MS(m/z): 326 (M⁺) 80%.

3,5-Diamino(diacetyl)-5'-methyl-6'-phenyl-4'H-spiro[pyrazole-4,3'-thieno[2,3-c]pyridazin]-4'-one (4a,b)

A mixture of **3a,b** (0.01 mol) and hydrazine hydrate (99–100%) (7 mL, 0.03 mol) was refluxed in ethanol (30mol) containing a catalytic amount of piperidine for 5 h. The reaction mixture was cooled, filtered off, and recrystallized from the proper solvent to produce **4a,b**.

3,5-Diamino-5'-methyl-6'-phenyl-4'H-spiro[pyrazole-4,3'-thieno[2,3-c]pyridazin]-4'-one (4a). Compound **4a** was obtained by refluxing **3a** (2.92 g, 0.01 mol) and hydrazine hydrate in ethanol containing a catalytic amount of piperidine. The solid product recrystallized from dioxane to produce **4a** as deep red crystals; IR spectrum (KBr) cm⁻¹: 3313, 3180 (NH₂) and 1660 (CO); ¹H NMR (DMSO-d₆) δ ppm: 2.22 (s, 3H, CH₃), 3.81 (br.,2H, NH₂, D₂O exchangeable), 7.40–7.57 (m, 5H, aromatic protons) and 9.44 (br.,2H, NH₂, D₂O exchangeable); MS (m/z): 324 (M⁺) 14.80%.

3,5-Diacetyl-5'-methyl-6'-phenyl-4'H-spiro[pyrazole-4,3'-thieno[2,3-c]pyridazin]-4'-one (4b). Compound **4b** was obtained by refluxing **3b** (3.26 g, 0.01 mol) and hydrazine hydrate in ethanol containing a catalytic amount of piperidine. The solid product recrystallized from dioxane to produce **4b** as deep red crystals; IR spectrum (KBr) cm^{-1} : 1670 (CO); ^1H NMR (DMSO-d_6) δ ppm: 2.23 (s, 6H, 2CH_3), 3.72 (s, 3H, CH_3) and 7.32–7.44 (m, 5H, aromatic protons); ^{13}C NMR (75 MHz, CDCl_3): δ 190 (s) ppm, 165 (s, 2C), 147 (s), 136 (s), 134 (s, 3C), 129 (d, 3C), 128(d, 2C), 80 (s), 19 (q, 2C), 5 (s); MS (m/z): 322 (M^+) 65%.

4,6-Diamino(diacetyl)-5'-methyl-6'-phenyl-2-thioxo-2H-4'H-spiro[pyrimidine-5,3'-thieno[2,3-c]pyridazin]-4'-one (5a,b)

A mixture of **3a,b** (0.01 mol) and thiourea (0.01, mol) was refluxed in ethanol containing a catalytic amount of piperidine for 5 h. The reaction mixture was cooled and poured into water. The solid product that precipitated was filtered off, dried, and recrystallized from the proper solvent to produce **5a,b**.

4,6-Diamino-5'-methyl-6'-phenyl-2-thioxo-2H-4'H-spiro[pyrimidine-5,3'-thieno[2,3-c]pyridazin]-4'-one (5a). Compound **5a** was obtained by refluxing of **3a** (2.92 g, 0.01 mol) and thiourea (0.76 g, 0.015 mol). The solid product recrystallized from dioxane to produce **5a** as yellow crystals; IR spectrum (KBr) cm^{-1} : 3310, 3150 (NH_2) and 1654 (CO); ^1H NMR (DMSO-d_6) δ ppm: 2.33 (s, 3H, CH_3), 3.88 (br., 2H, NH_2 , D_2O exchangeable), 7.22–7.47 (m, 5H, aromatic protons) and 9.44 (br., 2H, NH_2 , D_2O exchangeable); ^{13}C NMR (75 MHz, CDCl_3): δ 229(s) ppm, 191 (s), 165 (s, 2C), 147 (s), 135 (s), 134 (s, 3C), 129 (d, 3C), 128 (d, 2C), 72 (s), 5 (q); MS (m/z): 368(M^+) 96%.

4,6-Diacetyl-5'-methyl-6'-phenyl-2-thioxo-2H-4'H-spiro[pyrimidine-5,3'-thieno[2,3-c]pyridazin]-4'-one (5b). Compound **5b** was obtained by refluxing **3b** (3.26g, 0.01 mol) and thiourea (0.76 g, 0.015 mol). The solid product recrystallized from dioxane to produce **5b** as orange crystals; IR spectrum (KBr) cm^{-1} : 1673 (CO); ^1H NMR (DMSO-d_6) δ ppm: 2.21 (s, 3H, CH_3), 3.20 (s, 6H, 2CH_3) and 7.22–7.48 (m, 5H, aromatic protons); ^{13}C NMR (75 MHz, CDCl_3): δ 229(s) ppm, 192 (s), 165 (s, 2C), 147 (s), 135 (s), 134 (s, 3C), 129 (d, 3C), 128 (d, 2C), 72 (s), 15 (q, 2c), 5 (q); MS (m/z): 366(M^+) 70%.

2,4-Diamino(diacetyl)-5'-methyl-6'-phenyl-4'H-spiro[1,5-benzodiazepine-3,3'-thieno[2,3-c]pyridazin]-4'-one (6a,b)

A mixture of **6a,b** (0.01 mol) and *o*-phenylenediamine hydrochloride (0.01 mol) was refluxed in boiling ethanol containing a catalytic amount of piperidine for 5 h. The reaction mixture was cooled and poured into water. The formed solid was filtered off, dried, and recrystallized from the appropriate solvent to produce **6a,b**.

2,4-Diamino-5'-methyl-6'-phenyl-4'H-spiro[1,5-benzodiazepine-3,3'-thieno[2,3-c]pyridazin]-4'-one (6a). Compound **6a** was obtained by refluxing **3a** (2.92 g, 0.01 mol) and *o*-phenylenediamine hydrochloride. The solid product recrystallized from dilute dimethyl formamide (80%) to produce **6a** as yellow crystals; IR spectrum (KBr) cm^{-1} : 3430–3360 (br, 2NH_2), 1680 (CO); ^1H NMR (DMSO-d_6) δ ppm: 2.13 (s, 3H, CH_3), 2.74 (br, 4H, 2NH_2 , D_2O exchangeable), 7.01–7.41 (m, 9H, aromatic protons). ^{13}C NMR (75 MHz, CDCl_3): δ 190 (s) ppm, 160 (s, 2C), 145 (s), 140 (s, 2C), 136 (s), 134 (s, 2C),

132 (s), 130 (d, 2C), 129 (d), 128 (d, 2C), 127 (d, 2C), 122 (d, 2C), 75 (s), 50 (q). MS (m/z): 400 (M^+) 70%.

2,4-Diacetyl-5'-methyl-6'-phenyl-4'H-spiro(1,5-benzodiazepine-3,3'-thieno [2,3-c]pyridazin)-4'-one (6b). Compound **6b** was obtained by refluxing **3b** (3.26g, 0.01 mol) and o-phenylenediamine. The solid product recrystallized from dilute dimethyl formamide (80%) to produce **6b**; 1710 (2CO), 1670 (CO); ^1H NMR (DMSO- d_6) δ ppm: 2.13 (s, 3H, CH_3), 2.24 (s, 6H, 2CH_3), 7.01–7.41 (m, 9H, aromatic protons); MS (m/z): 398 (M-86) 20%.

Biology

Measurement of Antimicrobial Activity Using Diffusion Disc Method.

The prepared compounds were tested against one strain of Gram positive bacteria (*Staphylococcus aureus* G^+), Gram-negative bacteria (*Escherichia coli* G^-), and fungi (*Aspergillus flavus* and *Candida albicans*).^{11,12}

REFERENCES

1. R. E. Dolle, B. Le Bourdonnec, A. J. Goodman, G. A. Morales, J. M. Salvino, and W. Zhang, *J. Comb. Chem.*, **9**, 855 (2007).
2. J. A. Gonzalez-Vera, T. Garcia-Lopez, and R. Herranz, *J. Org. Chem.*, **70**, 3660 (2005).
3. C. Ramalingan, S. Balasubramanian, and S. Kabilan, *Synth. Commun.*, **4**, 1105 (2004).
4. C. J. F. Bichard, E. P. Mitchell, M. R. Wormald, K. A. Waston, L. N. Johnson, S. E. Zographos, D. D. Koutra, N. G. Oikonomakos, and G. W. Fleet, *Tetrahedron Lett.*, **36**, 2145 (1995).
5. F. Risitano, G. Grassi, F. Foti, G. Bruno, and A. Rotondo, *Heterocycles*, **60**, 857 (2003).
6. V. Nair, K. V. Radhakrishnan, K. C. Sheela, and N. P. Rath, *Tetrahedron*, **55**, 14199 (1999).
7. K. C. Liu, and R. K. Howe, *J. Org. Chem.*, **48**, 4590 (1983).
8. (a) T. Ichiba, and P. J. Scheuer, *J. Org. Chem.*, **58**, 4149 (1993); (b) C. Lacy and P. J. Scheuer, *J. Nat. Prod.*, **63**, 119 (2000).
9. A. M. S. Youssef, R. A. M. Faty, and M. M. Youssef, *J. Korean Chem. Soc.*, **45**, 448 (2001).
10. E. S. Gewald and H. Bottcher, *Ber.*, **99**, 94 (1966).
11. D. N. Muanza, B. W. Klm, K. L. Euler, and L. Williams, *J. Pharmacog.*, **32**, 337 (1994).
12. O. N. Irob, M. Moo-Young, and W. A. Anderson, *J. Pharmacog.*, **34**, 87 (1996).