This article was downloaded by:

On: 30 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t713618290

The Reaction of Phosphonium Ylids with o-Quinones and Triketones

F. M. Soliman^a; Kh. M. Khalil^a; G. Abd-el-naim^a

^a National Research Centre, Cairo, Egypt

To cite this Article Soliman, F. M., Khalil, Kh. M. and Abd-el-naim, G.(1987) 'The Reaction of Phosphonium Ylids with o-Quinones and Triketones', Phosphorus, Sulfur, and Silicon and the Related Elements, 30: 3, 807

To link to this Article: DOI: 10.1080/03086648708079299 URL: http://dx.doi.org/10.1080/03086648708079299

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

The Reaction of Phosphonium Ylids with o-Quinones and Triketones

F. M. Soliman, Kh. M. Khalil and G. Abd-El-Naim
National Research Centre
Dokki, Cairo, Egypt.

The reaction of 1,2-benzo a phenazine-8,9-dione 1 and/or 1,2,3-indantrione 2, with phosphonium ylides has been studied.

When 1 was reacted with two molar amounts of methoxy-(3a) and/or ethoxycarbonylmethylenetriphenylphosphorane (3b), in THF, at the reflux temp. for 3 hrs, dimethyl (4a) and/or diethyl 1,2-dihydro benzo a furo [3,2-b] phenazine-1,2-dicarboxylate (4b), along with triphenylphosphine oxide and triphenylphosphine were obtained.

On the other hand, reaction of equimolar amounts of ylides 2 with the red trione 2 in THF at room temp., afforded colourless crys C tals of 2',4'-dihydroxyspiro [indan-2,3'(2'H)-indeno [1,2-b] pyran]-1,3,5'(4'H)-trione diacetate (5a) or dipropionate (5b), together with triphenylphosphine oxide. Formation of 6-membered dihydro C aromatic ring like 5, is considered as a new reaction of phos 2 phoranes. The structure of the new compounds 4 and 5 was confirmed and the reaction mechanisms are discussed.

$$\frac{1}{4} R R = C(0)OCH_3; b, R = C(0)OC_2H_5$$