

This article was downloaded by:

On: 29 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

A NEW STRUCTURAL FORM OF TIN OCTAHEDRALLY COORDINATED IN A DRUM-SHAPED MOLECULE

V. Chandrasekhar^a; Roberta O. Day^a; Robert R. Holmes^a

^a Department of Chemistry, University of Massachusetts, Amherst, Massachusetts

To cite this Article Chandrasekhar, V. , Day, Roberta O. and Holmes, Robert R.(1995) 'A NEW STRUCTURAL FORM OF TIN OCTAHEDRALLY COORDINATED IN A DRUM-SHAPED MOLECULE', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 98: 1, 417 — 421

To link to this Article: DOI: 10.1080/10426509508036965

URL: <http://dx.doi.org/10.1080/10426509508036965>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A NEW STRUCTURAL FORM OF TIN OCTAHEDRALLY COORDINATED IN A DRUM-SHAPED MOLECULE¹

V. CHANDRASEKHAR, ROBERTA O. DAY and ROBERT R. HOLMES*


*Department of Chemistry, University of Massachusetts,
 Amherst, Massachusetts 01003*

(Received February 11, 1985)

Although oligomeric organotin oxycarboxylates based on the composition $[\text{RSn}(\text{O})\text{O}_2\text{CR}]_n$ have been known for some time,² no indications of their possible structural features are available. As early as 1921 some members of this class were reported³ to be hexameric, e.g., $[\text{CH}_3\text{Sn}(\text{O})\text{O}_2\text{CR}]_6$.

We report herein the synthesis and X-ray crystal structure of hexameric phenyltin oxycyclohexanecarboxylate, $[\text{PhSn}(\text{O})\text{O}_2\text{CC}_6\text{H}_{11}]_6$. The X-ray characterization shows tin(IV) present in a new structural environment appearing as a drum-shaped molecule containing hexacoordinated tin atoms.

During the course of our work^{4,5} on triorganotin esters of carboxylic acids, we first obtained the hexamer while recrystallizing triphenyltin cyclohexanoate, $\text{Ph}_3\text{SnO}_2\text{C}(\text{C}_6\text{H}_{11})$. Presumably, it formed as a result of slow hydrolysis of the latter compound with the loss of two molecules of benzene. A more direct route involves the reaction of PhSnCl_3 with the sodium salt of the acid in CCl_4 solution, followed by a hydrolysis step which gave the compound as a powdery substance in modest yield⁶ present in a mixture of products:



Although a pure product has yet to be obtained by this route, both the crystalline hexamer and the powder had similar Nujol mull infrared spectra, melted with decomposition at 340°C, and were insoluble in organic solvents and in water. It thus appears that the hexamer is hydrolytically and thermally stable.

The schematic in Figure 1 portrays the drum-shaped structure resulting from the X-ray analysis⁷ and shows the atom labeling. The high symmetry associated with the molecule is revealed in Figure 2. The hexamer has crystallographic C_i symmetry. However, the noncrystallographic symmetry of the molecule is higher, namely S_6 , assuming free rotation about the C—C single bonds.

Reprinted with permission from *Inorg. Chem.*, 1985, 24, 1970. Copyright 1985 American Chemical Society.




FIGURE 1 Schematic representation of the drum structure of $[\text{PhSn}(\text{O})\text{O}_2\text{CC}_6\text{H}_{11}]_6$




FIGURE 2 ORTEP plot of $[\text{PhSn}(\text{O})\text{O}_2\text{CC}_6\text{H}_{11}]_6$ viewed down the S_6 axis with thermal ellipsoids at the 50% probability level

The six tin atoms are chemically equivalent, as are the six trivalent oxygen atoms. The Sn—O framework of the molecule can be described as a drum with top and bottom faces each being comprised of a six-membered $(-\text{Sn}-\text{O}-)_3$ tristannoxane ring.

The drum faces are joined together by six Sn—O bonds containing tricoordinated oxygen atoms. The sides of the drum are thus comprised of six four-membered ($-\text{Sn}-\text{O}-$)₂ distannoxane rings, each of which is spanned by a carboxylate group that forms a symmetrical bridge between two tin atoms. The sides of the structure are reminiscent of the ladder arrangement in dimeric distannoxanes recently found for aryl⁸ and alkyl^{9,10} derivatives, e.g., $[\text{Ph}_2(\text{Cl})\text{SnOSnOPh}_2(\text{Cl})]_2$. Concomitant with the equivalence of the Sn—O bonds to a particular carboxylate group is the corresponding equivalence of the pairs of C—O carboxylate bond lengths. The hexamer provides the first example of tin atoms symmetrically bridged by a carboxylate group. A summary of important bond parameters are listed in Table I.

The distannoxane ring units of the sides of the drum (i.e., $\text{Sn1}-\text{O1}-\text{Sn2}-\text{O2}$) are not planar but are folded along the Sn—Sn vectors so that the oxygen atoms are directed toward the interior of the cavity. This is apparent in Figure 2. Similarly, the tristannoxane ring faces of the drum (i.e., $\text{Sn1}-\text{O2}-\text{Sn3}-\text{O1}'-\text{Sn2}'-\text{O3}'$) are not planar but have the oxygen atoms directed toward the interior of the cavity

TABLE I
Summary of Selected Bond Distances (Å) and Angles (deg) for
 $[\text{PhSn}(\text{O})\text{O}_2\text{CC}_6\text{H}_{11}]_6$

Tristannoxane Ring, $(-\text{Sn}-\text{O}-)_3^a$			
Sn3'-O1-Sn2	133.7 (2)	Sn2-O3-Sn1'	135.2 (2)
Sn1-O2-Sn3	133.8 (2)	av	134.2
O3'-Sn1-O2	102.2 (1)	O2-Sn3-O1'	104.1 (1)
O1-Sn2-O3	102.7 (1)	av	103.0
Sn1-O2	2.083 (3)	Sn3-O2	2.080 (3)
Sn1-O3'	2.073 (3)	Sn3-O1'	2.069 (3)
Sn2-O1	2.081 (3)	av	2.079
Sn2-O3	2.089 (3)		
Distannoxane Ring, $(-\text{Sn}-\text{O}-)_2$			
Sn1-O1-Sn2	99.3 (1)	Sn3-O3-Sn2	99.6 (1)
Sn1-O1-Sn3'	99.6 (1)	Sn3-O3-Sn1'	100.3 (1)
Sn2-O2-Sn1	100.1 (1)	av	99.8
Sn2-O2-Sn3	100.0 (1)		
O1-Sn1-O2	77.7 (1)	O3-Sn3-O2	78.0 (1)
O1-Sn1-O3'	77.6 (1)	O3-Sn3-O1'	78.2 (1)
O2-Sn2-O1	78.4 (1)	av	78.0
O2-Sn2-O3	77.9 (1)		
Sn1-O1	2.124 (3)	Sn3-O3	2.098 (3)
Sn2-O2	2.096 (3)	av	2.106
Carboxylate			
Sn1-O11	2.165 (3)	Sn3-O31	2.139 (4)
Sn1-O22	2.155 (4)	Sn3-O12'	2.149 (4)
Sn2-O21	2.153 (4)	av	2.153
Sn2-O32	2.154 (4)		
C11-O11	1.266 (6)	C31-O31	1.242 (8)
C11-O12	1.261 (6)	C31-O32	1.277 (7)
C21-O21	1.259 (6)	av	1.260
C21-O22	1.252 (6)		

^aSimilar Sn—O—Sn (133.1–134.5°) and O—Sn—O (105.9–106.9°) bond angles and slightly shorter Sn—O bond lengths (1.952–1.978 Å) are present in the oxides (*t*-Bu₂SnO)₃ and (*t*-Am₂SnO)₃.¹²

an average of 0.24 Å, relative to the Sn atoms. This contrasts with the planar six-membered tin–oxygen rings found in the di-*tert*-butyltin^{11,12} and di-*tert*-amyltin¹² oxides, (t-Bu₂SnO)₃ and (t-Am₂SnO)₃, containing tetracoordinate tin. The distance between the plane defined by Sn1, Sn2', Sn3 and Sn1', Sn2, Sn3' is 2.315 Å. The corresponding distance between the planes defined by the oxygen atoms is 1.844 Å.

Thus, the interior of the cavity is defined by a crown of six oxygen atoms, in a trigonal-antiprismatic arrangement, where the average O—O distance is 2.633 Å and the average distance from the center of the cavity to the O atoms is 2.093 Å. On the basis of a van der Waals radius¹³ of 1.40 Å for oxygen, the interior of the cavity could thus host a species with a radius of approximately 0.69 Å. The entrance to the cavity is defined by three oxygen atoms arranged as an approximate equilateral triangle with an average edge length (oxygen center to oxygen center) of 3.254 Å. Again with 1.40 Å as the van der Waals radius of oxygen, species with radii up to about 0.5 Å could gain entrance to the cavity. Thus, encapsulation of ions as large as Li⁺, Be²⁺, or Mg²⁺ might be possible.

ACKNOWLEDGEMENT

The support of this research by the National Science Foundation (Grant No. CHE 8205411) and the donors of the Petroleum Research Fund, administered by the American Chemical Society, is gratefully acknowledged, as is the generous allocation of computing time by the University of Massachusetts Computing Center.

Supplementary Material Available: Atomic coordinates, anisotropic thermal parameters, hydrogen atom parameters, and bond lengths and bond angles for [PhSn(O)O₂CC₆H₁₁]₆ (Tables SI–SIV, respectively) and a listing of observed and calculated structure factors (26 pages). Ordering information is given on any current masthead page.

REFERENCES AND NOTES

1. "Abstracts of Papers," 189th National Meeting of the American Chemical Society, Miami, FL, April 1985; American Chemical Society: Washington, DC, 1985; INOR P044.
2. Anderson, H. H. *Inorg. Chem.* **1964**, *3*, 912.
3. Lambourne, H. J. *Chem. Soc.* **1921**, *121*, 2533.
4. Swisher, R. G.; Vollano, J. F.; Chandrasekhar, V.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1984**, *23*, 3147.
5. Vollano, J. F.; Day, R. O.; Rau, D. N.; Chandrasekhar, V.; Holmes, R. R. *Inorg. Chem.* **1984**, *23*, 3153.
6. Phenyltin trichloride (1.1 g, 3.6 mmol) and the sodium salt of cyclohexanecarboxylic acid (0.55 g, 3.6 mmol) were reacted in carbon tetrachloride (100 mL) at reflux temperature for 2 h. Filtration of NaCl and removal of the solvent in vacuo yielded a residue that was heated with a mixture of 95% ethanol (10 mL), acetonitrile (10 mL), and water (5 mL) for 2 h. A white insoluble compound was isolated (0.3 g); 25% yield, mp 340°C dec. The infrared spectrum of this compound in Nujol shows $\nu(\text{CO}_2)$ at 1530 cm⁻¹.
7. The hexameric stannoxane [PhSn(O)O₂CC₆H₁₁]₆ crystallizes in the monoclinic space group *P*2₁/*n* with *a* = 13.919 (3) Å, *b* = 19.361 (3) Å, *c* = 15.089 (2) Å, β = 90.81 (1)°, *Z* = 2, $\mu_{\text{MoK}\alpha}$ = 1.891 mm⁻¹. A total of 5984 independent reflections were measured at room temperature with use of the θ – 2θ scan mode and graphite-monochromated MoK α radiation, for $2^\circ \leq 2\theta_{\text{MoK}\alpha} \leq 47^\circ$, on an Enraf-Nonius CAD4 diffractometer. No corrections were made for absorption. Full-matrix least-squares refinement (nonhydrogen atoms anisotropic, hydrogen atoms fixed isotropic, function minimized $\sum w(|F_o| - |F_c|)^2$, $w^{1/2} = 2F_o L p / \sigma_f$) led to *R* = 0.030 and *R_w* = 0.044 for the 5093 reflections having $I \geq 2\sigma_f$.
8. Vollano, J. F.; Day, R. O.; Holmes, R. R. *Organometallics* **1984**, *3*, 745.

9. Harrison, P. G.; Begley, M. J.; Molloy, K. C. *J. Organomet. Chem.* **1980**, *186*, 213.
10. Puff, H.; Friedrichs, E.; Visel, F. Z. *Anorg. Allg. Chem.* **1981**, *477*, 50.
11. Puff, H.; Schuh, W.; Sievers, R.; Zimmer, R. *Angew. Chem., Int. Ed. Engl.* **1981**, *20*, 591.
12. Puff, H.; Schuh, W.; Sievers, R.; Wald, W.; Zimmer, R. *J. Organomet. Chem.* **1984**, *260*, 271.
13. Pauling, L. "The Nature of the Chemical Bond," 3rd ed.; Cornell University Press: Ithaca, NY, 1960.