

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Stereoselective Epoxidation of bis-Sulfinyl Alkenes and Application to the Asymmetric Synthesis of α -Substituted Carboxylic Acids

Varinder K. Aggarwal; Julia M. Worrall; Rikki Alexander

To cite this Article Aggarwal, Varinder K. , Worrall, Julia M. and Alexander, Rikki(1997) 'Stereoselective Epoxidation of bis-Sulfinyl Alkenes and Application to the Asymmetric Synthesis of α -Substituted Carboxylic Acids', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 120: 1, 351 – 352

To link to this Article: DOI: 10.1080/10426509708545544

URL: <http://dx.doi.org/10.1080/10426509708545544>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Stereoselective Epoxidation of bis-Sulfinyl Alkenes and Application to the Asymmetric Synthesis of α -Substituted Carboxylic Acids

Varinder K. Aggarwal,^{a*} Julia M. Worrall,^a and Rikki Alexander.^b


^a*Department of Chemistry, University of Sheffield, Sheffield, S3 7HF, UK.*

^b*Celltech Therapeutics Limited, 216 Bath Road, Slough SL1 4EN, UK.*

Ketene thioacetals **3** can be easily prepared in optically pure form and react diastereoselectively with metal peroxides. The resulting bis-sulfinyl oxiranes can be converted in a single step to α -substituted acids.

INTRODUCTION


1,3-dithiane-1,3-dioxide **1** can be prepared in optically pure form via Modena oxidation of 1,3-dithiane-2-ethylcarboxylate.¹ It has been found to be effective as a diastereoselective carbonyl anion equivalent in reactions with aldehydes (scheme 1).²



Scheme 1 Reagents: i, (R,R)-DET, $\text{Ti}(\text{OiPr})_4$, PhCMe_2OOH , CH_2Cl_2 ; ii, NaOH , H_2O ; iii, NaHMDS , RCHO , THF , py

RESULTS AND DISCUSSION

We initially envisaged expanding on this chemistry by using 2-halo-1,3-dithiane-1,3-dioxide as a carbonyl anion equivalent in Darzens type reactions, providing access to spirocyclic epoxides **2**. Diastereoselective addition to aldehydes was achieved^{3,4} but the ring closure step proved problematic, giving low overall yields of epoxides.



Excellent diastereoselectivities in epoxidations of electron deficient alkenes have been achieved by Jackson, using metal peroxides as nucleophilic oxidants.⁵ We decided to apply his conditions to the oxidation of the ketene thioacetals **3**, prepared by aldol condensation of **1** with aldehydes. This was found to be a successful, highly diastereoselective alternative route to **2** (table 1). A more general route to the ketene thioacetals **3** was developed using Wadsworth-Emmons type reactions of phosphonate **4** with aldehydes. Subjection of dithiane phosphonate **4** to the Modena asymmetric oxidation conditions afforded the dioxide **5** in high enantiomeric excess (scheme 2).

Scheme 2 Oxidation route to the target epoxides




Table 1 Diastereoselective nucleophilic epoxidation of ketene thioacetals **3**

The masked carboxylic acid functionality α - to the newly formed stereocentre was revealed by nucleophilic opening of the epoxide.

Acknowledgements: We thank Celltech Therapeutics Ltd and Sheffield University for financial support.

References:

1. V. K. Aggarwal, G. Evans, E. Moya, J. Dowden, *J. Org. Chem.*, **57**, 6390-6391 (1992).
2. V. K. Aggarwal, R. Franklin, J. Maddock, G. R. Evans, A. Thomas, M. F. Mahon, K. C. Molloy, M. J. Rice, *J. Org. Chem.*, **60**, 2174-2182 (1995).
3. V. K. Aggarwal, J. M. Worrall, H. Adams, R. Alexander, *Tetrahedron Lett.*, **35**, 6167-6170 (1994).
4. V. K. Aggarwal, G. Boccardo, J. M. Worrall, H. Adams, R. Alexander, *J. Chem. Soc., Perkin Trans. I*, Submitted.
5. A. D. Briggs, R. F. W. Jackson, W. Clegg, M. R. J. Elsegood, J. Kelly, P.A. Brown, *Tetrahedron Lett.*, **35**, 6945-6948 (1994).