


This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

SYNTHESIS OF BIHETEROCYCLIC α -AMINOPHOSPHONIC ACID DERIVATIVES

S. Achamlale^a; A. Elachqar^a; A. El Hallaoui^a; A. Alami^a; S. Elhajji^a; M. L. Roumestant^{ab}; Ph. Viallefont^c

^a Laboratoire de Chimie Organique, Faculté des Sciences Dhar El Mehraz, Université Sidi Mohamed Ben Abdellah, Fes, Maroc ^b Lab. de Chimie Org., Université Montpellier II, Montpellier Cedex, France ^c UMR 5810, Université MONTPELLIER II, Montpellier, France

To cite this Article Achamlale, S. , Elachqar, A. , Hallaoui, A. El , Alami, A. , Elhajji, S. , Roumestant, M. L. and Viallefont, Ph.(1998) 'SYNTHESIS OF BIHETEROCYCLIC α -AMINOPHOSPHONIC ACID DERIVATIVES', Phosphorus, Sulfur, and Silicon and the Related Elements, 140: 1, 103 – 111

To link to this Article: DOI: 10.1080/10426509808035736

URL: <http://dx.doi.org/10.1080/10426509808035736>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS OF BIHETEROCYCLIC α -AMINOPHOSPHONIC ACID DERIVATIVES

S. ACHAMLALE^a, A. ELACHQAR^a, A. EL HALLAOUTI^a, A. ALAMI^a,
S. ELHAJJI^a, M.L. ROUMESTANT^{b*} and PH. VIALLEFONT^b

^a*Laboratoire de Chimie Organique, Faculté des Sciences Dhar El Mehraz Université Sidi Mohamed Ben Abdellah, Fes, Maroc* and ^b*UMR 5810, Université MONTPELLIER II, Montpellier, France*

(Received March 19, 1998)

We report here the synthesis of biheterocyclic α -aminophosphonic acid derivatives by 1,3 dipolar cycloaddition of acetylenic compounds, prepared in the laboratory, on α -azido α -amino phosphonic esters **1**^[1a].

Keywords: α -aminophosphonic acid; biheterocycle; 1,2,3-triazole; tetrazole

INTRODUCTION


α -aminophosphonic acids, analogues of α -amino acids display diverse and useful biological properties^[2-4]; these compounds are found to be substrates and inhibitors of enzymes, as well as plant growth regulators and herbicides. They also display antibacterial properties and neuronal activities^[3-4].

In this paper we describe new heterocyclic analogues of α -amino acids able to present an interesting biological activity due to the presence of heterocyclic cores as in the case of tetrazolic glycine^[5] and proline^[6] derivatives. This approach consists of preparing, firstly, tetrazolic dipolarophiles by nucleophilic substitution of propargyl bromide by 5-aryl substituted tetrazoles.

* Address to Author: Lab. de Chimie Org., Université Montpellier II, 5 Av. Ch. Flahaut, F-34060 Montpellier Cedex, France.


RESULTS

Tetrazole derivatives were obtained as described in the literature^[7], the yields (35 to 75%) being improved in our team^[1c,7]. There is literature^[8] report which illustrates the existence of the two tetrazole tautomeric forms (scheme 1).




SCHEME 1

Treatment of 5-aryl substituted tetrazoles with propargyl bromide in the presence of triethylamine in acetone at room temperature during 4 h afforded the two regioisomers (2,5-disubstituted / 1,5-disubstituted) in good yields (70–85%) in a ratio of approximately 80/20, separable by column chromatography, whose structures were determined unambiguously by X-Ray diffraction analysis (scheme 2)^[9].



SCHEME 2

The 2,5 and 1,5-disubstituted tetrazoles thus prepared were submitted to cycloaddition reaction with azide **1** (scheme 3). The azide **1** was obtained by reaction^[1a] of sodium azide on the α -bromo- α -amino phosphonic ester. Results are summarised in table I.



SCHEME 3

TABLE I Synthesis of biheterocyclic α-amino phosphonic acid derivatives (2-8) a,b

Product	Ar	m.p. (°C)	Time (h)	Yield (%)	Ratio of isomers 1,5/1,4
2b	C ₆ H ₅	127–129	48	80	20
2a	C ₆ H ₅	139–141			80
3b	p-Me-C ₆ H ₄	145–147	48	75	47
3a	p-Me-C ₆ H ₄	116–118			53
4b	p-MeO-C ₆ H ₄	110–112	48	73	40
4a	p-MeO-C ₆ H ₄	98–100			60
5b	α-thienyl	101–103	48	69	25
5a	α-thienyl	122–124			75
6a	o-HO-C ₆ H ₄	186–188	48	75	>98
7b	α-furyl	151–153	48	72	10
7a	α-furyl	139–140			90
8a	α-pyrryl	170–172	48	79	>98

If we consider now the triazole ring after cycloaddition reaction, two regioisomers (1,4-isomer and 1,5-isomer) were formed in good yields and separated. The regioselectivity was excellent, only the 1,4-isomer was obtained in two cases: from **6a** (Ar = o-HO-C₆H₄) and **8a** (Ar = α -pyrryl).

The structures of the two regioisomers (**2-8**) **a,b** were assigned on the basis of literature data^[1d,8,10-12] concerning the chemical shifts of triazolic protons^[1d,10,11] and the chemical shifts of the carbons of the triazole ring in positions 4 and 5. The studies carried out^[1d,10,11] have shown that the triazolic proton signal for the 1,5-isomer lies downfield from the corresponding signal for the 1,4-isomer but in ¹³C NMR^[8,12] the carbon signal (HC₄) for the 1,5-isomer lies upfield from the corresponding carbon signal (HC₅) for the 1,4-isomer. (Table II)

TABLE II The chemical shifts of triazolic protons and carbons in ¹H NMR and ¹³C NMR for 1,4 and 1,5-isomers

Ar	Isomer 1,5	Isomer 1,4	¹ H NMR		¹³ C NMR	
			δ_{H_4} ppm	δ_{H_5} ppm	δ_{C_4H} ppm	δ_{C_5H} ppm
C ₆ H ₅	2b	-	7.76	-	135.04	- 58
	-	2a	-	8.28	-	124.58
p-Me-C ₆ H	3b	-	7.75	-	135.06	-
	-	3a	-	8.35	-	124.63
p-MeO-C ₆ H ₄	4b	-	7.75	-	135.00	-
	-	4 a	-	8.3 1	-	124.57
α -thienyl	5b	-	7.75	-	135.03	-
	-	5a	-	8.30	-	124.65
o-HO-C ₆ H ₄	-	6	-	8.52	-	125.30
	7b	-	7.57	-	134.56	-
α -furyl	-	7 a	-	8.30	-	124.5 1
	-	8	-	8.39	-	124.21

By this strategy we have obtained different biheterocyclic α -aminophosphonic acid derivatives in good yields.

EXPERIMENTAL

Melting points were obtained on a electrothermal melting point apparatus and are uncorrected. ^1H NMR spectra were obtained on VARIAN EM – 360 (60 MHz) and BRUCKER (250 MHz) instruments, TMS as internal standard. ^{13}C NMR spectra were obtained on BRUCKER (200 MHz) instrument. Microanalyses were performed by the ENCT (Toulouse). Mass spectra were measured on a JEOL-JMS-DX 300 FAB instrument and Desorption in chemical ionisation with NH_3 D/CI instrument. 5-substituted tetrazoles have been prepared using Alami's method^[1c].

Cycloaddition reaction: General procedure

The azide **1** and the dipolarophile were magnetically stirred in a minimum volume of benzene at reflux (see table II for the reaction conditions). After evaporation of the solvent, the residue was chromatographed over silica gel or by preparative liquid chromatography (HPLC). The solid compounds are recrystallized from ether/dichloromethane and/or from acetone.

Minor isomer **2b**: Rf = 0.24 (ether).

^1H NMR (CDCl_3) δ : 1.08 (t, 3H, J = 7Hz), 1.31 (t, 3H, J = 7Hz), 3.75–4.35 (m, 2x2H), 5.93 (s, 2H), 7.09 (dd, 1H, J_1 = 10Hz, J_2 = 15.5Hz), 7.35–7.4 (m, 6H), 7.52 (dd, NH, J_1 = 10Hz, J_3 = 4Hz), 7.76 (s, 1H), 7.79–7.88 (m, 2H), 8.07–8.14 (m, 2H).

^{13}C NMR (CDCl_3) δ : 16.32 (2x $\text{CH}_3\text{CH}_2\text{O}$), 48.03 ($-\text{CH}_2-$), 64.09 (d, $-\text{CH}-$, $^1J_{\text{C-P}}$ = 180.2Hz), 64.45 (2x CH_2O), 126.9, 127.1, 127.6, 128.8, 128.93, 130.52, 132.35, 132.72 (2x C_6H_5), 135.04 ($\text{CH}_4=\text{C}$), 141.88 ($\text{CH}_4=\text{C}$), 165.63 ($\text{C}_6\text{H}_5-\text{CN}_4$), 166.8 (C=O). $\text{C}_{22}\text{H}_{25}\text{N}_8\text{O}_4\text{P}$ (496): M.S [D.C.I/ NH_3 ($\text{M}+\text{H}$) $^+$ = 497 and ($\text{M}+\text{NH}_4$) $^+$ = 514].

Major isomer **2a**: m.p. = 139–141°C Rf = 0.16 (ether).

^1H NMR (CDCl_3) δ : 1.06 (t, 3H, J = 7Hz), 1.29 (t, 3H, J = 7Hz), 3.75–4.35 (m, 2x2H), 5.98 (s, 2H), 7.08 (dd, 1H, J_1 = 10Hz, J_2 = 16Hz), 7.26–7.5 (m, 6H), 7.64–7.69 (m, 2H), 8.04–8.09 (m, 2H), 7.28 (s, 1H), 8.6 (dd, NH, J_1 = 10Hz, J_3 = 4Hz).

^{13}C NMR (CDCl_3) δ : 16.2 (2x $\text{CH}_3\text{CH}_2\text{O}$), 48.3 ($-\text{CH}_2-$), 60.4 (d, $-\text{CH}-$, $^1J_{\text{C-P}}$ = 182Hz), 64.76 (2x CH_2O), 124.58 ($\text{CH}_5=\text{C}$), 126.9, 127.2, 127.89,

128.6, 128.87, 130.41, 132.23, 132.7 (2x C_6H_5), 140.55 ($\text{CH}_5=\text{C}$), 165.46 ($\text{C}_6\text{H}_5-\text{CN}_4$), 167.5 ($\text{C}=\text{O}$). Anal. calcd. for $\text{C}_{22}\text{H}_{25}\text{N}_8\text{O}_4\text{P}$: C, 53.22, H, 5.04, N, 22.58.

Found. C, 52.93, H, 4.96, N, 22.50.

Minor isomer **3b**: $R_f = 0.3$ (ether).

^1H NMR (CDCl_3) δ : 1.06 (t, 3H, $J = 7\text{Hz}$), 1.31 (t, 3H, $J = 7\text{Hz}$), 2.38 (s, 3H), 3.7–4.31 (m, 2x2H), 5.91 (s, 2H), 7.1 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 15.5\text{Hz}$), 7.2–7.6 (m, 5H), 7.75 (s, 1H), 7.8–8.03 (m, 5H).

^{13}C NMR (CDCl_3) δ : 16.4 (2x $\text{CH}_3\text{CH}_2\text{O}$), 21.52 (CH_3), 47.96 ($-\text{CH}_2-$), 64.05 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 180.4\text{Hz}$), 64.47 (2x CH_2O), 124.29, 126.81, 127.35, 127.64, 128.8, 129.63, 132.72, 140.73 (C_6H_5 and p-Me- C_6H_4), 135.06 ($\text{CH}_4=\text{C}$), 141.95 ($\text{CH}_4=\text{C}$), 165.7 (Me- $\text{C}_6\text{H}_4-\text{CN}_4$), 166.8 ($\text{C}=\text{O}$).

Anal. calcd. for $\text{C}_{23}\text{H}_{27}\text{N}_8\text{O}_4\text{P}$: C, 54.12, H, 5.3, N, 21.96.

Found. C, 53.87, H, 5.36, N, 21.93.

Major isomer **3a**: $R_f = 0.22$ (ether).

^1H NMR (CDCl_3) δ : 1.02 (t, 3H, $J = 7\text{Hz}$), 1.26 (t, 3H, $J = 7\text{Hz}$), 2.35 (s, 3H), 3.64–4.28 (m, 2x2H), 5.97 (s, 2H), 7.15 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 16.7\text{Hz}$), 7.16–7.56 (m, 5H), 7.84–7.96 (m, 4H), 8.35 (s, 1H), 8.95 (dd, NH, $J_1 = 10\text{Hz}$, $J_3 = 4\text{Hz}$).

^{13}C NMR (CDCl_3) δ : 16.26 (2x $\text{CH}_3\text{CH}_2\text{O}$), 21.5 (CH_3), 48.22 ($-\text{CH}_2-$), 60.4 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 182.78\text{Hz}$), 64.8 (2x CH_2O), 124.63 ($\text{CH}_5=\text{C}$), 124.38, 126.78, 128, 128.5, 129.56, 132.3, 132.6, 140.56 (C_6H_5 and p-Me- C_6H_4), 140.6 ($\text{CH}_5=\text{C}$), 165.5 (Me- $\text{C}_6\text{H}_5-\text{CN}_4$), 167.6 ($\text{C}=\text{O}$).

$\text{C}_{23}\text{H}_{27}\text{N}_8\text{O}_4\text{P}$ (510): M.S [D.C.I/NH_3 ($\text{M}+\text{H}$) $^+ = 511$ and ($\text{M}+\text{NH}_4$) $^+ = 528$].

Minor isomer **4b**: $R_f = 0.33$ (AcOEt/DCM 3/7).

^1H NMR (CDCl_3) δ : 1.07 (t, 3H, $J = 7\text{Hz}$), 1.31 (t, 3H, $J = 7\text{Hz}$), 3.84 (s, 3H), 3.7–4.31 (m, 2x2H), 5.9 (s, 2H), 7.09 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 15.5\text{Hz}$), 7.3–7.87 (m, 5H+NH), 7.56 (AA'BB' System, 4H, $J_{\text{AB}} = J_{\text{A'B'}} = 9\text{Hz}$), 7.75 (s, 1H).

^{13}C NMR (CDCl_3) δ : 16.31 (2x $\text{CH}_3\text{CH}_2\text{O}$), 48.33 ($-\text{CH}_2-$), 55.4 (CH_3O), 64.05 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 180.2\text{Hz}$), 64.45 (2x CH_2O), 114.33, 119.68, 127.61, 128.41, 128.68, 130.43, 132.35, 132.72, 161.43 (C_6H_5 and p-MeO- C_6H_4), 135 ($\text{CH}_4=\text{C}$), 141.99 ($\text{CH}_4=\text{C}$), 165.5 (MeO- $\text{C}_6\text{H}_4-\text{CN}_4$), 166.8 ($\text{C}=\text{O}$).

Anal. calcd. for $C_{23}H_{27}N_8O_5P$: C, 52.47, H, 5.13, N, 21.29.

Found. C, 52.53, H, 5.10, N, 21.57.

Major isomer **4a**: $R_f = 0.2$ (AcOEt/DCM 3/7).

1H NMR ($CDCl_3$) δ : 1.04 (t, 3H, $J = 7$ Hz), 1.27 (t, 3H, $J = 7$ Hz), 3.82 (s, 3H), 3.80–4.30 (m, 2x2H), 5.96 (s, 2H), 7.1 (dd, 1H, $J_1 = 10$ Hz, $J_2 = 15.5$ Hz), 7.24–7.95 (m, 5H), 7.45 (AA'BB'System, 4H, $J_{AB} = J_{A'B'} = 9$ Hz), 8.31 (s, 1H), 8.8 (dd, NH, $J_1 = 10$ Hz, $J_3 = 4$ Hz)

^{13}C NMR ($CDCl_3$) δ : 16.35 (2x $\underline{CH_3CH_2O}$), 48.10 ($-\underline{CH_2}-$), 55.38 ($\underline{CH_3O}$), 60.4 (d, $-\underline{CH}$, $^1J_{C-P} = 182.5$ Hz), 64.81 (2x $\underline{CH_2O}$), 124.57 ($\underline{CH_5=C}$), 114.27, 119.78, 127.74, 127.94, 128.38, 128.57, 128.72, 132.23, 132.66, 161.32 ($\underline{C_6H_5}$ and $p\text{-MeO-}\underline{C_6H_4}$), 140.63 ($\underline{CH_5=C}$), 165.3 ($\text{Me-}\underline{C_6H_5-CN_4}$), 167.5 (C=O).

$C_{23}H_{27}N_8O_5P$ (526): M.S [D.C.I/ NH_3 ($M+H$) $^+$ = 527 and ($M+NH_4$) $^+$ = 544].

Minor isomer **5b**: $R_f = 0.36$ (AcOEt/Et₂O 1/2).

1H NMR ($CDCl_3$) δ : 1.08 (t, 3H, $J = 7$ Hz), 1.32 (t, 3H, $J = 7$ Hz), 3.72–4.35 (m, 2x2H), 5.9 (s, 2H), 7.1 (dd, 1H, $J_1 = 10$ Hz, $J_2 = 15.5$ Hz), 7.11–7.13 (m, 1H), 7.39–7.9 (m, 8H), 7.75 (s, 1H).

^{13}C NMR ($CDCl_3$) δ : 16.31 (2x $\underline{CH_3CH_2O}$), 48.01 ($-\underline{CH_2}-$), 64.12 (d, $-\underline{CH}$, $^1J_{C-P} = 180.16$ Hz), 64.47 (2x $\underline{CH_2O}$), 127.63, 127.98, 128.1, 128.18, 128.8, 132.34, 132.7 ($\underline{C_6H_5}$ and $\underline{C_4H_3S}$), 135.03 ($\underline{CH_4=C}$), 141.7 ($\underline{CH_4=C}$), 161.9 ($\alpha\text{-Thienyl-}\underline{CN_4}$), 166.8 (C=O). Anal. calcd. for $C_{20}H_{23}N_8O_4PS$: C, 47.81, H, 4.58, N, 22.31.

Found. C, 47.41, H, 4.55, N, 22.16.

Major isomer **5a**: $R_f = 0.28$ (AcOEt/Et₂O 1/2).

1H NMR ($CDCl_3$) δ : 1.06 (t, 3H, $J = 7$ Hz), 1.28 (t, 3H, $J = 7$ Hz), 3.74–4.32 (m, 2x2H), 5.96 (s, 2H), 7.07–7.1 (m, 1H), 7.14 (dd, 1H, $J_1 = 10$ Hz, $J_2 = 15.5$ Hz), 7.26–7.94 (m, 6H), 7.7–7.73 (m, 1H), 7.3 (s, 1H), 8.77 (dd, NH, $J_1 = 10$ Hz, $J_3 = 4$ Hz).

^{13}C NMR ($CDCl_3$) δ : 16.2 (2x $\underline{CH_3CH_2O}$), 48.27 ($-\underline{CH_2}-$), 60.45 (d, $-\underline{CH}$, $^1J_{C-P} = 182.43$ Hz), 64.78 (2x $\underline{CH_2O}$), 124.65 ($\underline{CH_5=C}$), 127.42, 127.94, 128, 128.55, 128.82, 132, 132.26, 132.64 ($\underline{C_6H_5}$ and $\underline{C_4H_3S}$), 140.36 ($\underline{CH_5=C}$), 161.5 ($\alpha\text{-Thienyl-}\underline{CN_4}$), 167.5 (C=O).

Anal. calcd. for $C_{20}H_{23}N_8O_4PS$: C, 47.81, H, 4.58, N, 22.31.

Found. C, 47.72, H, 4.59, N, 21.85.

Major isomer **6a**: Rf = 0.2 (AcOEt/DCM 2/3).

^1H NMR ($\text{CDCl}_3 + \text{DMSO } d_6$) δ : 1.06 (t, 3H, $J = 7\text{Hz}$), 1.24 (t, 3H, $J = 7\text{Hz}$), 3.73–4.27 (m, 2x2H), 6.04 (s, 2H), 6.8–8 (m, 9H), 7.2 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 15.5\text{Hz}$), 8.52 (s, 1H), 9.68 (dd, NH, $J_1 = 10\text{Hz}$, $J_3 = 4\text{Hz}$), 11.5 (s, 1H).

^{13}C NMR ($\text{CDCl}_3 + \text{DMSO } d_6$) δ : 16.8 (2x $\text{CH}_3\text{CH}_2\text{O}$), 49.13 ($-\text{CH}_2-$), 62.96 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 180.1\text{Hz}$), 64.7 (2x CH_2O), 125.3 ($\text{CH}_5=\text{C}$), 112.15, 118, 120.5, 126.4, 128.6, 128.8, 132.7, 132.9, 133.1, 156.64 (C_6H_5 and o- $\text{HO}-\text{C}_6\text{H}_4$), 140.67 ($\text{CH}_5=\text{C}$), 165.1 (o- $\text{HO}-\text{C}_6\text{H}_4-\text{CN}_4$), 168 (C=O).

Anal. calcd. for $\text{C}_{22}\text{H}_{25}\text{N}_8\text{O}_5\text{P}$: C, 51.56, H, 4.88, N, 21.87.

Found: C, 51.40, H, 4.71, N, 21.55.

Minor isomer **7b**: Rf = 0.67 (ether).

^1H NMR (CDCl_3) δ : 1.01 (t, 3H, $J = 7\text{Hz}$), 1.29 (t, 3H, $J = 7\text{Hz}$), 3.63–4.28 (m, 2x2H), 5.92 (s, 2H), 6.54–6.56 (m, 1H), 7.1 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 16\text{Hz}$), 7.25–7.27 (m, 1H), 7.3–7.84 (m, 5H), 7.62–7.64 (m, 1H), 7.57 (s, 1H), 8.1 (dd, NH, $J_1 = 10\text{Hz}$, $J_3 = 4\text{Hz}$).

^{13}C NMR (CDCl_3) δ : 16.2 (2x $\text{CH}_3\text{CH}_2\text{O}$), 43.71 ($-\text{CH}_2-$), 64.04 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 180.52\text{ Hz}$), 64.4 (2x CH_2O), 112.56, 115.44, 139.1, 145.5 ($\text{C}_4\text{H}_3\text{O}$), 127.50, 128.56, 132.23, 132.7 (C_6H_5), 134.56 ($\text{CH}_4=\text{C}$), 142.51 ($\text{CH}_4=\text{C}$), 145.91 ($\alpha\text{-Furyl}-\text{CN}_4$), 167 (C=O). $\text{C}_{20}\text{H}_{23}\text{N}_8\text{O}_5\text{P}$ (486): M.S [D.C.I./ NH_3 ($\text{M}+\text{H}$) $^+$ = 487 and ($\text{M}+\text{NH}_4$) $^+$ = 504].

Major isomer **7a**: Rf = 0.48 (ether).

^1H NMR (CDCl_3) δ : 1.02 (t, 3H, $J = 7\text{Hz}$), 1.25 (t, 3H, $J = 7\text{Hz}$), 3.58–4.32 (m, 2x2H), 5.97 (s, 2H), 6.54–6.57 (m, 1H), 7.12 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 16\text{Hz}$), 7.18–7.9 (m, 7H), 8.3 (s, 1H), 9 (dd, NH, $J_1 = 10\text{Hz}$, $J_3 = 4\text{Hz}$).

^{13}C NMR (CDCl_3) δ : 16.3 (2x $\text{CH}_3\text{CH}_2\text{O}$), 43.8 ($-\text{CH}_2-$), 60.42 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 182.43\text{ Hz}$), 64.7 (2x CH_2O), 124.51 ($\text{CH}_5=\text{C}$), 112.4, 115.36, 139.18, 145.8 ($\text{C}_4\text{H}_3\text{O}$), 128, 128.47, 132.3, 132.6 (C_6H_5 and $\text{C}_4\text{H}_3\text{S}$), 141.2 ($\text{CH}_5=\text{C}$), 146.4 ($\alpha\text{-Furyl}-\text{CN}_4$), 167 (C=O).

Anal. calcd. for $\text{C}_{20}\text{H}_{23}\text{N}_8\text{O}_5\text{P}$: C, 49.38, H, 4.73, N, 23.04.

Found: C, 48.98, H, 4.66, N, 22.71.

Major isomer **8a**: Rf = 0.17 (AcOEt/DCM 1/1).

^1H NMR ($\text{CDCl}_3 + \text{DMSO } d_6$) δ : 1.03 (t, 3H, $J = 7\text{Hz}$), 1.21 (t, 3H, $J = 7\text{Hz}$), 3.65–4.27 (m, 2x2H), 5.85 (s, 2H), 6.19–6.28 (m, 1H), 6.86–6.87 (m, 1H), 6.97–6.98 (m, 1H), 7.14 (dd, 1H, $J_1 = 10\text{Hz}$, $J_2 = 16\text{Hz}$),

7.3–7.8 (m, 5H), 8.39 (s, 1H), 9.75 (dd, NH, $J_1 = 10\text{Hz}$, $J_3 = 4\text{Hz}$), 11.88 (m, NHpyrryl).

^{13}C NMR ($\text{CDCl}_3 + \text{DMSO } d_6$) δ : 16.49 ($2 \times \text{CH}_3\text{CH}_2\text{O}$), 43.42 ($-\text{CH}_2-$), 60.67 (d, $-\text{CH}-$, $^1J_{\text{C-P}} = 182.43 \text{ Hz}$), 64.22 ($2 \times \text{CH}_2\text{O}$), 110.38, 112.53, 114.32, 123.4 ($\text{C}_4\text{H}_4\text{N}$), 124.21 ($\text{CH}_5=\text{C}$), 128.35, 128.52, 132.5, 132.84 (C_6H_5), 141.56 ($\text{CH}_5=\text{C}$), 148.53 (α -Pyrryl- CN_4), 167.3 ($\text{C}=\text{O}$).

Anal. calcd. for $\text{C}_{20}\text{H}_{24}\text{N}_9\text{O}_4\text{P}$: C, 49.48, H, 4.95, N, 25.98.

Found: C, 49.14, H, 4.83, N, 25.95.

References

- [1] a) A. Elachqar, A. El Hallaoui, M. L. Roumestant, Ph. Viallefont, *Synthetic Comm.*, **24**, 1279 (1994).
b) S. Achamlale, A. Elachqar, A. El Hallaoui, M. L. Roumestant, Ph. Viallefont, *Aminoacids*, **12**, 257 (1997).
c) A. Alami, A. El Hallaoui, A. Elachqar, M. L. Roumestant, Ph. Viallefont, *Bull. Soc. Chim. Belg.*, **105**, 769 (1996).
d) A. Atmani, S. Elhajji, A. El Hallaoui, M. L. Roumestant, Ph. Viallefont, *Synthetic Comm.*, **21**, 2383 (1991).
e) F. Zaïd, *Thesis*, Fès, (1996).
- [2] P. Kafarski, B. Lejczak, *Phosphorus, Sulfur and Silicon*, **63**, 193 (1991).
- [3] B. Lejczak, P. Kafarski, H. Sztajer, P. Mastalerz, *J. Med. Chem.*, **29**, 2212 (1986).
- [4] V. P. Kukhar, N. M. Solodenko, V. A. Solodenko, *Ukr. BioKhim. Zh.*, **60**, 95 (1988).
- [5] a) W. H. W. Lunn, D. D. Schoepp, D. O. Calligaro, R. T. Vasileff, L. J. Heinz, C. R. Salhoff, P. J. O'Malley, *J. Med. Chem.*, **35**, 4608 (1992).
b) D. D. Schoepp, C. L. Smith, D. Lodge, J. D. Millar, J. D. Leander, A. I. Sacaan, W. H. W. Lunn, *European J. Pharmacology*, **203**, 237 (1992).
- [6] J. A. Monn, M. J. Valli, R. A. True, D. D. Schoepp, J. D. Leander, D. Lodger, *Bioorg. Med. Chem. Lett.*, **3**, 95 (1993).
- [7] A. Antonowa, S. Hauptmann, *Z. Chem.*, **16**, 17 (1976).
- [8] J. Elguero, C. Marzin, J. D. Roberts, *J. Org. Chem.*, **39**, 357 (1974).
- [9] S. Achamlale, A. Elachqar, A. El Hallaoui, S. Elhajji, A. Alami, M. L. Roumestant, Ph. Viallefont, *J. Heterocyclic Chem.*, in press.
- [10] L. Birkofer, A. Ritter, H. Uhlenbravick, *Chem. Ber.*, **96**, 3280 (1963).
- [11] G. I. Tsypin, T. N. Timofeeva, V. V. Mel'Nikov, B. V. Gidasov, *Zh. Org. Khim.*, **11**, 1395 (1975), and **13**, 2275 (1977).
- [12] M. Begtrup, *J. Chem. Soc. Perkin Trans. II*, 736 (1976).