

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

THE CHLORINATION REACTION OF O-ALKYL S-ALKYL(ARYL) THIOPHOSPHORIC(-NIC) ACID DERIVATIVES WITH PHOSPHORUS OXYCHLORIDE

Zheng-Jie He^a; Wen-Bing Chen^a; Fu-Peng^a; Zheng-Hong Zhou^a; Chu-Chi Tang^a

^a Institute of Elemento-Organic Chemistry, State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin, P.R. China

To cite this Article He, Zheng-Jie , Chen, Wen-Bing , Fu-Peng, Zhou, Zheng-Hong and Tang, Chu-Chi(1998) 'THE CHLORINATION REACTION OF O-ALKYL S-ALKYL(ARYL) THIOPHOSPHORIC(-NIC) ACID DERIVATIVES WITH PHOSPHORUS OXYCHLORIDE', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 142: 1, 49 – 55

To link to this Article: DOI: 10.1080/10426509808029665

URL: <http://dx.doi.org/10.1080/10426509808029665>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

THE CHLORINATION REACTION OF O-ALKYL S-ALKYL(ARYL) THIOPHOSPHORIC(-NIC) ACID DERIVATIVES WITH PHOSPHORUS OXYCHLORIDE

ZHENG-JIE HE, WEN-BING CHEN, FU-PENG MA,
ZHENG-HONG ZHOU and CHU-CHI TANG*

Institute of Elemento-Organic Chemistry, State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, P.R. China

(Received 15 March, 1998; Revised 26 June, 1998)

It is reported that a variety of O-alkyl S-alkyl(aryl) thiophosphoric(-nic) acid derivatives **4** can be readily chlorinated with phosphorus oxychloride giving S-alkyl(aryl) thiophosphoro(-no)chloridates **2** and O-alkyl phosphorodichloridates **3**.

Keywords: Chlorination; Phosphoro(-no)thiolate; Thiophosphoro(-no)chloridate; Phosphorus oxychloride

1. INTRODUCTION

In our laboratory, the isomerization/chlorination of a variety of O,O-dialkyl phosphoro(-no) thionates **1** with phosphorus oxychloride have been systematically studied. It was found that when R' equals aryloxy^[1], alkylthio^[2], arylthio^[2], dialkylamino^[3], phenyl^[4], methyl^[5], and nitrogen heterocyclic group^[6] in **1**, respectively, this reaction can proceed smoothly and gives the desired products **2** and **3**. Hence, it provides a general synthetic method for S-alkyl thiophosphoro(-no)chloridates, especially for the asymmetric ones.


During the course of investigating the isomerization/chlorination mechanism^[7], it was found that the isomerization product of **1**, O,S-dialkyl phosphorothiolate **4**, can be readily chlorinated with phosphorus oxychloride leading its one alkoxy to be replaced by a chlorine atom to give S-alkyl

* Correspondence author.




R = R'' = C₁₋₄ Alkyl, 2-Chloroethyl

R' = Alkoxy, Aryloxy, Alkylthio, Dialkylamino, Me, Ph,



thiophosphorochloridate **2** and O-alkyl phosphorodichloridate **3**. In this paper, we hope that the chlorination of **4** with phosphorus oxychloride can be developed a new convenient method for synthesis of S-alkyl thiophosphoro(-no)chloridate **2**, which is a key intermediate for preparation unsymmetric S-alkyl phosphoro(-no)thiolate possessing extensive biological activity. Because compound **4** is easily prepared from some cheap material, such as dialkoxy phosphite, this reaction is of some value to the synthesis.



2. RESULTS AND DISCUSSION

Phosphoro(-no)thiolate **4** reacted with equivalent amount of phosphorus oxychloride at 65–100°C until **4** disappeared from the reaction mixture (TLC control). After removal of by-product **3** under reduced pressure, the crude product **2** was purified by column chromatography on silica gel or by distillation under vacuum. The structures of product **2** were confirmed by IR, ¹H NMR spectra and elementary analyses (Table II). When R

equals alkyl or phenyl, R' is alkoxy, aryloxy, alkylthio, dialkylamino, nitrogen-containing heterocyclic group, methyl and phenyl, and R'' is C_{1-4} alkyl in compounds **4**, respectively, the chlorination of **4** can proceed smoothly giving the corresponding chlorinated products **2** in fair yields (Table I). Results show that the presence of alkylthio or arylthio RS group in **4** plays a critical role in the occurrence of this chlorination reaction. When trialkyl orthophosphoric acid ester, e.g. O,O,O-trimethyl phosphate, was treated with equivalent phosphorus oxychloride under a similar condition as **4a**, the desired product, O,O-dimethyl phosphorochloridate, was not obtained in a certain amount. In the previous literature^[8,9], it was reported that O-alkyl phosphoramidates or trialkyl phosphates give only pyrophosphoric acid derivatives by treatment of phosphorus oxychloride.

TABLE I The Chlorination Reaction of **4** with $POCl_3$ and Products **2**

Reactants						Products		
4	<i>R</i>	<i>R'</i>	<i>R''</i>	Reaction temp. (°C)	Reaction time (h)	2	n_D^{25}	Yield (%) [*]
a	Me	MeO	Me	75	2.5	a	1.4899	72.0
b	Pr	EtO	Et	80	5	b	1.4791	77.4
c	allyl	EtO	Et	65	10	c	1.4978	63.2
d	PhCH ₂	EtO	Et	85	5	d	1.5451	66.7
e	Ph	EtO	Et	100	5	e	1.5492	71.4
f	Et	PhO	Et	100	5	f	1.5508	42.3
g	Et	PhO	Pr	100	5	g	1.5590	71.5
h	Pr	PhO	Me	95	5	h	1.5436	85.5
i	Pr	PhO	Et	100	4	i	1.5440	55.9
j	Et	PhO	Bu	100	11	j	1.5580	74.7
k	Pr	2,4-ClBrC ₆ H ₃ O	Et	100	4	k	1.4770	43.9
l	Pr	MeS	Et	85	4	l	1.5490	42.8
m	Et	PrS	Et	75	5	m	1.5382	62.0
n	Et	Et ₂ N	Pr	100	6	n	1.4995	58.6
o	Ph	Et ₂ N	Et	100	5	o	1.5549	75.5
p	Pr	1-Piperidyl	Et	80	5	p	1.5189	66.7
q	Et	Me	Et	90	5	q	1.5042	75.7
r	Bu	Ph	Et	100	5	r	1.5120	62.5

* The isolated yields by column chromatography

TABLE II Data of 2 Prepared

2	IR(film), $\nu(\text{cm}^{-1})$		^1H NMR (CDCl_3/TMS) δ , J_{HH} (Hz)	Elementary Analyses			
	P=O	P-Cl		C%		H%	
				Calcd.	Found	Calcd.	Found
a	1223	591	2.50(d,3H,J=18.4), 3.90(d,3H,J=19.3)	14.95	14.52	3.91	3.74
b	1263	594	1.00(t,3H), 1.40(t,3H), 1.82(m,2H),	29.63	29.74	5.93	6.00
		2.92(dt,2H,J=16.8), 4.20(dq,2H,J=14.1)					
c	1267	587	1.35(t,3H), 3.64(dd,2H,J=18.1), 4.29(dq,2H,	29.92	29.79	4.98	5.13
		J=10.4), 5.22(d,2H), 5.88(m,1H)					
d	1262	588	1.30(t,3H), 4.11(dq,2H,J=14.2),	43.10	42.84	4.79	4.82
		4.31(d,2H,J=16.2), 7.30(m,5H)					
e	1266	596	1.35(t,3H), 4.29(dq,2H,J=14.3), 7.40(m,5H)	40.59	40.30	4.23	4.53
f	1264	592	1.45(t,3H), 3.00(dq,2H,J=16.1), 7.26(m,5H)	40.59	40.68	4.23	4.52
g	1265	592	1.44(t,3H), 2.96(dq,2H,J=16.2), 7.28(m,5H)	40.59	40.85	4.23	4.48
h	1260	587	0.98(t,3H), 1.78(m,2H), 3.08(dt,2H,J=18.7), 7.30(m,5H)	43.11	43.19	4.79	4.85
i	1261	586	0.98(t,3H), 1.88(m,2H), 3.09(dt,2H,J=18.5), 7.30(m,5H)	43.11	43.25	4.79	4.54
j	1265	593	1.44(t,3H), 2.95(dq,2H,J=16.2), 7.28(m,5H)	40.59	40.76	4.23	4.36
k	1267	590	1.02(t,3H), 1.80(m,2H), 3.18(dt,2H,J=18.7), 7.40(m,3H)	29.75	30.08	2.75	2.73
l	1226	587	1.00(t,3H), 1.74(m,2H), 2.48(d,3H,J=18.7), 2.98(dt,2H,J=18.7)	23.47	23.73	4.89	4.80
m	1249	579	1.02(t,3H), 1.44(t,3H), 1.80(m,2H), 3.05(m,4H)	27.46	27.52	5.49	5.48
n	1240	564	1.18(t,6H), 1.40(t,3H), 2.92(dq,2H,J=17.3), 3.24(dq,4H,J=14.4)	33.41	33.72	6.96	6.75
o	1249	559	0.97(t,3H), 3.17-3.30(m,4H), 7.36(m,3H), 7.58(m,2H)	45.54	45.39	5.69	5.47
p	1249	576	1.00(t,3H), 1.60(m,6H), 1.75(m,2H), 3.22(dt,2H,J=16.7), 3.26(m,4H)	39.75	39.38	7.04	6.90
q	1232	593	1.44(t,3H), 2.36(d,3H,J=16.2), 3.12(dq,2H,J=21.6)	22.71	22.89	5.04	5.45
r	1222	589	1.00(t,3H), 1.20-1.60(m,4H), 2.64(dt,2H,J=15.1), 7.48(m,5H)	48.29	48.01	5.63	5.75

TABLE III Data of Compounds 4 Prepared

4	bp(°C/Pa)	n_D^{25}	Yield(%)	1H NMR ($CDCl_3/TMS$) δ , J_{HH} (Hz)
a	78–80/266	1.4650	81.5	2.20(d,3H,J=15.1), 3.73(d,6H,J=12.6)
b	76–80/267	1.4560	76.4	0.93(t,3H), 1.27(t,2×3H), 1.65(m,2H), 2.76(m,2H), 4.08(m,2×2H)
c		1.4724	85.7	1.36(t,2×3H), 3.48(dd,2H,J=14.4), 4.18(dq,2×2H,J=8.6), 5.20(d,2H), 5.79(m,1H)
d	140–142/400	1.5216	67.4	1.25(t,2×3H), 4.00(d,2H,J=14.4), 4.04(dq,2×2H,J=12.8), 7.28(m,5H)
e	134–138/27	1.5136	93.5	1.28(t,2×3H), 4.18(dq,2×2H,J=8.3), 7.28–7.60(m,5H)
f	113–115/13.3	1.5165	69.9	1.28(t,3H), 1.32(t,3H), 2.85(dq,2H,J=15.8), 4.21(dq,2H, J=9.8), 7.38(m,5H)
g		1.5310	83.1	0.98(t,3H), 1.31(t,3H), 1.70(m,2H), 2.80(dq,2H,J=15.0), 4.10(dt,2H,J=9.8), 7.18(m,5H)
h	155–156/400	1.5235	39.5	1.02(t,3H), 1.72(m,2H), 2.89(dt,2H,J=13.5), 3.90(d,3H, J=10.7), 7.29(m,5H)
i	111–112/27	1.5165	49.7	0.90(t,3H), 1.33(t,3H), 1.63(m,2H), 2.83(m,2H), 4.25(m, 2H), 7.35(m,5H)
j		1.5305	79.2	0.82(t,3H), 1.10–1.90(m,4H), 1.32(t,3H), 2.82(dq,2H, J=14.8), 4.18(dt,2H,J=9.7), 7.18(m,5H)
k	110/0.13	1.5511	48.6	1.00(t,3H), 1.36(t,3H), 1.72(m,2H), 2.88(dt,2H,J=15.9), 4.25(dq,2H,J=9.8), 7.43(m,3H)
l	111–112/13.3	1.5036	60.0	1.01(t,3H), 1.37(t,3H), 1.73(m,2H), 2.33(d,3H,J=16.9), 2.87(dt,2H,J=16.2), 4.21(dq,2H,J=9.4)
m	128–130/267	1.4964	87.7	0.99(t,3H), 1.35(t,3H), 1.37(t,3H), 1.73(m,2H), 2.88(m, 4H), 4.19(dq,2H,J=9.8)
n		1.4705	75.3	0.90–1.18(m,9H), 1.30(t,3H), 1.68(m,2H), 2.70(dq,2H, J=14.8), 3.10(m,4H), 3.98(dt,2H,J=10.2)
o		1.5221	57.1	0.96(t,2×3H), 1.32(t,3H), 3.04(m,2×2H), 4.19(m,2H), 7.46(m,5H)
p	107–109/5.3	1.4916	66.4	1.00(t,3H), 1.32(t,3H), 1.56–1.84(m,8H), 2.80(dt,2H, J=14.0), 3.16(m,4H), 4.06(dq,2H,J= 11.9)
q	71–74/267	1.4726	78.7	1.32(t,3H), 1.40(t,3H), 1.78(d,3H,J=18.0), 2.90(dq,2H, J=12.2), 4.12(dq,2H,J=10.8)
r		1.5239	41.7	0.76(t,3H), 1.32(t,3H), 1.04–1.64(m,4H), 2.68(dt,2H, J=15.1), 4.19(dq,2H,J=9.2), 7.68(m,5H)

The chlorination reaction of phosphoro(-no)thiolates **4** can be performed under mild conditions. It also provides a new synthetic pathway for various S-alkyl(aryl) thiophosphoro(no)chloridates, particularly for chiral ones, which are often those important intermediates for synthesis of S-alkyl(aryl) thiophosphoric(-nic) acids derivatives probably possessing extensive biological activity. For example, the achiral S-propyl thiophosphate **4b**, $(\text{EtO})_2\text{P}(\text{O})\text{SPr}$, can be converted into the chiral S-propyl thiophosphorochloridate **2b**, $(\text{EtO})(\text{PrS})\text{P}(\text{O})\text{Cl}$, in a good yield (Table I), **2b** is a key synthetic intermediate for several excellent S-propyl O-ethyl O-aryl thiophosphates insecticides, such as Profenofos, Diphenprofos, Heterophos, and Pyraclofos.

EXPERIMENTAL

All temperatures are uncorrected. IR spectra were recorded on a NICOLET 5DX spectrophotometer as thin film. ^1H NMR spectra were measured on a JEOL FX-90Q instrument at 90 MHz, using TMS as internal standard and CDCl_3 as solvent. Elementary microanalysis data were determined with a Yanaco MT-3 instrument. For column chromatography, Qingdao silica gel (200~300 mesh) was used as a stationary phase. Phosphorus oxychloride was used after redistilled. Other reagents are commercial.

O-Alkyl S-alkyl(aryl) thiophosphoric(-nic) acid derivatives **4**

According to a general procedure described in a previous literature^[10], **4a** was prepared from the reaction of O,O-dimethyl ammonium thiophosphate with methyl iodide. Similarly, **4b~d**, **4f** and **4i** were prepared from reactions of their corresponding ammonium or sodium thiophosphates with appropriate alkyl halides RX (R is the same as in **4**, X is Cl, Br), respectively. O-Ethyl S-propyl ammonium phosphorodithioate and O-ethyl sodium phenylphosphonothioate reacted separately with butyl bromide giving **4m** and **4r**. According to an ordinary method, S-ethyl thiophosphoryl dichloride reacted with diethylamine or phenol to produce the corresponding thiophosphorochloridate, which in turn condensed with propanol or butanol giving **4g**, **4j** and **4n**. **4h**, **4k** and **4l** were prepared as

described in the literatures^[2,11,12], Using a known procedure^[13], triethyl phosphorous acid ester reacted with phenylsulfur chloride to afford **4e**, which further reacted with phosphorus oxychloride followed by condensation with diethylamine to give **4o**. Compounds **4** prepared above were purified by distillation under reduced pressure or by column chromatography on silica gel. Their data are listed in Table III.

The chlorination reaction of thiophosphoric(-nic) acids derivatives **4 with phosphorus oxychloride (General procedure)**

A mixture of **4** and phosphorus oxychloride (equiv.) was heated with stirring at 65~100°C for 2.5~11 h until **4** disappeared from the reaction mixture (TLC control, petroleum ether / ethyl acetate 5 : 1 as eluent, iodine as detecting reagent). After the reaction was complete, by-product **3** was removed by distillation under reduced pressure. The crude product **2** was purified by vacuum liquid chromatography on silica gel (petroleum ether / ethyl acetate, gradient elution) to give pure **2** (Table I, II).

Acknowledgements

The authors wish to thank Natural Science Foundation of Tianjin City and State Key Laboratory of Elemento-Organic Chemistry for financial support.

References

- [1] C.C. Tang, G.P. Wu, G.Z. Zhang, *Synthesis*, 1991, 454.
- [2] C.C. Tang, G.P. Wu, *Chem. J. Chinese Univ.*, 1993, **14**, 642.
- [3] C.C. Tang, G.P. Wu, G.Y. Huang, Z. Li, G.Y. Jin, *Phosphorus, Sulfur and Silicon*, 1993, **84**, 159.
- [4] C.C. Tang, G.P. Wu, S.J. He, Z.J. He, *ibid*, 1995, **101**, 91.
- [5] C.C. Tang, F.P. Ma, K. Zhang, Z.J. He, Y.C. Jin, *Heteroatom Chem.*, 1995, **6**, 413.
- [6] C.C. Tang, H.F. Lang, Z.J. He, *ibid*, 1996, **7**, 207.
- [7] C. C. Tang, F. P. Ma, Z. J. He, *Chem. J. Chinese Univ.*, 1997, **18**, 229.
- [8] H. Tolkmith, U.S. Patent 2654780 (1953).
- [9] E. Cherbuliez, G. Cordahi, J. Rabinowitz, *Helv. Chim. Acta*, 1959, **42**, 590.
- [10] C.C. Tang, G.P. Wu, Y.X. Chai, *Chem. J. Chinese Univ.*, 1981, **2**, 70.
- [11] C.C. Tang, F.P. Ma, M.J. Zhang, Z. Li, *Chem. J. Chinese Univ.*, 1994, **15**, 1327.
- [12] C.C. Tang, L.Z. Liu, G.P. Wu, Z. Li, Chinese Patent 94 108485 (1994).
- [13] N.M. Yousif, K.Z. Gadalla, S.M. Yassin, *Phosphorus, Sulfur and Silicon*, 1991, **60**, 261.