

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### A CONVENIENT METHOD FOR BROMOSULFENYLATION: REACTIONS OF SULFENAMIDES WITH OLEFINS IN THE PRESENCE OF $\text{POBr}_3$

Nikolai V. Zyk<sup>a</sup>; Elena K. Beloglazkina<sup>a</sup>; Rimma Gazzaeva<sup>b</sup>; Vladimir S. Tyurin<sup>a</sup>; Igor D. Titanyuk<sup>a</sup>

<sup>a</sup> Department of Chemistry, Moscow State University, Moscow, Russia <sup>b</sup> Hetagurov North-Osetia State University, Vladikavkaz, Russia

**To cite this Article** Zyk, Nikolai V. , Beloglazkina, Elena K. , Gazzaeva, Rimma , Tyurin, Vladimir S. and Titanyuk, Igor D.(1999) 'A CONVENIENT METHOD FOR BROMOSULFENYLATION: REACTIONS OF SULFENAMIDES WITH OLEFINS IN THE PRESENCE OF  $\text{POBr}_3$ ', Phosphorus, Sulfur, and Silicon and the Related Elements, 155: 1, 33 — 45

**To link to this Article:** DOI: 10.1080/10426509908044968

**URL:** <http://dx.doi.org/10.1080/10426509908044968>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## A CONVENIENT METHOD FOR BROMOSULFENYLATION: REACTIONS OF SULFENAMIDES WITH OLEFINS IN THE PRESENCE OF $\text{POBr}_3$

NIKOLAI V. ZYK<sup>a\*</sup>, ELENA K. BELOGLAZKINA<sup>a</sup>,  
RIMMA GAZZAEVA<sup>b</sup>, VLADIMIR S. TYURIN<sup>a</sup> and  
IGOR D. TITANYUK<sup>a</sup>

<sup>a</sup>*Department of Chemistry, Moscow State University, Moscow, Russia 119899 and*

<sup>b</sup>*Hetagurov' North-Osetia State University, Kutuzova str., 60-3-113, Vladikavkaz,  
Russia 362000*

*(Received 03 November, 1998; In final form 05 January, 1999)*

A general method for the one-pot transformation of alkenes into abromoalkyl-arylsulfides has been proposed based on the sulfonylation reaction by means of sulfenamides in the presence of phosphorus oxibromide. The plausible reaction mechanism and the results of reactions with a number of model alkenes such as cyclohexene, 1-heptene, norbornene and other from the bicyclo[2.2.1]heptane series are discussed.

**Keywords:** sulfenamides; sulfonylation; phosphorus oxohalides

### INTRODUCTION


The sulfonylation reactions, the processes of electrophilic incorporation of the RS fragment to a molecule, are known for more than 100 years<sup>[1]</sup>. The most useful as sulfonylating agents are sulfonyl halides; other derivatives of sulfenic acid, such as sulfonyl carboxylates, sulfenamides and sulfonates are less frequently used<sup>[2-6]</sup>. Sulfonyl halide addition to  $\text{C}=\text{C}$  double bond is a classical model AdE reaction. There is a variety of works dealing with investigation of the mechanism and products of these reactions<sup>[7]</sup>. However, with some exceptions, all works on the issue are con-

\* Correspondence Author.

cerned with arylsulfenyl chlorides, and the data about other haloanhydrides of sulfenic acid are presented by few articles. Perhaps, this selection is due to availability and higher stability of sulfenyl chlorides as compared with the corresponding fluorides<sup>[6]</sup>, iodides<sup>[9]</sup>, and, to a lesser degree, bromides<sup>[10]</sup>.

## RESULTS AND DISCUSSION

We have found that arylsulfenamides can react with POBr<sub>3</sub> at -40 °C to form complexes which are capable of electrophilic addition to alkene double bonds. After the phosphorus-containing products were removed on a silica gel column, the reaction mixture was identified as the bromosulfenylation product, based on the <sup>1</sup>H NMR spectra. The products were further purified by preparative TLC method.



1a, Ar = C<sub>6</sub>H<sub>5</sub>, R<sub>2</sub> = (CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>;

1b, Ar = 2-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, R<sub>2</sub> = (CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>;


1c, Ar = 4-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, R<sub>2</sub> = (CH<sub>2</sub>)<sub>2</sub>O(CH<sub>2</sub>)<sub>2</sub>;

1d, Ar = 4-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, R = CH<sub>3</sub>

The structures and yields of products obtained are presented in table I.

The same reaction of sulfenamides with olefins, activated by POCl<sub>3</sub>, give chlorosulfides which are the analogs of arylsulfenyl chloride addition products.

The reaction is the alternative to the ArS-Hal addition to olefins in its result. Nevertheless, the details of the reaction proceeding allow us to reject the sulfenyl halide intermediate formation in the process under investigation. First, sulfenyl halide formation upon mixing of POBr<sub>3</sub> and sulfenamide in the absence of olefin was not detected by means of TLC. Second, sulfenyl halides addition to olefins, as a rule, is a slow process (taking several hours at room-temperature or upon heating) and is accom-



panied by formation of considerable amounts of diaryldisulfides, which are products of radical degradation of sulfenylating agents<sup>[11]</sup>. At the same time, the reaction was completed within several minutes at  $-40\text{ }^{\circ}\text{C}$  and the formation of by-products was virtually negligible.

The reagent, arylsulfenamide-POBr<sub>3</sub> complex, is stable at room temperature for several hours in the CH<sub>2</sub>Cl<sub>2</sub> solution. Complex 1b-POBr<sub>3</sub> was isolated as the spreading solid after mixing of reactant solutions in THF. In the CH<sub>2</sub>Cl<sub>2</sub> solution, the reagent slowly degrades to form diaryldisulfide, Br<sub>2</sub> and O=P(NR<sub>2</sub>)Br<sub>2</sub>.

The structure of the formed complex was examined by <sup>13</sup>P NMR method. In the spectrum of sulfenamide 1c and POBr<sub>3</sub> mixture additional signals were found (-50.2 and -28.0 ppm) besides the POBr<sub>3</sub> signal at -101.3 ppm. In another synthesis using POBr<sub>3</sub> and morpholine it has been proved that the peak at -28.0 ppm is the signal due to O=P(NF<sub>2</sub>)Br<sub>2</sub>. The signal at -50.2 disappeared after olefin addition to the reaction mixture, thus being attributed to the sulfonylation complex.

We assume that, first, the reaction proceeds as acid-base interaction between POBr<sub>3</sub> as a Lewis acid and the nitrogen atom of sulfenamide, yielding B or C (see chart 1). Using AM-1 method the structure of the phenylthiomorpholine-POBr<sub>3</sub> complex was calculated. The structure with the optimised geometry was shown to be close to C.

Further, the complete substitution of a bromine atom at the phosphorus atom takes place, forming the ammonium salt D. After this, intermediate D can add to olefin giving trans-1,2-bromo-sulfide. The following finding testifies that it is the D species which attacks the olefin: calculated charges on the sulfur atom are 0.059 and 0.064 for structures B and C, respectively, which are nearly the same as that for the starting sulfenamide (0.052). Cal-





CHART 1


culuation made for structure D gives the much greater positive charge on sulfur, namely 0.182 and, which is more surprising, the sulfur (not nitrogen) atom is the electrophilic site of the molecule. Moreover, the semi-empirical calculations show that LUMO is localised on the O-P-Br fragment for all of the structures B, C, D and, therefore, the reaction under consideration is not an orbital-controlled process.

Cyclohexene is a convenient model for studying the stereochemistry of addition reactions. The addition of sulfenamides **1a-d** to this olefin in the presence of  $\text{POBr}_3$  have been found to occur stereoselectively and results in trans-1,2-adducts, in agreement with the electrophilic addition mechanism. The broadening of the  $^1\text{H}$  NMR signals for  $\alpha$ -protons at both arylthio- and bromo- groups on transition from nonpolar ( $\text{C}_6\text{D}_6$ ) to more polar ( $\text{CDCl}_3$ ) solvent unequivocally suggests the trans-configuration of bromosulfide **1c**.




1-Heptene as a terminal olefin, was chosen for determination of reaction regioselectivity. It is known that addition of sulfenyl chlorides to terminal olefins usually proceed without regioselectivity, with the formation of both possible products. The addition products formed in accordance with the Markovnikov rule is assumed to be more thermodynamically stable. In the same time, the anti-Markovnikov product is formed under the conditions

reaction kinetically controlled <sup>[12]</sup>. After the reaction of sulfenamide **1a-d** with 1-heptene at  $-40^{\circ}\text{C}$ , the mixture of both Markovnikov **3a-c** and anti-Markovnikov **4a''c** products has been isolated. In the case of compounds **3c** and **4c** their NMR spectra were measured again one month after their preparation and only the adduct **3c** (Markovnikov's) was found. This means that compounds **3** and **4** can be transformed into each other.



**3a, 4a**, Ar = C<sub>6</sub>H<sub>5</sub>, **3b, 4b**, Ar = 2-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, **3c, 4c**, Ar = 4-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>


To study the addition to norbornene was interesting for determination of the reagent's effective electrophilicity, the ability to cause a carbon skeleton rearrangements and the other phenomena typical of carbocation processes<sup>[13]</sup>. In this reaction only formation of 1,2-addition adducts **5a-c** has been observed. Neither the reaction conducted in the presence of LiClO<sub>4</sub> nor the interaction in a more polar solvent (CH<sub>3</sub>CN) lead to formation of the rearrangement products. The attempt to isolate the conjugated addition products were made by means of Hg(OCOCH<sub>3</sub>)<sub>2</sub> adding to the reaction mixture. But this reaction was resulted in the formation only bromsulfide **5**.




**5a**, Ar = C<sub>6</sub>H<sub>5</sub>, **5b**, Ar = 2-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, **5c**, Ar = 4-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>

However, the products of the Wagner-Meerwein rearrangement have been obtained when an other olefin from the bicyclo[2.2.1]heptane series, namely, 3,6-dimethoxy-benzonorbornadiene, was put into reaction. The interaction of sulfenamides **1b** and **1c** with this olefin in the presence of POBr<sub>3</sub> results in the formation of corresponding bromosulfides, having the structure of 2,7-disubstituted norbornanes. The reduction of yields in these

cases, perhaps, connected with the formation of by-products of sulfenylation of the substrate aromatic ring, activated by the methoxy substitution.




Products of the conjugated addition we can obtain using as a substrate other olefin from the bicyclo[2.2.1]heptane series, 5,6-diendo-norborn-2-enedimethylcarboxylate. The own carbomethoxy group of the substrate in this example acts as external nucleophile and  $\gamma$ -lactones **7b, c** have been isolated as the products.



The stereochemistry of electrophilic agents addition to 5-methylenenorbornene has not been yet established. It is known that its sulfenylation by arylsulfenyl chlorides results in nortricyclene structure product, besides, the authors supposed the beginning endo-attack of electrophile to the endocyclic double bond as the most likely<sup>[14]</sup>. We have made a careful NMR investigation of the methylenenorbornene bromosulfenylation products with the use of the nuclear Overhauser effect (NOE). Under the irradiation of the HCS-proton signal (3.42 ppm) the significant increasing of H<sub>6</sub> (1.67 ppm) and H<sub>3</sub> (1.53 ppm) signals intensity and the absence of the response on the H<sub>7</sub> (2.02 and 1.50 ppm) were observed. The irradiation of the CH<sub>2</sub>Br-group multiple (3.37 ppm) results in the increasing of the H<sub>3</sub>H<sub>4</sub>

(1.49 ppm) and  $H_6$  (2 protons, 1.67 and 1.62 ppm) signals intensity. These data demonstrated that, at least in the this reaction, not *endo*, but *exo*-attack of sulfenylating species to norbornene double bond and the final formation of the nortricyclane structure take place.



**8b**, Ar = 2-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>, **8c**, Ar = 4-NO<sub>2</sub>-C<sub>6</sub>H<sub>4</sub>

TABLE I The products of arylsulfenamides reactions with alkenes in the presence of POBr<sub>3</sub>

Alkene	ArS-NR <sub>2</sub>	Product	Yield, %
Cyclohexene	<b>1a</b>	<b>2a</b>	72
	<b>1b</b>	<b>2b</b>	95
	<b>1c</b>	<b>2c</b>	80
	<b>1d</b>	<b>2c</b>	85
1-Heptene	<b>1a</b>	<b>3a + 4a</b>	81
	<b>1b</b>	<b>3b + 4b</b>	92
	<b>1c</b>	<b>3c + 4c</b>	95
	<b>1d</b>	<b>3c + 4c</b>	62
Norbornene	<b>1a</b>	<b>5a</b>	79
	<b>1b</b>	<b>5b</b>	99
	<b>1c</b>	<b>5c</b>	75
	<b>1d</b>	<b>5c</b>	76
2,6-dimethoxy-benzonorbornadiene	<b>1b</b>	<b>6b</b>	69
	<b>1c</b>	<b>6c</b>	45
5,6-diendo-norborn-2-ene-dimethyl-carboxylate	<b>1b</b>	<b>7b</b>	86
	<b>1c</b>	<b>7c</b>	98
5-methylene-norbornene	<b>1b</b>	<b>8b</b>	92
	<b>1c</b>	<b>8c</b>	78



In conclusion we would noted that the proposed method for alkenes bromosulfenylation via generating of high-reactive arylsulfenamide-POBr<sub>3</sub> complexes can be a convenient synthetical method. The stability of the starting compounds and the simplicity of the experimental procedure make this method a suitable one for practical use.

## EXPERIMENTAL

### General

The purity of the reaction products was monitored by TLC. The preparative separation of the reaction product was carried out by TLC with Silufol plates using ethyl acetate – petroleum ether (1:3) as the eluent. All solvents were dried by standart technics. NMR spectra were recorded in CDCl<sub>3</sub>. IR spectra were obtained in nujol.

### Reactions between arylsulfenamides and alkenes in the presence of POHal<sub>3</sub> (general procedure)

An equivalent amount of POHal<sub>3</sub> in absolute CH<sub>2</sub>Cl<sub>2</sub> was slowly added to a solution of sulfenamide in CH<sub>2</sub>Cl<sub>2</sub> at –40 °C and then 1.5-fold excess of alkene in the same solvent was added. The mixture was heated to the room temperature, passed through a column-filter with silica gel (h = 5 cm). After evaporation of the solvent the residue was chromatographed.

#### *trans*-2-Phenylthio-cyclohexylbromide (2a)

R<sub>f</sub> 0.92. <sup>1</sup>H NMR δ 7.5–7.15 (m, 5H), 4.28 (m, 1H), 3.52 (m, 1H), 2.45–0.90 (m, 8H). Anal. Calcd. for C<sub>12</sub>H<sub>15</sub>BrS: C 53.14; H 5.57. Found: C 53.09; H 5.70.

#### *trans*-2-(2-Nitrophenylthio)-cyclohexylbromide (2b)

R<sub>f</sub> 0.85. <sup>1</sup>H NMR δ 8.08 (m, 1H), 7.58 (m, 1H), 7.43 (m, 1H), 7.30 (m, 1H), 4.40 (m, 1H), 3.75 (m, 1H), 2.50–1.20 (m, 8H). Anal. Calcd. for C<sub>12</sub>H<sub>14</sub>BrNO<sub>2</sub>S: C 45.58; H 4.46; N 4.43. Found: C 45.53; H 3.98; N 4.00.

***trans-2-(4-Nitrophenylthio)-cyclohexylbromide (2c)***

R<sub>f</sub> 0.92. <sup>1</sup>H NMR δ 8.15 (d, 2H, J = 9 Hz), 7.38 (d, 2H, J = 9 Hz) 4.28 (m, 1H), 3.72 (m, 1H), 2.45–0.90 (m, 8H). <sup>13</sup>C NMR δ 22.28, 22.51, 28.66, 32.30, 50.43, 53.25, 123.70, 124.00, 128.4. Anal. Calcd. for C<sub>12</sub>H<sub>14</sub>BrNO<sub>2</sub>S: C 45.58; H 4.47; N 4.43. Found: C 45.43; H 4.58; N 4.28.

***1-Phenylthio-hexyl-2-bromide (3a) and 2-Phenylthio-hexyl-1-bromide (4a)***

R<sub>f</sub> 0.90. <sup>1</sup>H NMR (for mixture) δ 7.38–7.10 (m, 5H of comp. **3a** and **4a**), 3.98 (m, 1H of comp. **3a**), 3.47 (dd, 2H of comp. **3a**, J = 4.9, 13.9 Hz), 3.17 (dd, 2H of comp. **4a**, J = 9.6, 13.8 Hz), 3.12 (m, 1H of comp. **4a**), 2.08–1.10 (m, 8H of comp. **3a** and **4a**), 0.82 (q, 3H of comp. **3a** and **4a**, J = 7.0 Hz). <sup>1</sup>H NMR (for **3a**) δ 7.38–7.10 (m, 5H), 3.98 (m, 1H), 3.47 (m, 2H, J = 4.9, 13.9 Hz), 2.08–1.10 (m, 8H), 0.82 (q, 3H). Anal. Calcd. for C<sub>13</sub>H<sub>19</sub>BrS: C 54.36; H 6.67. Found: C 54.11; H 6.18.

***1-(2-Nitrophenylthio)-hexyl-2-bromide (3b) and 2-(2-Nitrophenylthio)-hexyl-1-bromide (4b)***

R<sub>f</sub> 0.94. <sup>1</sup>H NMR δ 8.15 (m, 1H of comp. **3b** and **4b**), 7.58 (m, 1H of comp. **3b** and **4b**), 7.43 (m, 1H of comp. **3b** and **4b**), 7.25 (m, 1H of comp. **3b** and **4b**), 4.07 (m, 1H of comp. **3b**), 3.55 (dd, 2H of comp. **3b**, J = 4.0, 10.2 Hz), 3.32 (m, 1H of comp. **4b**), 3.23 (dd, 2H of comp. **4b**, J = 3.8, 13.5 Hz), 2.50–1.20 (m, 8H of comp. **3b** and **4b**), 1.08 (q, 3H, of comp. **3b** and **4b**, J = 7.0 Hz). Anal. Calcd. for C<sub>13</sub>H<sub>18</sub>BrNO<sub>2</sub>S: C 46.99; H 5.46; N 4.21. Found: C 46.50; H 4.98; N 4.20.

***1-(4-Nitrophenylthio)-hexyl-2-bromide (3c) and 2-(4-Nitrophenylthio)-hexyl-1-bromide (4c)***

R<sub>f</sub> 0.90. <sup>1</sup>H NMR δ 8.12 (d, 2H of comp. **3c** and **4c**, J = 9.0 Hz), 7.38 (d, 2H of comp. **3c** and **4c**, J = 9.0 Hz), 4.05 (m, 1H of comp. **3c**), 3.63 (dd, 2H of comp. **3c**, J = 5.0, 13.9 Hz), 3.56 (dd, 2H of comp. **4c**, J = 3.5, 10.0 Hz), 3.51 (m, 1H of comp. **4c**), 2.15–1.20 (m, 8H of comp. **3c** and **4c**), 0.90 (q, 3H of comp. **3c** and **4c**, J = 7.0 Hz). Anal. Calcd. for C<sub>13</sub>H<sub>18</sub>BrNO<sub>2</sub>S: C 46.99; H 5.46; N 4.21. Found: C 47.18; H 5.50; N 4.21.

***exo-2-Phenylthio-endo-3-norbornylbromide (5a)***

$R_f$  0.93.  $^1\text{H}$  NMR  $\delta$  7.25–6.95 (m, 5H), 3.88 (m, 1H), 3.00 (dd, 1H,  $J = 2.7, 4.0$  Hz), 2.51 (m, 1H), 2.45 (m, 1H), 2.30–1.14 (m, 6H). Anal. Calcd. for  $\text{C}_{13}\text{H}_{15}\text{BrS}$ : C 55.13; H 5.34. Found: C 55.14; H 5.48.

***exo-2-(2-Nitrophenylthio)-endo-3-norbornylbromide (5b)***

$R_f$  0.93.  $^1\text{H}$  NMR  $\delta$  8.10 (d, 1H,  $J = 7.5$  Hz), 7.50–7.05 (m, 3H), 4.05 (m, 1H), 3.15 (dd, 1H,  $J = 2.3, 4.2$  Hz), 2.60 (m, 1H), 2.48 (m, 1H), 2.37–1.10 (m, 6H). Anal. Calcd. for  $\text{C}_{13}\text{H}_{14}\text{BrNO}_2\text{S}$ : C 47.56; H 4.30; N 4.27. Found: C 48.00; H 4.29; N 3.99.

***exo-2-(4-Nitrophenylthio)-endo-3-norbornylbromide (5c)***

$R_f$  0.84.  $^1\text{H}$  NMR  $\delta$  8.01 (d, 2H,  $J = 8.8$  Hz), 7.20 (d, 2H,  $J = 8.8$  Hz), 3.95 (m, 1H), 3.24 (dd, 1H,  $J = 3.8, 8.0$  Hz), 2.46 (m, 1H), 2.40 (m, 1H), 2.30–1.14 (m, 6H). Anal. Calcd. for  $\text{C}_{13}\text{H}_{14}\text{BrNO}_2\text{S}$ : C 47.56; H 4.30; N 4.27. Found: C 47.78; H 4.33; N 4.24.

***3, 6-Dimethoxy-sin-1 1-(2-nitrophenylthio)-tricyclo[6.0.1.0<sup>2,7</sup>]undeca-2(7), 3, 5-trienyl-exo-9-bromide (6b)***

M.p. 60–61 °C, yellow crystals from heptane.  $R_f$  0.53.  $^1\text{H}$  NMR  $\delta$  8.09 (m, 1H), 7.51 (m, 2H), 7.25 (m, 1H), 6.15 (dd, 2H,  $J = 10.7, 16.0$  Hz), 3.99 (m, 1H), 3.78 (s, 1H), 3.76 (m, 1H), 3.73 (m, 1H), 3.72 (dd, 1H,  $J = 4.5, 8.0$  Hz), 3.65 (d, 1H,  $J = 4.5$  Hz), 2.79 (dt, 1H,  $J = 4.5, 14.0$  Hz), 2.17 (dd, 1H,  $J = 8.0, 14.0$  Hz).  $^{13}\text{C}$  NMR  $\delta$  36.58, 45.09, 45.64, 52.31, 55.58, 55.62, 61.92, 110.49, 110.73, 124.98, 125.70, 128.02, 132.90, 133.29, 134.20, 138.20, 147.26, 147.67. Anal. Calcd. for  $\text{C}_{19}\text{H}_{18}\text{BrNO}_4\text{S}$ : C 52.30; H 4.16; N 3.21. Found: C 52.69; H 4.07; N 3.30.

***3,6-Dimethoxy-sin-1 1-(4-nitrophenylthio)-tricyclo[6.0.1.0<sup>2,7</sup>]undeca-2(7), 3, 5-trienyl-exo-9-bromide (6c)***

$R_f$  0.40.  $^1\text{H}$  NMR  $\delta$  8.20 (d, 2H,  $J = 9.0$  Hz), 7.45 (d, 2H,  $J = 9.0$  Hz), 6.66 (d, 2H,  $J = 9.0$  Hz), 6.62 (d, 2H,  $J = 9.0$  Hz), 3.98 (m, 1H), 3.81 (m, 1H), 3.80 (s, 3H), 3.78 (s, 3H), 3.73 (m, 1H), 3.71 (m, 1H), 2.77 (dt, 1H,  $J = 4.0, 12.0$  Hz), 2.20 (dd, 1H,  $J = 7.0, 12.0$  Hz). Anal. Calcd. for  $\text{C}_{19}\text{H}_{18}\text{BrNO}_4\text{S}$ : C 52.30; H 4.16; N 3.21. Found: C 52.42; H 4.12; N 3.64.

***endo-3-Methoxycarbonyl-exo-5-(2-nitrophenylthio)-bicyclo[2.2.1]heptan-2-carboxylic acid  $\gamma$ -lactone (7b)***

M.p. 115–117 °C, yellow crystals from heptane.  $R_f$  0.06.  $^1\text{H}$  NMR  $\delta$  8.12 (m, 1H), 7.98 (m, 2H), 7.55 (m, 1H), 4.89 (dd, 1H,  $J = 3.2, 5.6$  Hz), 4.76 (d, 1H,  $J = 3.2$  Hz), 3.50 (s, 3H), 3.48 (dd, 1H,  $J = 5.6, 14.4$  Hz), 2.89 (dd, 1H,  $J = 5.6, 14.4$  Hz). IR 1790 (C=O, lactone), 1730 (COOMe). Anal. Calcd. For  $\text{C}_{16}\text{H}_{15}\text{NO}_6\text{S}$ : C 55.01; H 4.33; N 4.01. Found: C 55.31; H 4.27; N: 4.24.

***endo-3-Methoxycarbonyl-exo-5-(4-nitrophenylthio)-bicyclo[2.2.1]heptan-2-carboxylic acid  $\gamma$ -lactone (7c)***

M.p. 147–149 °C, yellow crystals from heptane (lit. m.p. 146 °C <sup>[15]</sup>).  $R_f$  0.09.  $^1\text{H}$  NMR  $\delta$  8.23 (d, 2H,  $J = 9.0$  Hz), 7.53 (d, 2H,  $J = 9.0$  Hz), 4.77 (dd, 1H,  $J = 3.2, 5.7$  Hz), 4.28 (d, 1H,  $J = 2.5$  Hz), 3.83 (s, 3H), 3.57 (dd, 1H,  $J = 2.5, 5.7$  Hz), 3.37 (dd, 1H,  $J = 3.6, 10.0$  Hz), 3.07 (dd, 1H,  $J = 5.5, 10.0$  Hz). IR 1800 (C=O, lactone), 1745 (COOMe). Anal. Calcd. for  $\text{C}_{16}\text{H}_{15}\text{NO}_6\text{S}$ : C 55.01; H 4.33; N 4.01. Found: C 55.36; H 4.43; N 4.33,

***2-Bromomethyl-5-(2-nitrophenylthio)-tricyclo[2.2.1.0<sup>[2,6]</sup>]heptane (8b)***

$R_f$  0.94.  $^1\text{H}$  NMR  $\delta$  3.42 (bs, 1H), 3.37 (m, 2H), 2.25 (bs, 1H), 2.02 (d, 1H,  $J = 10.8$  Hz), 1.67 (d, 1H,  $J = 10.8$  Hz), 1.62 (d, 1H,  $J = 10.8$  Hz), 1.53 (d, 1H,  $J = 5.3$  Hz), 1.50 (d, 1H,  $J = 10.8$  Hz), 1.49 (d, 1H,  $J = 5.3$  Hz).  $^{13}\text{C}$  NMR  $\delta$  21.67, 24.68, 28.40, 31.99, 36.03, 36.26, 36.58, 51.25, 124.39, 126.01, 127.43, 132.95, 137.38, 146.56. Anal. Calcd. for  $\text{C}_{14}\text{H}_{15}\text{BrNO}_2\text{S}$ : C 49.28; H 4.43; N 4.10. Found: C 49.08; H 3.91; N 4.39.

***2-Bromomethyl-5-(4-nitrophenylthio)-tricyclo[2.2.1.0<sup>[2,6]</sup>]heptane (8c)***

$R_f$  0.88.  $^1\text{H}$  NMR  $\delta$  7.95 (d, 2H,  $J = 9.0$  Hz), 7.25 (d, 2H,  $J = 9.0$  Hz), 3.83 (dd 2H,  $J = 5.1, 11.0$  Hz), 3.66 (bs, 1H), 2.30 (s, 1H), 1.90–1.37 (m, 6H). Anal. Calcd. for  $\text{C}_{14}\text{H}_{15}\text{BrNO}_2\text{S}$ : C 49.28; H 4.43; N 4.10. Found: C 49.55; H 3.98; N 4.00.

***trans-2-(4-Nitrophenylthio)-cyclohexylchloride (9)***

M.p. 68 °C, yellow crystals from heptane (lit. m.p. 66 °C <sup>[16]</sup>).  $R_f$  0.80.  $^1\text{H}$  NMR  $\delta$  8.02 (d, 2H,  $J = 9.0$  Hz), 7.34 (d, 2H,  $J = 9.0$  Hz), 3.99 (m, 1 H), 3.60 (m, 1 H), 2.50–1.30 (m, 8H).

***exo-2-(4-Nitrophenylthio)-endo-3-norbornylchloride (10)***

$R_f$  0.80.  $^1\text{H}$  NMR  $\delta$  8.05 (d, 2H,  $J = 9.0$  Hz), 7.25 (d, 2H,  $J = 9.0$  Hz), 3.95 (m, 1H), 3.15 (m, 1H), 2.60–1.12 (m, 8H). Anal. Calcd. for  $\text{C}_{13}\text{H}_{14}\text{ClNO}_2\text{S}$ : C 55.02; H 4.97; N 4.94. Found: C 54.80; H 4.88; N 4.66.

***Acknowledgements***

We wish to thank the Russian Foundation of Basic Research for the financial support (Grant N 96–03–32570).

***References***

- [1] (a) E. Kuhle, *Synthesis*, **N 11**, 561 (1970).  
 (b) E. Kuhle, *Synthesis*, **N 11**, 563 (1971).  
 (c) E. Kuhle, *Synthesis*, **N 12**, 617 (1971).
- [2] (a) N. S. Zefirov, N. K. Sadovaja, A. M. Magerramov, I. V. Bodrikov, V. R. Kartashov, *Tetrahedron*, **31**, 2948 (1975).  
 (b) N. S. Zefirov, N. K. Sadovaja, L. A. Novgorodtseva, I. V. Bodrikov, *Zh. Org. Khim.*, **14**, 1806 (1978).  
 (c) V. N. Ermolaeva, N. K. Sadovaja, N. S. Zefirov, *Zh. Org. Khim.*, **17**, 1554 (1981).
- [3] (a) P. Brownbridge, *Tetrahedron Lett.*, **25**, 3779 (1984).  
 (b) R. A. Abramovitch, J. Pilski, *J. Chem. Soc., Chem. Comm.*, **N 14**, 704 (1981).  
 (c) M. C. Caserio, J. K. Kim, *J. Am. Chem. Soc.*, **104**, 3231 (1982).  
 (d) L. Benati, P. C. Montevecchi, P. Spagnolo, *Tetrahedron Lett.*, **25**, 2039 (1984).  
 (e) N. S. Zefirov, N. V. Zyk, A. G. Kutateladze, E. K. Butina, *Zh. Org. Khim.*, **26**, 671 (1990).  
 (f) N. S. Zefirov, N. V. Zyk, A. G. Kutateladze, E. K. Butina, E.K. *Izv. Akad. Nauk SSSR, Ser. Khim.*, **N 12**, 2870 (1989).
- [4] (a) A. J. Havlik, N. Kharasch, *J. Am. Chem. Soc.*, **78**, 1207 (1956).  
 (b) L. Field, P. M. Giles, D. L. Tuleren, *J. Org. Chem.*, **36**, 623 (1971).  
 (c) P. A. Bell, D. R. Hogg, A. Robertson, *J. Chem. Soc., P-1*, 1246 (1978).  
 (d) B. M. Trost, M. Ochaiai, P. G. Mc.Dougal, *J. Am. Chem. Soc.*, **100**, 7103 (1978).
- [5] G. Capozzi, G. Modena and L. Pasquato, *The Chemistry of Functional Groups. The Chemistry of Sulphenic Acids and their Derivatives*. (Ed. S.Patai) Wiley, London, 1990, 403–516.
- [6] G.H. Schmidt and D. G. Garratt *The Chemistry of Functional Groups. The Chemistry of Double-Bonded Functional Groups. Supplement A. Part 2* (Ed.s. Patai) Wiley, London 1977, 828.
- [7] For selected examples, see: (a) V. A. Smit, N. S. Zefirov, I. V. Bodrikov, M. Z. Krimer, *Acc. Chem. Res.*, **12**, 282 (1979).  
 (b) N. S. Zefirov, N. K. Sadovaja, L. A. Novgorodtseva, I. V. Bodrikov, *Tetrahedron*, **N 9**, 1373 (1978).  
 (c) N. S. Zefirov, N. K. Sadovaja, L. A. Novgorodtseva, I. V. Bodrikov, *Zh. Org. Khim.*, **14**, 463 (1978).
- [8] (a) F. Seel, W. Gombler, R. Budenz, *Angew. Chem.*, **79**, 686 (1967).  
 (b) F. Seel, W. Gombler, R. Budenz, *Angew. Chem.*, **81**, 789 (1969).  
 (c) E. Cober, *J. Am. Chem. Soc.*, **81**, 4810 (1959).
- [9] (a) H. Rheinboldt, E. Motzkus, *Ber. Dtsch. Chem. Ges.*, **72**, 657 (1939).  
 (b) A. Burawoy, S. S. Mistry, *J. Chem. Soc.*, 3877 (1959).
- [10] (a) N. Kharasch, C. M. Buess, G. I. Gleason, *J. Am. Chem. Soc.*, **72**, 1796 (1950).  
 (b) M. Huerzeler, B. Berhet, A. Vasella, *Helv. Chim. Acta*, **75**, 557 (1992).  
 (c) J. Michalski, B. Iiszka-Krawiecka, A. Skowranska, *Roczniki Chem.*, **37**, 1963 (1963).

- (d) S. Kato, S. Min, Beniya, M. Ishida, *Tetrahedron Lett*, **28**, 6473 (1987).  
(e) N. Kharasch, R. B. Langford, *rg. Chem.*, **28**, 1903 (1963).
- [11] I. L. Knunyants, I. N. Rojkov, *Izv. Akad. Nauk SSSR, Ser. Khim.*, N 10, 2264 (1970).
- [12] (a) L. P. Rasteikene, D. I. Greitchute, M. T. Linkova, I. A. Knunyants, *Uspekhi Khimii*, **46**, 041 (1977).  
(b) W H. Mueller, P. E Bulter, *J. Am. Chem. Soc.*, **90**, 2075 (1968).  
(c) C. L. Dean, D. G. Garratt, T. T. Tichvell, G. H. Schmid, *J. Am Chem. Soc.*, **69**, 4958 (1947).
- [13] (a) See ref. 5(a).  
(b) N. S. Zefirov, V. A. Smit, I. V Bodrikov, M. Z. Krimer, *M.Z Doklady Akad. Nauk SSSR*, **240**, 858 (1978).
- [14] N S. Zefirov, N. K. Sadovaja, R. Sh. Ahmedova, I V. Bodrikov, *Zh. Org. Chim*, **15**, 217 (1979).
- [15] A. G Kutateladze, *Ph. D. Thesis*, Moscow, Moscow State University, 1989.
- [16] W H Mueller, P E. Butler, *J. Org. Chem.*, **33**, 1533 (1968).