

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### A Flexible Route to Mannose 6-Phosphonate Functionalized Derivatives

Sébastien Vidal<sup>a</sup>; Jean-Louis Montero<sup>a</sup>; Alain Leydet<sup>a</sup>; Alain Morère<sup>a</sup>

<sup>a</sup> Laboratoire de Chimie Biomoléculaire, Ecole Nationale Supérieure de Chimie de Montpellier, Université Montpellier II, Montpellier, France

Online publication date: 27 October 2010

**To cite this Article** Vidal, Sébastien , Montero, Jean-Louis , Leydet, Alain and Morère, Alain(2010) 'A Flexible Route to Mannose 6-Phosphonate Functionalized Derivatives', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 177: 10, 2363 – 2377

**To link to this Article:** DOI: 10.1080/10426500214297

**URL:** <http://dx.doi.org/10.1080/10426500214297>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



## A FLEXIBLE ROUTE TO MANNOSE 6-PHOSPHONATE FUNCTIONALIZED DERIVATIVES

Sébastien Vidal, Jean-Louis Montero, Alain Leydet,  
and Alain Morère  
Laboratoire de Chimie Biomoléculaire, Université  
Montpellier II, Ecole Nationale Supérieure de Chimie de  
Montpellier, Montpellier, France

(Received March 19, 2002)

*A new approach for the synthesis of a mannose 6-phosphonate isosteric analog of mannose 6-phosphate is reported. The mannosylphosphonate has been prepared in a multistep synthesis involving an homologation reaction of the methyl  $\alpha$ -D-mannopyranoside followed by an Arbuzov reaction between a bromohomomannosyl derivative and the tris(trimethylsilyl)phosphite. This approach, avoiding the deprotection of dialkylphosphonate, allowed us to prepare the mannose 6-phosphonate in good yield. The described method was successfully extended to the preparation of a mannose 6-phosphonate linked to a cholesteryl moiety. This strategy affords a more general route for a wide range of functionalized mannose 6-phosphonate derivatives.*

**Keywords:** Glycolipid; lectin; mannose 6-phosphate; mannose 6-phosphate/insulin-like growth factor; mannose 6-phosphonate/triethyleneglycolcholesteryl; phosphonation; *tris*(trimethylsilyl)phosphite

## INTRODUCTION


Mannose 6-Phosphate (M6P) is a recognition marker involved in the selective targeting of newly synthesized enzymes to lysosomes.<sup>1</sup> While two different receptors recognize specifically the M6P residues, only the larger M6P receptor mediates the endocytosis of extracellular

This research was supported by the Scientific Council of the Université Montpellier II and the Fondation pour la Recherche Médicale (to S.V.). We also wish to thank Mayoly-Spindler Laboratories (Chatou—France) for financial support.

Address correspondence to Alain Morère, Laboratoire de Chimie Biomoléculaire (UMR 5032), Université Montpellier II, Ecole Nationale Supérieure de Chimie de Montpellier, 8 Rue de l'Ecole Normale, F-34296 Montpellier Cedex 05, France. E-mail: morere@univ-montp2.fr

M6P-containing ligands.<sup>2</sup> This receptor is a type I glycoprotein of 275 kDa, which apart from M6P residues, is also known to bind retinoic acid<sup>3</sup> through distinct binding sites (RA) and insulin-like growth factor II<sup>4</sup> (IGF-II). This receptor is the first example of a protein able to bind three different classes of ligands, that is, a saccharide (M6P), a peptide (IGF-II), and a lipid (RA). Moreover, the M6P/IGFIIR plays a fundamental role in the control of cell growth in fetal development and carcinogenesis.<sup>6</sup> In our ongoing research program, we focus on the ability of the M6P/IGF-IIR to recognize and then internalize M6P analogs. We recently described<sup>7</sup> the synthesis of two mannose 6-phosphonate (M6Pn) analogs (**I** and **II**) of M6P (Figure 1), and evaluated their affinity towards the M6P/IGFIIR.

Isosteric compound **I** has been shown to have an affinity to the M6P/IGFIIR similar to natural M6P, whereas nonisosteric compound **II** was only weakly recognized. In our previous syntheses<sup>7,8</sup> of the mannosyl phosphonate **I**, the key reaction in the synthetic scheme was the last step, involving the deprotection of the diethylphosphonate moiety according to Rabinowitz's procedure.<sup>9</sup> However, the conversion of diethylphosphonates into the corresponding free phosphonic acids using the method of transesterification by  $\text{Me}_3\text{SiBr}$  can be troublesome. Indeed, degradation problems, partial deprotection, or even complete failure of the deprotection has been described in several cases, such as with aminoacids or glycopyranose frameworks.<sup>10–12</sup> Recently, these difficulties were met with the preparation of a glycoconjugate we called




**FIGURE 1** Phosphonate analogs of M6P.

M6Pn/CSA,<sup>13</sup> where M6Pn and CSA stand for mannose 6-phosphonate and cholesteryl-succinyl-anilinyI respectively (Figure 1). Partial deprotection of the diethylphosphonate of this molecule and partial cleavage of the anomeric bond and/or amide bonds were observed under Rabinowitz's procedure. It is worth pointing out that this amphiphilic steroidal mannose 6-phosphonate incorporated into the lipidic bilayer of a liposome<sup>13</sup> constitute an efficient drug delivery system in selective targeting of breast cancer cells.

## RESULTS AND DISCUSSION

In order to circumvent the Rabinowitz deprotection step in the synthesis of M6Pn **1** and its conjugates, we decided to introduce the phosphonate moiety by an Arbuzov reaction between *tris*(trimethylsilyl)phosphite and a halogenohomomannosyl derivative. This attractive approach could furnish the *bis*(trimethylsilyl)ester which can be easily hydrolyzed with saturated aqueous NaHCO<sub>3</sub> solution.<sup>14</sup> The key step of this alternative synthesis involves the preparation of a mannoheptose derivative by a one carbon chain extension at the C-6 position. Various procedures are described in the literature, and many reactants can be used such as cyanide anion,<sup>15–20</sup> triphenylmethoxymethylphosphonium salt,<sup>21</sup> or iron-carbonyl-cyclopentadienyl complexes.<sup>22,23</sup> Homologation can be also accomplished via alkyne<sup>24–28</sup> or epoxysilane<sup>29,30</sup> derivatives. Another method is possible using a cyclic sulfate intermediate.<sup>31</sup>


The preparation of the mannose 6-phosphonate was realized (Scheme 1) via an homologation reaction involving condensation of the 2-lithio-1,3-dithiane<sup>32</sup> with the triflate derivative of alcohol **1**.<sup>33</sup> Compound **2** was then reacted with ceric ammonium nitrate (CAN)<sup>34–36</sup> in wet acetonitrile affording a crude aldehyde (<sup>1</sup>H NMR analysis gave satisfactory data showing the resonance of the aldehydic proton at 9.75 ppm) which was reduced by sodium borohydride to furnish the homologated alcohol **3** in a one-pot procedure. Reactions proceed quickly affording expected compounds in good yields. Conversion of homoalcohol **3** to homoalkylbromide **5** was carried out by reaction of **3** with methanesulfonylchloride (affording **4** in 95% yield) followed by bromination with lithium bromide<sup>37,38</sup> in butanone, affording **5** (95%) without further purification. This two-step procedure was preferred to the one-step PPh<sub>3</sub>/CBr<sub>4</sub><sup>39</sup> method, in order to avoid problems with the removal of triphenylphosphine oxide byproduct by column chromatography. The phosphonate moiety was introduced by an Arbuzov reaction between *tris*(trimethylsilyl)phosphite and **5**, affording



**SCHEME 1** Reagents and conditions: (a)  $\text{TiF}_2\text{O}$ , 2,6-di-*tert*-butyl-4-methylpyridine,  $\text{CH}_2\text{Cl}_2$ ,  $-40^\circ\text{C}$ , 10 min (85%). (b) 1,3-dithiane, *n*-BuLi, HMPA, THF,  $-78^\circ\text{C}$ , 10 min (85%). (c) CAN, wet  $\text{CH}_3\text{CN}$ , rt, 15 min. (d)  $\text{NaBH}_4$ , EtOH/ $\text{H}_2\text{O}$ , rt, 5 min (75% for two steps). (e) MsCl, DIEA,  $\text{CH}_2\text{Cl}_2$ , rt, 15 min (95%). (f) LiBr, butanone,  $85^\circ\text{C}$ , 1 h (95%). (g)  $\text{P}(\text{OSiMe}_3)_3$ ,  $160^\circ\text{C}$ , 16 h (85%). (h)  $\text{H}_2/\text{Pd-C}$  10%, EtOH/ $\text{H}_2\text{O}$ , rt, 17 h then DOWEX 50WX2 ( $\text{Na}^+$  form) (90%).

a *bis*(trimethylsilyl)phosphonate, which was hydrolyzed with saturated aqueous  $\text{NaHCO}_3$  solution, giving the salt of the corresponding phosphonic acid **6**. Benzyl protecting groups were removed by hydrogenolysis over Pd-C (10%) to obtain the title compound **I** in good overall yield (29%).

We therefore decided to use the same strategy to prepare the M6Pn-based glycolipid **13** (Scheme 2) in which the hydrophilic carbohydrate head is linked to the hydrophobic steroidal tail through a triethyleneglycol (TEG) spacer arm. Compound **3** was deprotected to afford the mannoheptose derivative **7** which was activated at the 6-position followed by peracetylation<sup>32,33,40</sup> and subsequent bromination to afford **8** in 55% yield over three steps. The methyl glycoside **8** was then converted to the 1-*O*-acyl glycoside **9** then activated to provide a glycosyl donor intermediate **10**. Koenigs-Knorr glycosylation<sup>41,42</sup> of **10**, using known 8-(cholest-5-en-3 $\beta$ -yloxy)-3,6-dioxaoctan-1-ol<sup>43</sup> as the glycosyl acceptor, affords the orthoester derivative **11** as the major product, which upon subsequent treatment with a Lewis acid, underwent rearrangement<sup>44-49</sup> to furnish the desired *O*-glycoside **12**. These two isomers could easily be differentiated according to their respective NMR data (Figure 2 and Table I) and also by TLC.




**SCHEME 2** Preparation of M6Pn/TEGC **13**—reagents and conditions: (a)  $H_2/Pd-C$  10%, EtOH/ $H_2O$ , rt, 16 h, (95%). (b) TsCl,  $C_5H_5N$ , rt, 5 h. (c)  $Ac_2O$ ,  $C_5H_5N$ ,  $0^\circ C$  to rt, 1 h. (d) LiBr, 2-butanone,  $85^\circ C$ , 1 h, (55% over three steps). (e)  $Ac_2O/AcOH/H_2SO_4$ (cat.), rt, 24 h, (85%). (f) HBr/AcOH,  $CH_2Cl_2$ , rt, 16 h. (g) 8-(cholest-5-en- $3\beta$ -yloxy)-3,6-dioxaoctan-1-ol,  $CH_2Cl_2$ , AgOTf, *sym*-collidine,  $0^\circ C$  to rt, 15 min, (45% over both steps). (h) TMSOTf,  $CH_2Cl_2$ , molecular sieves (3Å),  $-5^\circ C$ , 5 min, (40%). (i)  $P(OSiMe_3)_3$ ,  $160^\circ C$ , 16 h then  $NaHCO_3$ (aq). (j) NaOMe, MeOH, rt, 16 h, (50% over both steps).

The final step in our synthesis involved an Arbuzov reaction between the alkyl bromide derivative **12** and *tris*(trimethylsilyl)phosphite, followed by deprotection of the acetate under Zemplén conditions affording the final M6Pn/TEGC **13** in good yields. Characterization of this glycolipid was troublesome but evidence of the structure was provided by NMR spectroscopy ( $^1H$  and  $^{31}P$ ) and also by mass spectrometric analyses.

**TABLE I** Comparison of Compounds **11** and **12** NMR Data

Compound	$J_{1-2}^a$	$\delta_{H-1}^b$	$\delta_C^c$	$\delta_{CH_3}^d$	$\delta_{CH_3}^e$
Orthoester <b>11</b>	2.6	5.44	124.7	—	1.71
Glycoside <b>12</b>	1.5	4.75	—	21.5	1.91

<sup>a</sup>Coupling constant, in Hz.<sup>b</sup><sup>1</sup>H NMR chemical shift, in ppm.<sup>c</sup><sup>13</sup>C NMR chemical shift for the quaternary carbon (see\* in Figure 2), in ppm.<sup>d</sup><sup>13</sup>C NMR chemical shift for the methyl (see † in Figure 2), in ppm.<sup>e</sup><sup>1</sup>H NMR chemical shifts for the methyl groups, in ppm.**FIGURE 2** Partial NMR analyses for compounds **11** and **12**. a) <sup>13</sup>C NMR in CDCl<sub>3</sub> shows the signal corresponding to the quaternary carbon (C\*) from the orthoester derivative **11** while b) does not display this peak for **12**. c) <sup>1</sup>H NMR in CDCl<sub>3</sub> shows two signals for the acetates and one more singlet corresponding to the orthoester-methyl (CH<sub>3</sub>†) group for **11** while d) displays the three signals corresponding to each acetyl groups for **12**.

## CONCLUSION

The compound **I** was obtained in good overall yield (29%), demonstrating the usefulness of the Arbuzov phosphonation—using the *tris*(trimethylsilyl)phosphite—for the preparation of glycopyranosylphosphonic acids, such as the isosteric analog (**I**) of mannose 6-phosphate. This strategy was successfully applied to the preparation of a steroidal derivative M6Pn/TEGC **13**. Preparation of liposomes incorporating M6Pn/TEGC **13** and evaluation of their interactions with MCF7 cells now are being investigated in our group with the aim of developing drug delivery systems. Although the overall yield is still very low (1.75% over 16 steps starting from methyl  $\alpha$ -D-mannopyranoside) for the preparation of M6Pn/TEGC **13**, this strategy provides a more general route to a wide variety of functionalized M6Pn where the anomeric center will be substituted with any biologically active molecule during the glycosylation.

## EXPERIMENTAL

Analytical TLC were performed using aluminum-coated TLC plates 60-F<sub>254</sub> (Merck). Plates were developed with: (1) UV light (254 nm), (2) immersion in a 10% H<sub>2</sub>SO<sub>4</sub>/EtOH solution followed by charring, (3) immersion in a 5% rhodanine/EtOH solution followed by charring (for aldehydes) or (4) immersion in a phosphomolybdic solution (for phosphorus containing compounds). Silica gel column chromatography was performed with silica gel 60A (Carlo Erba). Optical rotations were measured at the sodium D-line with a Perkin-Elmer-241 polarimeter. Fast Atom Bombardment (FAB) mass spectra were recorded on a Jeol JMS-DX300 spectrometer in either positive (FAB+) or negative (FAB-) modes and using either 3-nitrobenzyl alcohol (NBA) or glycerol/thioglycerol (1:1) mixture (G/T). <sup>1</sup>H NMR spectra were recorded on a Bruker DRX 400 (400 MHz), at 25°C. Chemical shifts ( $\delta$ ) are given in ppm and referenced using residual solvent signals (7.26 ppm for CHCl<sub>3</sub> and 4.79 ppm for HOD). The following abbreviations were used to explain the signal multiplicities or characteristics: s (singlet), d (doublet), dd (double doublet), t (triplet), td (triplet doublet), m (multiplet). <sup>13</sup>C NMR spectra were recorded on a Bruker DRX 400 (100.6 MHz). Chemical shifts ( $\delta$ ) are given in ppm relative to TMS as an external reference. <sup>31</sup>P NMR spectra were recorded on a Bruker DPX 200 (81.0 MHz). Chemical shifts ( $\delta$ ) are given in ppm relative to phosphoric acid (85%) as an external reference.



### Methyl 2,3,4-Tri-O-benzyl-6-deoxy-6-(1,3-dithianyl)- $\alpha$ -D-mannopyranoside **2**

To a solution of **1**<sup>33</sup> (1 g, 2.16 mmol) and 2,6-di-*tert*-butyl-4-methylpyridine (555 mg, 2.7 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added dropwise triflic anhydride (633  $\mu$ L, 2.37 mmol), at  $-40^{\circ}\text{C}$ . After stirring for 30 min, the reaction was neutralized with NaHCO<sub>3</sub> (50 mL, 1 g/100 mL H<sub>2</sub>O) and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2  $\times$  50 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane then hexane/ether 4:1) affording a pale pink oil (1.09 g, 85%). To a solution of 1,3-dithiane (706 mg, 5.87 mmol) dissolved in THF (2 mL) was added dropwise HMPA (300  $\mu$ L, 1.67 mmol). The mixture was cooled to  $-78^{\circ}\text{C}$  before addition of *n*-BuLi (5 mL, 5.87 mmol, 1.6 M in hexane). After 5 min, a solution of the triflate derivative previously prepared (1 g, 1.67 mmol) in THF (5 mL) was added dropwise at  $-78^{\circ}\text{C}$ . After 10 min, the reaction mixture was quenched with NH<sub>4</sub>Cl (50 mL, 1 g/100 mL H<sub>2</sub>O) and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2  $\times$  50 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ether 95:5 then 3:1) affording **2** as pale yellow oil (807 mg, 85%).  $R_f = 0.67$  (hexane/ether 2:3).  $[\alpha]_D^{20} = +20.8^{\circ}$  (*c* 0.72/CHCl<sub>3</sub>). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.80–2.20 (m, 3H, H-6 CH<sub>2</sub>CH<sub>2</sub>S), 2.30–2.50 (m, 1H, H-6'), 2.70–3.00 (m, 4H, CH<sub>2</sub>S), 3.41 (s, 3H, OCH<sub>3</sub>), 3.73 (t, 1H,  $J$  = 9.3 Hz, H-4), 3.84 (dd, 1H,  $J$  = 1.9 and 2.8 Hz, H-2), 3.90–4.05 (m, 2H, H-3 H-5), 4.32 (dd, 1H,  $J$  = 3.7 and 11.0 Hz, H-7), 4.65 (s, 2H, CH<sub>2</sub>Ph), 4.69 (d, 1H,  $J$  = 12.0 Hz, CH<sub>2</sub>Ph), 4.72 (d, 1H, H-1), 4.76 (d, 1H,  $J$  = 12.5 Hz, CH<sub>2</sub>Ph), 4.83 (d, 1H, CH<sub>2</sub>Ph), 5.01 (d, 1H, CH<sub>2</sub>Ph), 7.20–7.50 (m, 15H, 15H-ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 24.0, 24.3, 29.0, 36.5, 42.1, 54.0, 67.0, 70.0–73.0 (3C), 73.7, 77.7, 79.2, 98.1, 125.0–127.0 (15C), 137.0–137.5 (3C). MS (FAB+, NBA):  $m/z$ : 589 [M + Na]<sup>+</sup>, 567 [M + H]<sup>+</sup>, 535 [M–OCH<sub>3</sub>]<sup>+</sup>, 475 [M–PhCH<sub>2</sub>]<sup>+</sup>, 91 [PhCH<sub>2</sub>]<sup>+</sup>.

### Methyl 2,3,4-Tri-O-benzyl-6-deoxy- $\alpha$ -D-mannoheptopyranoside **3**

To a solution of **2** (440 mg, 0.78 mmol) dissolved in CH<sub>3</sub>CN/H<sub>2</sub>O (15:1, 16 mL) was added ceric ammonium nitrate (1.285 g, 2.34 mmol). After stirring at rt for 15 min, the reaction mixture was evaporated under reduced pressure before diluting with EtOAc (100 mL). The organic layer was washed with H<sub>2</sub>O (2  $\times$  75 mL) and the organic layer extracted with EtOAc (2  $\times$  100 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>),

filtered, and evaporated under reduced pressure. The crude oil was dissolved in absolute ethanol (10 mL) before addition of a solution of  $\text{NaBH}_4$  (75 mg, 1.95 mmol) in  $\text{EtOH}/\text{H}_2\text{O}$  (1:1, 20 mL). The reaction mixture was stirred at rt for 5 min before being neutralized with  $\text{HCl}$  (1 M) and then diluted with  $\text{H}_2\text{O}$  (100 mL). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 100$  mL). Organic layers were combined, dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (hexane/ether 7:3 then 1:1) affording **3** as pale yellow oil (278 mg, 75%).  $R_f = 0.44$  (hexane/ether 1:4).  $[\alpha]_{\text{D}}^{20} = +21.5^\circ$  ( $c$  0.65/ $\text{CHCl}_3$ ).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.80\text{--}2.30$  (m, 2H, H-6 H-6'), 3.35 (s, 3H,  $\text{OCH}_3$ ), 3.75–3.95 (m, 6H, H-2 H-3 H-4 H-5 H-7 H-7'), 4.65 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 4.66 (d, 1H,  $J = 10.9$  Hz,  $\text{CH}_2\text{Ph}$ ), 4.70 (d, 1H,  $J = 1.7$  Hz, H-1), 4.73 (d, 1H,  $J = 12.4$  Hz,  $\text{CH}_2\text{Ph}$ ), 4.82 (d, 1H,  $\text{CH}_2\text{Ph}$ ), 5.05 (d, 1H,  $\text{CH}_2\text{Ph}$ ), 7.30–7.50 (m, 15H, 15H-ar).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 34.2, 55.2, 61.6, 71.9, 72.5, 73.4, 74.9, 75.7, 78.6, 80.6, 99.5, 128.0\text{--}129.0$  (15C), 138.6–138.9 (3C). MS (FAB+, NBA):  $m/z$ : 501  $[\text{M} + \text{Na}]^+$ , 479  $[\text{M} + \text{H}]^+$ , 447  $[\text{M} - \text{OCH}_3]^+$ , 91  $[\text{PhCH}_2]^+$ .

### Methyl 2,3,4-Tri-*O*-benzyl-6-deoxy-7-*O*-methanesulfonyl- $\alpha$ -D-manno-heptopyranoside **4**

To a solution of **3** (250 mg, 0.52 mmol) dissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added DIEA (116  $\mu\text{L}$ , 0.68 mmol) and then methanesulfonyl chloride (50  $\mu\text{L}$ , 0.63 mmol). After stirring for 15 min at rt, the reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (150 mL) and the organic layer was washed with  $\text{NaHCO}_3$  (10% aqueous,  $2 \times 100$  mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated under reduced pressure affording **4** as a white oil (275 mg, 95%).  $R_f = 0.55$  (hexane/ether 1:4).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.80\text{--}2.05$  (m, 1H, H-6), 2.25–2.50 (m, 1H, H-6'), 2.95 (s, 3H,  $\text{CH}_3\text{SO}_2$ ), 3.34 (s, 3H,  $\text{OCH}_3$ ), 3.60–4.00 (m, 4H, H-2 H-3 H-4 H-5), 4.35–4.50 (m, 2H, H-7 H-7'), 4.65 (s, 2H,  $\text{CH}_2\text{Ph}$ ), 4.66 (d, 1H,  $J = 11.0$  Hz,  $\text{CH}_2\text{Ph}$ ), 4.71 (d, 1H,  $J = 1.8$  Hz, H-1), 4.73 (d, 1H,  $J = 12.3$  Hz,  $\text{CH}_2\text{Ph}$ ), 4.81 (d, 1H,  $\text{CH}_2\text{Ph}$ ), 5.01 (d, 1H,  $\text{CH}_2\text{Ph}$ ), 7.30–7.50 (m, 15H, 15H-ar).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 31.7, 37.8, 55.3, 67.0, 67.8, 72.5, 73.4, 75.0, 75.7, 78.7, 80.7, 99.5, 128.0\text{--}129.0$  (15C), 138.0–139.0 (3C). MS (FAB+, NBA):  $m/z$ : 579  $[\text{M} + \text{Na}]^+$ , 557  $[\text{M} + \text{H}]^+$ , 525  $[\text{M} - \text{OCH}_3]^+$ , 91  $[\text{PhCH}_2]^+$ .

### Methyl 2,3,4-Tri-*O*-benzyl-7-bromo-6,7-dideoxy- $\alpha$ -D-manno-heptopyranoside **5**

A solution of **4** (200 mg, 0.36 mmol) and  $\text{LiBr}$  (625 mg, 7.2 mmol) in 2-butanone (10 mL) was stirred at  $85^\circ\text{C}$  for 1 h. The reaction mixture

was cooled to rt then diluted with  $\text{CH}_2\text{Cl}_2$  (130 mL) and the organic layer was washed with  $\text{H}_2\text{O}$  ( $2 \times 100$  mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated under reduced pressure affording **5** as a pale yellow oil (185 mg, 95%).  $R_f = 0.68$  (hexane/ether 2:3).  $[\alpha]_{\text{D}}^{20} = +28.6^\circ$  ( $c$  0.21/ $\text{CHCl}_3$ ).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 2.00\text{--}2.25$  (m, 1H, H-6),  $2.35\text{--}2.50$  (m, 1H, H-6'),  $3.41$  (s, 3H,  $\text{OCH}_3$ ),  $3.86$  (dd, 1H,  $J = 1.8$  and  $3.1$  Hz, H-2),  $3.96$  (dd, 1H,  $J = 8.9$  Hz, H-3),  $3.55\text{--}3.90$  (m, 4H, H-4 H-5 H-7 H-7'),  $4.67$  (s, 2H,  $\text{CH}_2\text{Ph}$ ),  $4.69$  (d, 1H,  $J = 11.0$  Hz,  $\text{CH}_2\text{Ph}$ ),  $4.74$  (d, 1H, H-1),  $4.77$  (d, 1H,  $J = 12.4$  Hz,  $\text{CH}_2\text{Ph}$ ),  $4.85$  (d, 1H,  $\text{CH}_2\text{Ph}$ ),  $5.02$  (d, 1H,  $\text{CH}_2\text{Ph}$ ),  $7.30\text{--}7.50$  (m, 15H, 15H-ar).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 28.7, 34.0, 53.9, 67.9, 71.1, 71.9, 73.5, 74.1, 77.2, 79.3, 98.0, 126.5\text{--}128.0$  (15C),  $137.0\text{--}137.5$  (3C). MS (FAB+, NBA):  $m/z$ : 563  $[\text{M} + \text{Na}]^+$ , 509  $[\text{M} - \text{OCH}_3]^+$ , 449  $[\text{M} - \text{PhCH}_2]^+$ , 417  $[\text{M} - \text{PhCH}_2 - \text{OCH}_3]^+$ , 91  $[\text{PhCH}_2]^+$ .

### Diammonium Methyl 2,3,4-Tri-*O*-benzyl-6-deoxy-6-dihydroxyphosphonylmethylene- $\alpha$ -D-mannopyranoside **6**

A solution of **5** (130 mg, 0.24 mmol) dissolved in *tris*(trimethylsilyl) phosphite (50 eq, 11.96 mmol) was stirred at  $160^\circ\text{C}$  for 16 h. The reaction mixture was then diluted with  $\text{CH}_2\text{Cl}_2$  (150 mL) and poured into satd.  $\text{NaHCO}_3$  (100 mL) at  $0^\circ\text{C}$ . The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (*i*-PrOH/ $\text{NH}_4\text{OH}/\text{H}_2\text{O}$  6:3:1) affording **6** as white foam after freeze drying from  $\text{H}_2\text{O}$  (118 mg, 85%).  $R_f = 0.50$  (*i*-PrOH/ $\text{NH}_4\text{OH}/\text{H}_2\text{O}$  6:3:1).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 1.50\text{--}2.30$  (m, 4H, H-6 H-6' H-7 H-7'),  $3.17$  (s, 3H,  $\text{OCH}_3$ ),  $3.40\text{--}3.85$  (m, 4H, H-2 H-3 H-4 H-5),  $4.40\text{--}4.90$  (m, 7H, H-1  $3\text{CH}_2\text{Ph}$ ),  $6.70\text{--}7.00$  (m, 8H,  $2\text{NH}_4$ ),  $7.10\text{--}7.40$  (m, 15H, 15H-ar).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 23.3$  (d,  $J = 138.6$  Hz),  $24.3$  (d,  $J = 2.3$  Hz),  $53.7, 70.6\text{--}79.3$  (3C),  $70.7$  (d,  $J = 15.5$  Hz),  $70.8, 71.0, 74.1, 97.7, 126.4\text{--}127.9$  (15C),  $137.1\text{--}137.7$  (3C).  $^{31}\text{P}$  NMR ( $\text{CDCl}_3$ ):  $\delta = 27.2$  (s,  $\text{P}(\text{O})(\text{ONH}_4)_2$ ). MS (FAB+, NBA):  $m/z$ : 565  $[\text{M} - 2\text{NH}_4 + \text{Na}]^+$ , 543  $[\text{M} - 2\text{NH}_4 + \text{H}]^+$ , 511  $[\text{M} - 2\text{NH}_4 - \text{OCH}_3]^+$ , 91  $[\text{PhCH}_2]^+$ .

### Disodium Methyl 6-Deoxy-6-dihydroxyphosphonylmethylene- $\alpha$ -D-mannopyranoside **1**

A suspension of **6** (100 mg, 0.24 mmol) and Pd-C 10% (250 mg) in EtOH/ $\text{H}_2\text{O}$  (1:1, 15 mL) was stirred at rt under  $\text{H}_2$  atmosphere for 16 h. The reaction mixture was filtered through celite and the filtrate evaporated under reduced pressure. The residue was dissolved in  $\text{H}_2\text{O}$  (30 mL) and ion exchange resin (Dowex 50WX2,  $\text{Na}^+$  form, 10 g) was

added and the solution was kept at rt for 24 h before filtering-off the resin and freeze drying from H<sub>2</sub>O to afford **1** as a white foam (48 mg, 90%).  $R_f = 0.16$  (*i*-PrOH/NH<sub>4</sub>OH/H<sub>2</sub>O 6:3:1). <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta = 1.20$ – $2.10$  (m, 4H, H-6 H-6' H-7 H-7'),  $3.26$  (s, 3H, OCH<sub>3</sub>),  $3.30$ – $3.70$  (m, 3H, H-3 H-4 H-5),  $3.79$  (dd, 1H,  $J = 1.4$  and  $3.2$  Hz, H-2),  $4.59$  (d, 1H, H-1). <sup>13</sup>C NMR (D<sub>2</sub>O):  $\delta = 24.2$  (d,  $J = 134.4$  Hz),  $25.4$  (d,  $J = 2.2$  Hz),  $55.0$ ,  $70.2$ – $70.8$  (3C),  $72.7$  (d,  $J = 17.1$  Hz),  $101.1$  (C-1). <sup>31</sup>P NMR (D<sub>2</sub>O):  $\delta = 26.7$  (s, *P*(O)(ONa)<sub>2</sub>). MS (FAB+, NBA):  $m/z$ : 273 [M–2Na + 3H]<sup>+</sup>. MS (FAB–, NBA):  $m/z$ : 271 [M–2Na–H]<sup>–</sup>.

### Methyl 6-Deoxy- $\alpha$ -D-manno-heptopyranoside **7**

A suspension of **3** (1.84 g, 3.84 mmol) and Pd-C 10% (250 mg) in EtOH/H<sub>2</sub>O (1:1, 20 mL) was stirred at rt under H<sub>2</sub> atmosphere for 16 h. The reaction mixture was filtered through celite and the filtrate evaporated under reduced pressure affording **7** as a white foam (760 mg, 95%).  $R_f = 0.50$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH 4:1). <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta = 1.50$ – $1.70$  (m, 1H, H-6),  $1.90$ – $2.10$  (m, 1H, H-6'),  $3.27$  (s, 3H, OCH<sub>3</sub>),  $3.38$  (t, 1H,  $J = 9.5$  Hz, H-4),  $3.45$ – $3.75$  (m, 4H, H-3 H-5 H-7 H-7'),  $3.81$  (dd, 1H,  $J = 1.6$  and  $3.4$  Hz, H-2),  $4.58$  (d, 1H, H-1).

### Methyl 2,3,4-Tri-*O*-acetyl-7-bromo-6,7-dideoxy- $\alpha$ -D-manno-heptopyranoside **8**

To a solution of **7** (760 mg, 3.65 mmol) in C<sub>5</sub>H<sub>5</sub>N (20 mL) was added a solution of TsCl (1.05 g, 5.48 mmol) in C<sub>5</sub>H<sub>5</sub>N (10 mL). After stirring for 5 h at rt, acetic anhydride (5 mL) was added at 0°C. The reaction was stirred at rt for 1 h then poured into satd. NaHCO<sub>3</sub> (150 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  150 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure and C<sub>5</sub>H<sub>5</sub>N was coevaporated with toluene (3  $\times$  100 mL). The residue was dissolved in 2-butanone (50 mL) and LiBr (6.35 g, 73.07 mmol) was added. The reaction was stirred at 85°C for 1 h then cooled to rt and diluted with H<sub>2</sub>O (100 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  100 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (light petroleum/ether 9:1 then 2:3) to afford **8** (800 mg, 55%).  $R_f = 0.57$  (light petroleum/ether 2:3). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.80$ – $2.05$  (m, 2H, H-6 H-6'),  $1.87$  (s, 3H, CH<sub>3</sub>CO),  $1.95$  (s, 3H, CH<sub>3</sub>CO),  $2.02$  (s, 3H, CH<sub>3</sub>CO),  $3.34$  (s, 3H, OCH<sub>3</sub>),  $3.40$ – $3.50$  (m, 2H, H-7 H-7'),  $3.88$  (td, 1H,  $J = 3.3$  and  $9.8$  Hz, H-5),  $4.58$  (d, 1H,  $J = 1.7$  Hz, H-1),  $4.99$  (t, 1H, H-4),  $5.11$  (dd,  $J = 3.4$  Hz, H-2),  $5.18$  (dd, 1H, H-3).

### 1,2,3,4-Tetra- *O*-acetyl-7-bromo-6,7-dideoxy-*D*-manno-heptopyranoside **9**

A solution of **8** (100 mg) in AcOH/Ac<sub>2</sub>O (4:1, 10 mL) and two drops of conc. H<sub>2</sub>SO<sub>4</sub> was stirred at rt for 24 h. The reaction mixture was neutralized with satd. NaHCO<sub>3</sub> (100 mL) and the aqueous layer was extracted with EtOAc (3 × 100 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (light petroleum/ether 95:5 then 3:7) affording **9** as a pale yellow oil (91 mg, 85%) and a portion of **8** (15 mg) was recovered. *R*<sub>f</sub> = 0.61 (light petroleum/ether 3:7). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 1.80–2.00 (m, 2H, H-6 H-6'), 2.00 (s, 3H, CH<sub>3</sub>CO), 2.07 (s, 3H, CH<sub>3</sub>CO), 2.15 (s, 3H, CH<sub>3</sub>CO), 3.40–3.70 (m, 2H, H-7 H-7'), 4.08 (td, 1H, *J* = 3.0 and 9.9 Hz, H-5), 5.16 (t, 1H, H-4), 5.26 (dd, 1H, *J* = 1.8 and 3.5 Hz, H-2), 5.34 (dd, 1H, H-3), 5.99 (d, 1H, H-1). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ = 20.9, 21.1, 21.2, 29.2, 33.9, 68.4–69.6 (4C), 90.6, 170.1, 170.2, 170.3, 170.4. MS (FAB+, NBA): *m/z*: 447 [M + Na]<sup>+</sup>, 365 [M–OAc]<sup>+</sup>, 345 [M–Br]<sup>+</sup>, 245 [M–2OAc–2H]<sup>+</sup>, 203 [M–2OAc–CH<sub>3</sub>CO–H]<sup>+</sup>, 43 [CH<sub>3</sub>CO]<sup>+</sup>.

### 2,3,4-Tri- *O*-acetyl-7-bromo-6,7-dideoxy-1,2- *O*-[8-(cholest-5-en-3β-yloxy)-3,6-dioxaoctan-1-yloxyethylidene]-β-*D*-manno-heptopyranose **11**

A solution of **9** (220 mg, 0.52 mmol) and HBr (5.7 M in AcOH, 2.28 mL, 13 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 mL) was stirred at rt for 16 h. The reaction mixture was poured into satd. NaHCO<sub>3</sub> (50 mL) and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 70 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure affording **10** used for next step as the crude product. To a solution of 8-(cholest-5-en-3β-yloxy)-3,6-dioxaoctan-1-ol<sup>43</sup> (287 mg, 0.57 mmol) and AgOTf (146 mg, 0.57 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added *sym*-collidine (76 μL, 0.57 mmol) and a solution of **10** in CH<sub>2</sub>Cl<sub>2</sub> (15 mL). After stirring at rt for 15 min, the reaction mixture was filtered through celite and the filtrate evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (light petroleum/ether 4:1 then ether) affording **11** as a white gum (206 mg, 45%). *R*<sub>f</sub> = 0.41 (light petroleum/ether 1:4). <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ = 0.67 (s, 3H, H-18<sub>Chol</sub>), 0.86 (d, 6H, *J* = 6.6 Hz, H-26<sub>Chol</sub>), 0.91 (d, 3H, *J* = 6.4 Hz, H-21<sub>Chol</sub>), 0.99 (s, 3H, H-19<sub>Chol</sub>), 0.80–2.50 (m, 30H, 28H-Chol H-6 H-6'), 1.71 (s, 3H, CH<sub>3</sub>†), 2.07 (s, 3H, CH<sub>3</sub>CO), 2.11 (s, 3H, CH<sub>3</sub>CO), 3.10–3.30 (m, 1H, H-3<sub>Chol</sub>), 3.45–3.70 (m, 15H, OCH<sub>2</sub>CH<sub>2</sub>O H-5 H-7 H-7'), 4.59 (dd, 1H, *J* = 2.6 and 3.5 Hz, H-2), 5.00–5.20 (m, 2H, H-3

H-4), 5.30–5.38 (m, 1H, H-6<sub>Chol</sub>), 5.44 (d, 1H, H-1). <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 12.2, 15.6–41.7 (C-Chol CH<sub>3</sub>† C-6), 21.1, 21.2, 29.1, 50.6, 56.5, 57.2, 62.4, 66.2, 67.7, 69.0, 70.3, 70.9, 71.0, 71.1, 71.3, 76.8, 79.9, 97.9, 124.7 (C\*), 141.4, 170.2, 170.7.

### 8-(Cholest-5-en-3 $\beta$ -yloxy)-3,6-dioxaoctan-1-yl-2,3,4-tri-*O*-acetyl-7-bromo-6,7-dideoxy- $\alpha$ -D-*manno*-heptopyranoside **12**

To a suspension of **11** (160 mg, 0.18 mmol) and molecular sieves (3Å) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added dropwise TMSOTf (3.3  $\mu$ L, 18  $\mu$ mol) between –5°C and 0°C (ice-salt bath). The suspension was stirred for 5 min before neutralizing with Et<sub>3</sub>N (50  $\mu$ L) and diluting with H<sub>2</sub>O (50 mL). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  50 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (light petroleum/ether 9:1 then ether) affording **12** as a white gum (64 mg, 40%). *R*<sub>f</sub> = 0.58 (light petroleum/ether 1:4). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.60 (s, 3H, H-18<sub>Chol</sub>), 0.79 (d, 3H, *J* = 6.6 Hz, H-26<sub>Chol</sub>), 0.80 (d, 3H, H-26<sub>Chol</sub>), 0.84 (d, 3H, *J* = 6.5 Hz, H-21<sub>Chol</sub>), 0.92 (s, 3H, H-19<sub>Chol</sub>), 0.70–2.35 (m, 28H, 28H-Chol), 1.85–1.95 (m, 1H, H-6), 1.91 (s, 3H, CH<sub>3</sub>CO), 1.99 (s, 3H, CH<sub>3</sub>CO), 2.07 (s, 3H, CH<sub>3</sub>CO), 2.00–2.10 (m, 1H, H-6'), 3.05–3.15 (m, 1H, H-3<sub>Chol</sub>), 3.40–3.50 (m, 1H, H-7), 3.50–3.70 (m, 13H, OCH<sub>2</sub>CH<sub>2</sub>O H-7'), 3.97 (td, 1H, *J* = 2.5 and 9.9 Hz, H-5), 4.75 (d, 1H, *J* = 1.5 Hz, H-1), 5.04 (t, 1H, H-4), 5.21 (dd, 1H, *J* = 3.5 Hz, H-2), 5.25–5.30 (m, 1H, H-6<sub>Chol</sub>), 5.27 (dd, 1H, H-3). <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 12.3, 19.1–42.7 (C-Chol), 21.1, 21.2, 21.5, 29.2, 34.9, 50.6, 56.6, 57.2, 62.6, 67.7, 67.9, 69.5, 69.8, 70.1, 70.5, 71.0, 71.2, 71.3, 79.9, 97.8, 121.9, 141.4, 170.2, 170.3, 170.5.

### Disodium 8-(Cholest-5-en-3 $\beta$ -yloxy)-3,6-dioxaoctan-1-yl-6-deoxy-6-dihydroxyphosphonylmethylene- $\alpha$ -D-mannopyranoside M6Pn/TEGC **13**

A solution of **12** (67 mg, 76  $\mu$ mol) dissolved in *tris*(trimethylsilyl) phosphite (5 mL) was stirred at 160°C for 16 h. The reaction mixture was then diluted with CH<sub>2</sub>Cl<sub>2</sub> (15 mL) and poured into satd. NaHCO<sub>3</sub> (25 mL) at 0°C. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3  $\times$  50 mL). Organic layers were combined, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated under reduced pressure. The crude mixture (showing a single signal by <sup>31</sup>P NMR spectroscopy at 27 ppm) was dissolved in MeOH (5 mL) and NaOMe (35 mg, 610  $\mu$ mol) was added.

After 24 h, the reaction was neutralized with 1 M HCl and evaporated under reduced pressure. The resulting residue was purified by silica gel column chromatography (*i*-PrOH/NH<sub>4</sub>OH/H<sub>2</sub>O 6:3:1), then treated with ion exchange resin (Dowex 50WX2, Na<sup>+</sup> form, 5 g) affording M6Pn/TEGC **13** (31 mg, 50%).  $R_f = 0.52$  (*i*-PrOH/NH<sub>4</sub>OH/H<sub>2</sub>O 6:3:1). <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta = 0.70$ – $2.30$  (m, 47H, H-Chol H-6 H-6' H-7 H-7'),  $3.20$ – $4.00$  (m, 17H, OCH<sub>2</sub>CH<sub>2</sub>O H-2 H-3 H-4 H-5 H-3<sub>Chol</sub>),  $5.30$ – $5.40$  (m, 1H, H-6<sub>Chol</sub>). <sup>31</sup>P NMR (D<sub>2</sub>O):  $\delta = 23.0$  (s, P(O)(ONa)<sub>2</sub>). -MS (FAB+, G/T):  $m/z$ : 803 [M + H]<sup>+</sup>, 781 [M – Na + 2H]<sup>+</sup>. MS (FAB–, G/T):  $m/z$ : 757 [M – 2Na + H]<sup>–</sup>.

## REFERENCES

- [1] S. Kornfeld, *Annu. Rev. Biochem.*, **61**, 307 (1992).
- [2] M. Stein, J. E. Zijderhand-Bleekemolen, H. Geuze, A. Hasilik, and K. von Figura, *EMBO J.*, **6**, 2677 (1987).
- [3] J. X. Kang, Y. Li, and A. Leaf, *Proc. Natl. Acad. Sci. USA*, **95**, 13671 (1998).
- [4] B. Schmidt, C. Kiecke-Siemsen, A. Waheed, T. Bräulke, and K. von Figura, *J. Biol. Chem.*, **270**, 14975 (1995).
- [5] P. G. Marron-Terrada, M. A. Brzycki-Wessel, and N. M. Dahms, *J. Biol. Chem.*, **273**, 22358 (1998).
- [6] J. X. Kang, J. Bell, A. Leaf, R. L. Beard, and R. A. S. Chandraratna, *Proc. Natl. Acad. Sci. USA*, **95**, 13687 (1998).
- [7] C. Vidil, A. Morère, M. Garcia, V. Barragan, B. Hamdaoui, H. Rochefort, and J.-L. Montero, *Eur. J. Org. Chem.*, 447 (1999).
- [8] C. Vidil, S. Vidal, A. Morère, and J.-L. Montero, *Phosphorus, Sulfur, and Silicon*, **158**, 125 (2000).
- [9] R. Rabinowitz, *J. Org. Chem.*, **28**, 2975 (1963).
- [10] P. Le Marechal, C. Froussios, M. Level, and R. Azerad, *Carbohydr. Res.*, **94**, 1 (1981).
- [11] C. Garbay-Jaureguiberry, D. Ficheux, and B. P. Roques, *Int. J. Peptide Protein Res.*, **39**, 523 (1992).
- [12] M. F. Wang, M. M. L. Crilley, B. T. Golding, T. McInally, D. H. Robinson, and A. Tinker, *J. Chem. Soc., Chem. Commun.*, 667 (1991).
- [13] V. Barragan, F. M. Menger, K. L. Caran, C. Vidil, A. Morère, and J.-L. Montero, *Chem. Commun.*, 85 (2001).
- [14] A. F. Rosenthal and L. A. Vargas, *Tetrahedron Lett.*, 977 (1975).
- [15] K. Jones and W. W. Mood, *Carbohydr. Res.*, **155**, 217 (1986).
- [16] H. H. Baer, L. R. Breton, and Y. Shen, *Carbohydr. Res.*, **200**, 377 (1990).
- [17] H. H. Baer, Y. Shen, and X. Wu, *Carbohydr. Res.*, **241**, 117 (1993).
- [18] A. De Raadt, H. Griengl, and N. Klempier, *J. Org. Chem.*, **58**, 3179 (1993).
- [19] G. O. Aspinall, A. G. McDonald, and R. K. Sood, *Can. J. Chem.*, **72**, 247 (1994).
- [20] D. El Garrouj, S. Aliou, A. Aumelas, and J.-L. Borgna, *J. Med. Chem.*, **38**, 2339 (1995).
- [21] Z. Pakulski and A. Zamojski, *Polish J. Chem.*, **68**, 1109 (1994).
- [22] H. H. Baer and H. R. Hanna, *Carbohydr. Res.*, **102**, 169 (1982).
- [23] Z. Pakulski and A. Zamojski, *Tetrahedron*, **51**, 871 (1995).
- [24] D. Rouzaud and P. Sinaÿ, *J. Chem. Soc., Chem. Commun.*, 1353 (1983).
- [25] P. G. Goekjian, T.-C. Wu, H.-Y. Kang, and Y. Kishi, *J. Org. Chem.*, **56**, 6422 (1991).

- [26] C. Barbaud, M. Bols, and I. Lundt, *Tetrahedron*, **51**, 9063 (1995).
- [27] M. A. Leeuwenburgh, S. Picasso, H. S. Overkleeft, G. A. van der Marel, P. Vogel, and J. H. van Boom, *Eur. J. Org. Chem.*, 1185 (1999).
- [28] Y.-C. Xin, Y.-M. Zhang, J.-M. Mallet, C. P. J. Glaudemans, and P. Sinaÿ, *Eur. J. Org. Chem.*, 471 (1999).
- [29] R. Anderson, *Synthesis*, 717 (1985).
- [30] C. Burford, F. Cooke, G. Roy, and P. Magnus, *Tetrahedron*, **39**, 867 (1983).
- [31] P. A. M. van der Klein and J. H. van Boom, *Carbohydr. Res.*, **224**, 193 (1992).
- [32] Q. Shen, D. G. Sloss, and D. B. Berkowitz, *Synth. Commun.*, **24**, 1519 (1994).
- [33] D. B. Berkowitz, M. Bose, T. J. Pfannenstiel, and T. Doukov, *J. Org. Chem.*, **65**, 4498 (2000).
- [34] T.-L. Ho, H. C. Ho, and C. M. Wong, *J. Chem. Soc., Chem. Commun.*, 791 (1972).
- [35] T.-H. Chuang, J.-M. Fang, W.-T. Jiaang, and Y.-M. Tsai, *J. Org. Chem.*, **61**, 1794 (1996).
- [36] I. E. Marko, A. Ates, A. Gautier, B. Leroy, J.-M. Plancher, Y. Quesnel, and J.-C. Vanherck, *Angew. Chem. Int. Ed.*, **38**, 3207 (1999).
- [37] B. Bernet and A. Vasella, *Helv. Chim. Acta*, **62**, 1990 (1979).
- [38] B. Bernet and A. Vasella, *Helv. Chim. Acta*, **62**, 2400 (1979).
- [39] J. Hooz and S. S. H. Gilani, *Can. J. Chem.*, **46**, 86 (1968).
- [40] D. Seebach and E. J. Corey, *J. Org. Chem.*, **40**, 231 (1975).
- [41] S. Hanessian, *Preparative Carbohydrate Chemistry*, S. Hanessian, ed. (Marcel Dekker, New York, 1997), p 263.
- [42] P. Collins and R. Ferrier, *Monosaccharides: Their Chemistry and Their Roles in Natural Products* (Wiley & Sons: New York, 1995).
- [43] D. Lafont, P. Boullanger, and S. Chierici, *New J. Chem.*, **20**, 1093 (1996).
- [44] W. Wang and F. Kong, *J. Org. Chem.*, **63**, 5744 (1998).
- [45] W. Wang and F. Kong, *Tetrahedron Lett.*, **39**, 1937 (1998).
- [46] D. Crich, Z. Dai, and S. Gastaldi, *J. Org. Chem.*, **64**, 5224 (1999).
- [47] W. Wang and F. Kong, *Tetrahedron Lett.*, **40**, 1361 (1999).
- [48] W. Wang and F. Kong, *J. Org. Chem.*, **64**, 5091 (1999).
- [49] W. Wang and F. Kong, *Angew. Chem., Int. Ed. Engl.*, **38**, 1247 (1999).