

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

β -(N-Acylamino)vinylphosphonium Salts--Synthesis and Properties

Roman Mazurkiewicz^a; Beata Fryczkowska^b; Rafał Gabański^a; Roman Luboradzki^c; Andrzej Włochowicz^b; Wojciech Mól^a

^a Institute of Organic Chemistry and Technology, The Silesian University of Technology, Poland ^b

Department of Textile Engineering and Environmental Sciences, The Technical-Humanistic Academy,

Poland ^c Institute of Physical Chemistry, Polish Academy of Sciences, Poland

Online publication date: 27 October 2010

To cite this Article Mazurkiewicz, Roman , Fryczkowska, Beata , Gabański, Rafał , Luboradzki, Roman , Włochowicz, Andrzej and Mól, Wojciech(2002) ' β -(N-Acylamino)vinylphosphonium Salts--Synthesis and Properties', Phosphorus, Sulfur, and Silicon and the Related Elements, 177: 11, 2589 — 2598

To link to this Article: DOI: 10.1080/10426500214561

URL: <http://dx.doi.org/10.1080/10426500214561>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



β -(N-ACYLAMINO)VINYLPHOSPHONIUM SALTS—SYNTHESIS AND PROPERTIES

Roman Mazurkiewicz,^a Beata Fryczkowska,^b Rafał Gabański,^a
Roman Luboradzki,^c Andrzej Włochowicz,^b and Wojciech Mól^a
Institute of Organic Chemistry and Technology, The Silesian
University of Technology, Poland;^a Department of Textile
Engineering and Environmental Sciences,
The Technical-Humanistic Academy, Poland;^b and Institute
of Physical Chemistry, Polish Academy of Sciences, Poland^c

(Received February 19, 2002; accepted February 21, 2002)

A reaction of β -carbonyl phosphorus ylides with imidoyl halides gives hitherto unknown β -(N-acylamino)vinylphosphonium salts. The same product can be obtained using the N-monosubstituted amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ system instead of imidoyl halide. The key step of the reaction probably involves an intramolecular [1,3] O-to-N migration of the vinyl group, converting the primary O-imidoylation product into β -(N-acylamino)vinylphosphonium salt.

Keywords: β -Carbonyl ylides; β -(N-acylamino)vinylphosphonium salts; imidoylation; imidoyl halides; N-monosubstituted amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ system; rearrangement

INTRODUCTION


Vinylphosphonium salts **1** have been attracting significant attention of synthetic chemists since 1964, when Schweizer¹ realized that the addition of nucleophiles with a carbonyl function to vinylphosphonium salts results in phosphorus ylides **2**, which can undergo the intramolecular Wittig reaction (Scheme 1). Many carbo- and heterocyclic systems have been synthesized in this way.²

Recently, we have preliminarily communicated the synthesis of hitherto unknown β -(N-acylamino)vinylphosphonium salts **7** by imidoylation of β -carbonyl phosphorus ylides **4** with imidoyl halides

Address correspondence to Roman Mazurkiewicz, Institute of Organic Chemistry and Technology, The Silesian University of Technology, Krzywoustego 4, Gliwice, PL-44-100, Poland. E-mail: romanm@zeus.polsl.gliwice.pl

5 (Scheme 2).³ In this article we give a full account concerning this synthesis, including some novel modification of it, which consists in using the N-monosubstituted amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ system instead of imidoyl halide.

The treatment of the β -carbonyl phosphorus ylide with an imido-yl halide in acetonitrile at room temperature for 24 h results in β -(*N*-acylamino)vinylphosphonium salts **7** as stable, usually crystalline compounds in good yields (procedure A and B, Scheme 2, X = Cl or I). Imido-yl iodides were prepared in situ from imido-yl chlorides by exchange of chlorine with NaI. The obtained vinylphosphonium salts can be precipitated from the reaction mixture with Et₂O and purified further, if necessary, by crystallization from a mixture of MeCN or CH₂Cl₂ with Et₂O or by column chromatography.



The preparation of imidoyl halides is rather labor consuming. Apart from that, imidoyl halides, being highly reactive substances, must often be prepared as required, some of them tending to undergo self-condensation.⁴ Taking into account these disadvantages, we tried to use the N-monosubstituted amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ system as a source of an imidoylating agent generated in situ. In 1990 we provided some spectral evidence, that the new imidoylating agent generated from N-monosubstituted amides and Ph_3PBr_2 in the presence of Et_3N had the structure of N,N,N,N'-tetrasubstituted amidinium salts **9**, and could be detected based on a characteristic strong $\nu_{\text{C}=\text{N}}$ absorption band at about



As had been expected, the treatment of β -carbonyl phosphorus ylides with the N-monosubstituted amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ system in CH_2Cl_2 at room temperature for 24 h gives β -(N-acylamino)vinylphosphonium bromides **7**, usually in good yields (procedure C) (Table I). Before adding an ylide, in all the cases we detected in the reaction mixture a strong absorption band of the imidoylating agent at about 1700 cm^{-1} , as well as the characteristic set of absorption bands of triphenylphosphine oxide at 1439 , 1191 , and 1120 cm^{-1} , the intensity of which approximately corresponds to the total conversion of Ph_3PBr_2 into Ph_3PO .

It is obvious, that the final reaction product **7** cannot be formed in a simple, direct way from ylide **4** and imidoyl halide **5**. In order to explain our results we assume this reaction to involve the *O*-imidoylated intermediate **6** and the [1,3] O-to-N sigmatropic migration of its vinyl group. A similar kind of [1,3] sigmatropic migrations is well-known in the literature;⁶ e.g., *O*-imidoylated carboxamides undergo a similar rearrangement.^{5,7} An analogous rearrangement probably also takes place in the case of the similar, well-known acylation of β -carbonyl

TABLE I Synthesis of β -(N-Acyloamino)vinylphosphonium Salts **7**

Ylide 4		Imidoylating agent		β -(<i>N</i> -Acyloamino)vinylphosphonium salt 7				IR [cm ⁻¹]	Elemental analyses (calcd./found) [%]			
R ¹	R ²	R ³	R ⁴	X	No.	Procedure	Yield [%]		m.p. [°C]	C	H	N
H	H	Me	Ph	I	7a	B	91	133–134	61.21/61.12	4.59/4.66	2.55/2.41	5.64/5.77
H	H	Ph	Me	Cl	7b	A	71	238–239	73.44/73.51	5.50/5.74	3.06/3.00	6.76/6.63
H	H	Ph	Me	Br	7c	C	71	242–243	66.94/67.00	5.02/5.09	2.79/2.92	6.17/6.18
H	H	Ph	PhCH ₂	Cl	7d	A	64	273–275	76.47/76.70	5.47/5.30	2.62/2.78	5.80/6.00
H	Me	Ph	Me	Cl	7e	A	66	140–141	73.80/73.54	5.77/5.92	2.97/2.91	6.56/6.42
H	Me	Ph	Me	I	7f	B	85	182–183	61.71/62.09	4.83/4.59	2.49/2.74	5.50/5.42
H	Me	Ph	Me	Br	7g^d	C	80	196–198	67.45/67.19	5.27/5.30	2.71/2.85	6.00/5.97
H	Me	Ph	Ph	Cl	7h	A	72	192–193	76.47/76.21	5.47/5.22	2.62/2.32	5.80/5.61
H	Me	Ph	PhCH ₂	Cl	7i	A	87	175–177	76.70/76.48	5.70/5.91	2.56/2.62	5.65/5.48
H	Me	(CH ₂) ₅		Br	7j	C	79	205.5–206	65.59/65.31	5.91/5.88	2.83/3.00	6.26/6.24
Me	H	Ph	Me	Cl	7k	A	99	Resin	73.80/73.51	5.77/6.01	2.97/3.04	6.56/6.36
Me	H	(CH ₂) ₄		Br	7l	C	62	Resin	65.01/65.33	5.67/5.44	2.92/2.80	6.45/6.15

^aA mixture of two stereoisomers in the ratio of 68:32.

TABLE II ¹H and ¹³C-NMR Spectra of β-(N-Acyloamino)vinylphosphonium Salts **7**

No.	¹ H-NMR (CDCl ₃ /TMS, δ (ppm))	¹³ C NMR [CDCl ₃ /TMS, δ (ppm)/J _{C-P} (Hz)]							
		>C=O	P ⁺ -C=	C-N	Ph ₃ P ⁺				
					C ¹	C ²	C ³	C ⁴	Other carbons
7a	7.75-7.21 (m, 20H, Ph), 7.10 (dd, 1H, J _{P-H} = 13.8 Hz, J _{H-H} = 14.7 Hz, CH), 6.92 (dd, 1H, J _{P-H} = 17.0 Hz, J _{H-H} = 14.9 Hz, CH), 2.05 (s, 3H, Me)	176.6	86.1/99.5	151.1/17.6	119.2/91.1	133.7/10.7	130.2/13.1	134.5/3.0	135.6, 128.7, 125.2, 122.5 (Ph); 25.5 (Me)
	7b 7.8-7.2 (m, 22H, Ph and CH), 3.78 (s, 3H, Me)	170.9	83.5/100.5	150.8/17.8	119.4/91.4	133.8/10.6	130.2/12.9	134.7/3.0	132.8, 131.3, 128.6, 127.8 (Ph); 34.5 (Me)
7c	7.8-7.6 (m, 15H, Ph), 7.48-7.13 (m, 7H, Ph and CH), 3.78 (s, 3H, Me)	170.9	83.3/100.6	150.9/17.8	119.2/91.6	133.8/10.6	130.2/13.0	134.8/3.0	132.6, 131.3, 128.6, 127.9 (Ph); 34.8 (Me)
	7d 7.82-7.26 (m, 25H, Ph), 7.05 (dd, 1H, J _{P-H} = 14.1 Hz, J _{H-H} = 15.0 Hz, CH), 6.84 (dd, 1H, J _{P-H} = 17.4 Hz, J _{H-H} = 15.3 Hz, CH), 5.87 (s, 2H, CH ₂)	171.6	85.1/100.5	149.7/17.8	119.1/91.8	133.8/11.0	130.2/12.9	134.7/3.0	136.2, 133.0, 131.5, 128.9, 128.8, 128.6, 128.1, 127.5 (Ph); 47.9 (CH ₂)
7e	7.90-7.18 (m, 18H, Ph), 6.98 (d, 1H, J _{P-H} = 16.5 Hz, CH), 6.77 (d, 2H, J = 7.2 Hz, o-Ph), 2.98 (s, 3H, Me), 2.75 (s, 3H, Me)	170.4	99.4/97.6	161.6/8.6	120.6/92.5	133.8/10.3	130.0/12.8	134.3/3.1	133.2, 131.0, 128.4, 127.0 (Ph); 37.8 (NMe); 26.1/ 15.4 (CMe)

(Continued on next page)

No.	¹ H-NMR (CDCl ₃ /TMS, δ (ppm))	¹³ C NMR [CDCl ₃ /TMS, δ (ppm)/J _{C-P} (Hz)]					
		C=O	P ⁺ —C=	C—N	Ph ₃ P ⁺		
				C ¹	C ²	C ³	C ⁴
		>					Other carbons
7f^a	7.87–7.67 (m, 15H, Ph ₃ P),	169.2	98.2/94.9	160.9/8.0	120.6/92.5	133.5/10.7	129.7/12.8
	7.45–7.38, (m, 1H, Ph),						134.1/3.0
	7.32–7.25 (m, 2H, Ph),						133.3, 130.7,
	6.93 (d, 1H, J _{P-H} = 16.2 Hz, CH),						128.0, 127.1
7g^b	6.90–6.85 (m, 2H, Ph),						(Ph); 36.7 (NMe);
	2.84 (s, 3H, Me), 2.56						24.8/15.5 (CMe)
	(s, 3H, Me)						
	7.85–7.20 (m, 18H, Ph), 6.89	170.2	99.14/99.5	161.6/8.6	120.4/92.8	133.6/10.3	139.7/13.1
7g^c	(d, 1H, J _{P-H} = 16.8 Hz, CH),						134.0/3.1
	6.77 (d, 2H, J = 7.2 Hz, <i>o</i> -Ph),						128.2, 126.8 (Ph);
	2.99 (s, 3H, Me), 2.75 (s, 3H, Me)						37.8 (NMe); 26.0/
	7.85–7.20 (m, 20H, Ph), 5.85	172.5	90.3/102.2	165.0/11.5	119.6/90.7	133.2/10.6	130.5/13.1
7h	(d, 1H, J _{P-H} = 14.4 Hz, CH),						134.7/3.1
	3.62 (s, 3H, Me), 2.06 (d, 3H, J = 1.8 Hz, Me)						131.7, 128.7,
	7.9–6.5 (m, 26H, Ph and CH),	170.3	98.2/94.4	161.0/5.2	120.1/91.8	134.1/10.3	128.5 (Ph);
	2.45 (s, 3H, Me)						38.4 (NMe);
7h							23.3/5.7 (CMe)
							133.0, 130.8,
							127.8, 127.3
							(PhCO); 137.5,
7h							128.5, 124.1, 122.2
							(PhN); 27.4 (CMe)

7i	7.85–7.22 (m, 25H, Ph), 6.25 (d, 1H, CH, $J_{P-H} = 14.4$ Hz, CH), 5.52 (s, 2H, CH ₂), 1.66 (s, 3H, Me)	172.1	98.8/95.9	164.4/10.6	119.0/90.6	133.2/10.9	130.6/13.3	135.1/3.1	136.8, 135.7, 131.9, 129.00, 128.98, 128.6, 128.5, 127.8 (Ph); 52.3 (CH ₂); 24.3/5.3 (Me)
7j	7.89–7.62 (m, 15H, Ph), 6.86 (dd, 1H, $J_{P-H} = 16.8$ Hz, $J_{H-H} = 0.9$ Hz, CH), 3.34–3.27 (m, 2H, CH ₂), 2.60 (s, 3H, Me), 1.72–1.32 (m, 8H, (CH ₂) ₄)	175.8	99.7/96.1	161.7/4.9	119.9/92.2	134.1/10.4	129.7/13.1	134.2/3.1	49.1, 36.5, 28.8, 28.5, 22.4 (CH ₂); 26.0/15.5 (Me)
7k	7.88–7.18 (m, 20H, Ph), 6.91 (d, 1H, $J_{P-H} = 20.4$ Hz, CH), 3.75 (s, 3H, Me), 2.42 (d, 3H, $J_{P-H} = 15.6$ Hz, Me)	171.8	95.0/91.9	149.5/25.8	117.5/89.3	133.7/10.2	130.5/12.6	135.1/2.6	132.0, 131.1, 128.7, 127.9 (Ph); 36.0 (NMe); 16.6/8.3 (CMe)
7l	7.92–6.85 (m, 16H, Ph and CH), 3.19–3.11 (m, 2H, CH ₂), 1.81–1.40 (m, 6H, (CH ₂) ₃), 2.45 (d, 2H, $J_{P-H} = 14.9$ Hz, Me)	170.8	97.2/94.4	151.1/19.2	118.3/91.1	133.9/10.5	130.1/12.9	134.8/3.0	47.2, 37.9, 23.6, 23.8, 17.1 (Me)

^a In DMSO-*d*₆.^b Minor stereoisomer.^c Major stereoisomer.




FIGURE 1 ORTEP-Plot of the β -(*N*-benzoylamino)- β -methylvinyltriphenylphosphonium chloride **7e**.

ylides; however, being a degenerate rearrangement, it cannot be directly observed.

CONCLUDING REMARKS

The reported reactions offer a convenient way for the synthesis of β -(*N*-acyloamino)vinylphosphonium salts. The phosphonium salts **7** can be considered to be prospective precursors for the synthesis of amino derivatives of carbo- and heterocyclic systems (see Scheme 1), and some important reagents for organic synthesis like *N*-acylaminamines⁸ (by β -elimination of Ph_3P and HX if $\text{R}^2 = \text{H}$) or *N*-vinylamides (by hydro-dephosphonation of phosphonium salts **7**).

EXPERIMENTAL

General

M.p.s, determined in capillary tubes, are uncorrected. IR spectra were recorded on a Zeiss Specord M 80 spectrophotometer; the measurements were carried out in CHCl_3 (0.2 *M*) using cells of 0.105 mm. ^1H and ^{13}C NMR spectra were recorded in CDCl_3 on a Varian UNITY

INOVA-300 spectrometer at operating frequencies of 300 and 75.5 MHz, respectively, in the FT mode using *TMS* as an internal standard.

Starting Materials

Commercial grade acetonitrile and CH_2Cl_2 were distilled and dried over molecular sieves 4A. The following reagents were of commercial quality (Aldrich): (triphenylphosphoranylidene)acetaldehyde, 1-(triphenylphosphoranylidene)-2-propanone and 2-(triphenylphosphoranylidene)propionaldehyde. The synthesis and properties of the following compounds have been reported in the literature: N-phenylacetimidoyl chloride,⁹ N-methylbenzimidoyl chloride,¹⁰ N-benzylbenzimidoyl chloride,¹¹ and N-phenylbenzimidoyl chloride.¹²

Synthesis of β -(N-Acyloamino)vinylphosphonium Chlorides 7 Using Imidoyl Chlorides (General Procedure A)

To a solution of imidoyl halide **5** (2.4 mmol) in MeCN (3.6 ml) ylide **4** (2 mmol) was added and the mixture was left at room temperature for 24 h. The phosphonium salt was precipitated from the reaction mixture with Et_2O (5–8 ml). The product can be purified further, if necessary, by column chromatography on silica gel eluting with a mixture of CH_2Cl_2 and MeOH (97:3 or 99:1, v/v). Crude **7** can usually be recrystallized by dissolving in acetonitrile or CH_2Cl_2 and precipitating with diethyl ether (1:1–1:2; v/v).

Synthesis of β -(N-Acyloamino)vinylphosphonium Iodides 7 Using Imidoyl Chlorides (General Procedure B)

To a solution of imidoyl halide **5** (2.4 mmol) in MeCN (3.6 ml) NaI (0.30 g, 2 mmol) and ylide **4** (2 mmol) was added and the mixture was left at room temperature for 24 h. The precipitated NaCl was filtered off and the phosphonium salt was isolated from the reaction mixture as described above (procedure A).

Synthesis of β -(N-Acyloamino)vinylphosphonium Bromides 7 Using the N-monosubstituted Amide/ $\text{Ph}_3\text{PBr}_2/\text{Et}_3\text{N}$ System (General Procedure C)

To a solution of triphenylphosphine (0.63 g, 2.4 mmol) in CH_2Cl_2 (8 ml) a solution of bromine (0.38 g, 2.4 mmol) in CH_2Cl_2 (1 ml) was added at room temperature under argon atmosphere. After 30 min Et_3N (0.85 ml, 6 mmol) and amide (2.2 mmol) was added. After next 30 min ylide (2 mmol) was added and the mixture was left at room temperature

for 24 h. The solvent was evaporated and the phosphonium salt was isolated from the reaction mixture by column chromatography as described above (procedure A).

REFERENCES

- [1] E. E. Schweizer, *J. Am. Chem. Soc.*, **86**, 2744 (1964); E. E. Schweizer and K. K. Light, *J. Am. Chem. Soc.*, **86**, 2963 (1964); E. E. Schweizer and C. J. Berninger, *Chem. Soc., Chem. Commun.*, 92 (1965); E. E. Schweizer and G. J. O'Neill, *J. Org. Chem.*, **30**, 2082 (1965).
- [2] E. Zbiral, In *Organophosphorus Reagents in Organic Synthesis*, edited by J. I. G. Cadogan (London, 1979), p. 250; A. W. Johnson, W. C. Kaska, K. A. Ostojka Starzewski, and D. A. Dixon, *Ylides and Imines of Phosphorus* (Wiley, New York, 1993), p. 112; I. Burley and A. T. Newson, *Tetrahedron Lett.*, **35**, 7099 (1994).
- [3] R. Mazurkiewicz, B. Fryczkowska, R. Luboradzki, A. Włochowicz, and W. Mól, *Tetrahedron Lett.*, **42**, 8725 (2001); the erratum to this paper: *Tetrahedron Lett.*, **43**, 905 (2002).
- [4] R. Bunnet, In *The Chemistry of the Carbon-Nitrogen Double Bond*, edited by S. Patai (Interscience Publishers, London (1970), ch. 13.
- [5] R. Mazurkiewicz, *Acta Chim. Hung.*, **127**, 439 (1990).
- [6] D. Y. Curtin and L. L. Miller, *J. Am. Chem. Soc.*, **89**, 637 (1967).
- [7] M. G. Hitzler, M. Lutz, P. B. Shrestha-Dawadi, and J. C. Jochims, *Liebigs Ann. Chem.*, 247 (1996).
- [8] J. Collard-Motte and Z. Janousek, In *Topics in Current Chemistry* (Springer-Verlag, Berlin, 1986), p. 89.
- [9] O. Wallach and M. Hoffmann, *Liebigs Ann. Chem.*, **184**, 87 (1877).
- [10] R. Mazurkiewicz and T. Kiersznicki, *Pol. J. Chem.*, **55**, 547 (1981).
- [11] H. von Pechmann and B. Heinze, *Chem. Ber.*, **30**, 1783 (1897).
- [12] J. von Braun and W. Pinkernelle, *Chem. Ber.*, **67**, 1218 (1934).