

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### Part 1: Synthesis of Polyfused Heterocyclic Systems Derived From 3-Phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-b][1,3,4]thiadiazepine-6,8-dione

O. A. Abd Allah<sup>a</sup>

<sup>a</sup> Chemistry Department, Faculty of Science, South Valley University, Sohag, Egypt

Online publication date: 27 October 2010

**To cite this Article** Allah, O. A. Abd(2002) 'Part 1: Synthesis of Polyfused Heterocyclic Systems Derived From 3-Phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-b][1,3,4]thiadiazepine-6,8-dione', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 177: 12, 2871 – 2881

**To link to this Article:** DOI: 10.1080/10426500214876

**URL:** <http://dx.doi.org/10.1080/10426500214876>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



## PART 1: SYNTHESIS OF POLYFUSED HETEROCYCLIC SYSTEMS DERIVED FROM 3-PHENYL-5,6,7,8-TETRAHYDRO[1,2,4]TRIAZOLO- [3,4-*b*][1,3,4]THIADIAZEPINE-6,8-DIONE

O. A. Abd Allah  
Chemistry Department, Faculty of Science, South Valley  
University, Sohag, Egypt

(Received January 27, 2002; accepted April 2, 2002)

3-Phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6,8-dione (**1**) was condensed with *o*-aminothiophenol, 2-amino-ethanol or cystamine to afford compounds **2–4** respectively. Treatment of compound **1** with dimethylthiomethylenemalononitrile yielded the corresponding pyrano[3,2-*f*][1,2,4]triazolo[3,4-*b*]-[1,3,4]thiadiazepine derivative **5**. 7-[5-Amino-1,3-dithiolan-2-ylidene]-3-phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6,8-dione (**6**) was obtained by treating compound **1** with CS<sub>2</sub> and chloroacetonitrile. Thiation of compound **1** gave the corresponding thioanalog **7**, which in turn was condensed with malononitrile to give 3-phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6-one-8-ylidenemalononitrile (**8**). On treating compound **8** with benzaldehyde or *p*-nitrobenzaldehyde, pyrano[1,2,4]triazolo[1,3,4]thiadiazepine derivatives **9a,b**, respectively, were obtained. Compound **8** was treated with CS<sub>2</sub> and methyl iodide to give the corresponding dithiomethyl-methylene derivative **10** which was subjected to react with aniline to give pyrido[1,2,4]triazolo[1,3,4]thiadiazepine derivative **11**. Compound **8** was treated with 3-aminopyridine, *o*-aminothiophenol, or *o*-phenylenediamine to yield compounds **12** and **13a,b** respectively. Finally, tertiary amines or activated phenols were condensed with compound **8** to yield compounds **14** and **15a,b** respectively.

**Keywords:** CS<sub>2</sub>; malonyl dichloride; PTC; thiadiazepine; triazole

## INTRODUCTION

It has been reported that free and fused s-triazoles constitute an important class of compounds with a wide biological activities.<sup>1–4</sup> 1,2,4-Triazolo[3,4-*b*][1,3,4]thiadiazole derivatives have anti-bacterial<sup>5,6</sup>

Address correspondence to O. A. Abd Allah, Department of Chemistry, Faculty of Science, Sohag, Egypt. E-mail: omymatif@yahoo.com

and anti-inflammatory<sup>7</sup> activities, an interesting CNS depressant action,<sup>8</sup> and moderate anti-malarial and anti-tumor activities.<sup>9</sup> The s-triazolo(3,4-*b*) 1,3,4-thiadiazepines showed significant antimicrobial properties.<sup>10,11</sup> From this view and in the extension of our previous work<sup>12–14</sup> in the synthesis of free and fused s-triazole derivatives, it is reported here in the synthesis of a new series of polyfused 1,2,4-triazolo[3,4-*b*][1,3,4]thiadiazepines.

## RESULTS AND DISCUSSION


3-Phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6,8-dione (**1**) was prepared in a good yield via the reaction of 4-amino-5-phenyl-1,2,4-triazole-3-thione with malonyl dichloride and TEA in 1:1:2 molar ratio at room temperature. The structural assignment of this compound was established by elemental and spectral analyses (Scheme 1, Table I).

Compound **1** was condensed with *o*-aminothiophenol, 2-aminoethanol, or cystamine in refluxing diphenyl ether through intermolecular cyclization with the elimination of two water molecules to give the polyfused derivatives **2–4**, respectively.

Also, compound **1** was reacted in boiling DMF with dimethylthiomethylenemalononitrile, which was prepared via the reaction of malononitrile and CS<sub>2</sub> with two equivalents of methyl iodide in a one-pot reaction using liquid-liquid phase-transfer catalysis (PTC) technique [NaOH/dioxane/tetrabutylammonium bromide (TBAB)], to give 8-cyano-9-imino-7-methylthio-6-oxo-3-phenyl-5,6,8,9-tetrahydro-7*H*-pyrano[3,2-*f*][1,2,4]-triazolo[3,4-*b*][1,3,4]thiadiazepine (**5**) (Scheme 1, Table I).

The 7-[5-amino-1,3-dithiolan-2-ylidene]-3-phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6,8-dione (**6**) was obtained through the intermediate **M** by treating compound **1** with CS<sub>2</sub> and chloroacetonitrile in one-pot reaction under PTC experimental conditions [DMF/K<sub>2</sub>CO<sub>3</sub>/TBAB] (Scheme 1, Table I).

Thiation of compound **1** was performed with P<sub>2</sub>S<sub>5</sub> in boiling pyridine to give the corresponding thioanalog 3-phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6-oxo-8-thione (**7**). The IR spectra for this compound showed the absence of a signal corresponding to C=O and the presence of a C=S signal at 1115 cm<sup>-1</sup> (Scheme 2, Table I). The condensation of compound **7** with malononitrile in boiling DMF gave the dicyanomethylene derivative **8**. The IR and <sup>1</sup>H-nmr spectral data are consistent with the proposed structures (Scheme 2, Table I).



SCHEME 1

TABLE I Analytical and Spectral Data of the Prepared Compounds

Comp. no.	m.p. <sup>a</sup> (crys. solvent)	Yield (%)	M <sub>F</sub> /(M <sub>w</sub> )	Analytical data calc. (found) <sup>b</sup> %				IR (KBr) ν (cm <sup>-1</sup> ) <sup>c</sup>	<sup>1</sup> H-NMR (DMSO- <i>d</i> <sub>6</sub> ) <sup>d</sup> δ (ppm)
				C	H	N	S		
1	240 dioxane	79	C <sub>11</sub> H <sub>8</sub> N <sub>4</sub> SO <sub>2</sub> (260.29)	50.76 50.80	3.10 3.12	21.53 21.50	12.32 12.33	3432 (NH); 3060, 2920 (CH); 1708, 1635 (2C=O)	7.20 (s, 1H, NH); 7.15–6.65 (m, 5H, arom.); 2.97 (s, 2H, CH <sub>2</sub> )
2	162 DMF/H <sub>2</sub> O	61	C <sub>17</sub> H <sub>11</sub> N <sub>5</sub> S <sub>2</sub> (349.45)	58.43 58.52	3.18 3.15	20.04 20.14	18.35 18.40	3340 (NH); 3050 (CH); 1620 (C=N)	8.20 (s, 1H, NH); 8.00–6.30 (m, 9H, arom.); 6.20 (s, 1H, CH)
3	198–200 DMF/H <sub>2</sub> O	55	C <sub>13</sub> H <sub>11</sub> N <sub>5</sub> SO (285.33)	54.72 54.85	54.72 54.76	24.55 24.38	11.24 11.31	3434 (NH); 3150, 2937 (CH); 1636 (C=N)	7.32 (s, 1H, NH); 7.10–6.45 (m, 5H, arom.); 6.40 (s, 1H, CH); 4.11–3.91 (t, 2H, CH <sub>2</sub> ); 2.79–2.20 (t, 2H, CH <sub>2</sub> )
4	>300 DMF/H <sub>2</sub> O	51	C <sub>13</sub> H <sub>11</sub> N <sub>5</sub> S <sub>2</sub> (301.39)	51.81 52.00	3.68 3.75	23.24 23.30	21.28 21.39	3440 (NH); 3140, 2920 (CH); 1620 (C=N)	7.82–6.90 (m, 5H, arom.); 6.11 (s, 1H, CH); 5 (s, 1H, NH); 3.89–3.71 (t, 2H, CH <sub>2</sub> ); 2.85–2.51 (t, 2H, CH <sub>2</sub> )
5	Dec 300 n-butanol	59	C <sub>16</sub> H <sub>10</sub> N <sub>6</sub> S <sub>2</sub> O <sub>2</sub> (366.42)	52.45 52.65	2.75 2.88	22.94 22.90	17.50 17.39	3438, 3320 (2NH); 3150, 2954 (CH); 2208 (CN); 1648 (C=O); 1560 (C=N)	7.85–7.00 (m, 5H, arom.); 4.08–3.82 (br, 2H, 2NH); 2.91 (s, 3H, CH <sub>3</sub> )
6	160 DMF/H <sub>2</sub> O	49	C <sub>14</sub> H <sub>9</sub> N <sub>5</sub> S <sub>3</sub> O <sub>2</sub> (287.47)	16.71 16.53	3.16 3.27	24.36 24.48	33.46 33.41	3422, 3340 (NH <sub>2</sub> ); 3250 (NH); 3050, 2910 (CH); 1700 (C=O); 1632 (C=N)	8.00 (s, 1H, NH); 7.72–7.00 (m, 5H, arom.); 5.00–4.81 (br, 2H, NH <sub>2</sub> ); 3.73 (s, 1H, CH)
7	280 Pyridine/H <sub>2</sub> O	73	C <sub>11</sub> H <sub>8</sub> N <sub>4</sub> S <sub>2</sub> O (276.34)	47.81 47.92	2.92 3.00	20.27 20.39	23.21 23.32	3422 (NH); 3060, 2910 (CH); 1647 (C=O); 1060 (C=S)	7.51 (s, 1H, NH); 7.40–6.32 (m, 5H, arom.); 2.24 (s, 2H, CH <sub>2</sub> )


<b>8</b>	>300 DMF/H <sub>2</sub> O	68	C <sub>14</sub> H <sub>8</sub> N <sub>6</sub> SO (308.32)	54.54 54.70	2.62 2.60	27.26 27.19	10.40 10.49	3312 (NH); 3185, 2926 (CH); 2199, 2128 (2CN); 1641 (C=O)	7.30 (s, 1H, NH); 7.15–6.40 (m, 5H, arom.); 2.66 (s, 2H, CH <sub>2</sub> )
<b>9a</b>	>300 DMF/H <sub>2</sub> O	63	C <sub>21</sub> H <sub>14</sub> N <sub>6</sub> SO <sub>2</sub> (414.45)	60.86 60.99	3.41 3.50	20.28 20.18	7.74 7.80	3407, 3210 (2NH); 3060, 2928 (CH); 2203 (CN); 1620 (C=O).	7.40 (s, 1H, NH); 7.36–6.32 (m, 10H, arom.); 3.39 (s, 1H, NH); 2.66 (s, 1H, CH); 2.25 (s, 1H, Cl)
<b>9b</b>	>300 DMF/H <sub>2</sub> O	72	C <sub>21</sub> H <sub>13</sub> N <sub>7</sub> SO <sub>4</sub> (459.45)	54.90 54.79	2.85 3.00	21.34 21.25	6.98 6.89	3421, 3225 (2NH); 3030, 2926 (CH); 2204 (CN); 1621 (C=O); 1530, 1350 (NO <sub>2</sub> )	8.49 (s, 1H, NH); 8.12–7.00 (m, 9H, arom.); 4.30 (s, 1H, NH); 2.60 (s, 1H, CH); 2.42 (s, 1H, CH)
<b>10</b>	>300 DMF/H <sub>2</sub> O	50	C <sub>17</sub> H <sub>12</sub> N <sub>6</sub> S <sub>3</sub> O (412.52)	49.50 49.61	2.93 3.10	20.37 20.31	23.32 23.40	3429 (NH); 3085, 2930 (CH); 2200 (CN); 1653 (C=O)	7.41–6.80 (m, 5H, arom.); 3.00 (s, 1H, NH); 2.83 (s, 3H, CH <sub>3</sub> ); 2.68 (s, 3H, CH <sub>3</sub> )
<b>11</b>	>300 DMF/H <sub>2</sub> O	39	C <sub>22</sub> H <sub>15</sub> N <sub>7</sub> S <sub>2</sub> O (457.54)	57.75 57.74	3.30 3.30	21.43 21.50	14.02 14.00	3417, 3220 (2NH); 3050, 2927 (CH); 2205 (CN); 1620 (C=O)	7.82 (s, 1H, NH); 7.70–6.31 (m, 10H, arom.); 2.85 (s, 1H, NH); 2.41 (s, 3H, CH <sub>3</sub> )
<b>12</b>	Dec 300 DMF/H <sub>2</sub> O	83	C <sub>18</sub> H <sub>13</sub> N <sub>7</sub> SO (375.42)	57.74 57.90	3.49 3.35	26.19 26.02	8.56 8.47	3426, 3230 (2NH); 3045, 2928 (CH); 2201 (CN); 1631 (C=O)	8.69–8.00 (m, 4H, heterocyclic); 7.51 (s, 2H, 2NH); 7.31–6.32 (m, 5H, arom.); 2.55 (s, 2H, CH <sub>2</sub> )
<b>13a</b>	>300 DMF/H <sub>2</sub> O	75	C <sub>18</sub> H <sub>13</sub> N <sub>5</sub> OS <sub>2</sub> (379.47)	56.97 57.13	3.45 3.54	18.46 18.40	16.90 17.00	3441, 3250 (2NH); 3030, 2910 (CH); 1632 (C=O)	7.42 (s, 1H, NH); 7.35–5.20 (m, 9H, arom.); 2.32 (s, 2H, CH <sub>2</sub> ); 2.15 (s, 1H, NH)

(Continued on next page)

TABLE I Analytical and Spectral Data of the Prepared Compounds (Continued)

Comp. no.	m.p. °C <sup>a</sup> (crys. solvent)	Yield (%)	M <sub>F</sub> /(M <sub>w</sub> )	Analytical data calc. (found) <sup>b</sup> %				IR (KBr) ν (cm <sup>-1</sup> ) <sup>c</sup>	<sup>1</sup> H-NMR (DMSO- <i>d</i> <sub>6</sub> ) <sup>d</sup> δ (ppm)
				C	H	N	S		
13b	Dec 300 DMF/H <sub>2</sub> O	71	C <sub>18</sub> H <sub>14</sub> N <sub>6</sub> SO (362.42)	59.65	3.89	23.19	8.84	3441, 3250 (3NH); 3010, 2930 (CH); 1640 (C=O)	7.94 (s, 1H, NH); 7.78 (s, 2H, 2NH); 7.41–6.20 (m, 9H, arom.); 2.38 (s, 2H, CH <sub>2</sub> )
				59.78	4.00	23.12	8.76		7.68 (s, 1H, NH); 7.40–6.15 (m, 9H, arom.); 2.40 (s, 2H, CH <sub>2</sub> ); 2.20 (s, 6H, 2CH <sub>3</sub> )
14	>300 DMF/H <sub>2</sub> O	60	C <sub>21</sub> H <sub>18</sub> N <sub>6</sub> SO (402.48)	62.67	4.51	20.88	7.97	3425 (NH); 3040, 2905 (CH); 2200 (CN); 1632 (C=O)	8.30 (NH); 8.00–6.67 (m, 8H, arom.); 3.65 (s, 1H, OH); 2.82 (s, 2H, CH <sub>2</sub> ); 2.15 (s, 1H, OH)
				62.48	4.61	20.79	7.83		8.00 (s, 1H, NH); 7.32–6.31 (m, 8H, arom.); 2.56 (s, 2H, CH <sub>2</sub> ); 1.91 (s, 3H, CH <sub>3</sub> ); 1.10 (s, 1H, OH)
15a	>300 DMF/H <sub>2</sub> O	69	C <sub>19</sub> H <sub>13</sub> N <sub>5</sub> SO <sub>3</sub> (632.73)	36.07	2.07	11.07	5.07	3550 (2OH); 3434 (NH); 3060, 2905 (CH); 2204 (CN); 1639 (C=O)	
				36.26	2.20	11.15	5.20		
15b	>300 DMF/H <sub>2</sub> O	62	C <sub>20</sub> H <sub>15</sub> N <sub>5</sub> SO <sub>2</sub> (389.44)	61.68	3.88	17.98	8.23	3430 (NH); 3030, 2910 (CH); 2200 (CN); 1640 (C=O)	
				61.80	3.95	17.90	8.39		

<sup>a</sup>Uncorrected.  
<sup>b</sup>Satisfactory microanalyses; obtained; C, ±0.35%; H, ±0.11%; N, ±0.40%; S, ±0.21%.  
<sup>c</sup>Measured on Nicolet 710 FT-IR spectrophotometer.  
<sup>d</sup>Measured with a Varian EM 360L spectrometer at 60 MHz using TMS as internal standard.



SCHEME 2

Compound **8** was allowed to react with benzaldehyde or *p*-nitrobenzaldehyde to afford the corresponding 10-cyano-9-imino-6-oxo-3-phenyl-7-phenyl(*p*-nitrophenyl)-5,6,9,10-tetrahydro-7*H*-pyrano[3,4-*f*]-[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepines (**9a,b**) respectively. The



reaction mechanism was postulated to proceed through the nucleophilic attack of the active  $\text{CH}_2$  group of compound **8** at the carbonyl group of the aromatic aldehyde producing the corresponding aldol adduct, followed by intramolecular cyclization via the nucleophilic addition of the formed OH group at the CN group. The IR spectra of compounds **9a,b** showed the absence of a band corresponding to one CN group and the presence of the adsorption bands corresponding to  $=\text{NH}$ , CN and  $\text{C}=\text{O}_{\text{amidic}}$  groups at 3421, 2204 and  $1650\text{ cm}^{-1}$ , respectively (Scheme 2, Table I).

Compound **8** was treated with an equimolar ratio of  $\text{CS}_2$  and double molarity of methyl iodide to yield the corresponding *s*-methyl derivative **10**; the latter, in turn, underwent cyclization when treated with aniline in 1:1 or 1:2 molar ratio, affording 3,8-diphenyl-5,6,7,8,9,10-hexahydropyrido-9-imino-7-methylthio-6-oxo[3,4-*f*][1,2,4]triazolo[3,4-*b*][1,3,4]-thiadiazepine (**11**).

The behaviour of compound **8** toward amines and phenols was investigated. The products resulted from initial attack of the nucleophile at the C- $\alpha$  of the dicyanomethylene moiety.<sup>15</sup> Thus, compound **8** was condensed with 3-aminopyridine in a 1:3 molar ratio through the hydrogen cyanide elimination to yield the condensed product **12**. Also treatment of compound **8** with *o*-aminothiophenol or *o*-phenylenediamine in refluxing DMF yielded the cyclized products **13a,b** via elimination of two HCN molecules. Moreover, compound **8** was allowed to react with *N,N*-dimethylaniline or activated phenols, namely resorcinol or *o*-cresol, to furnish the desired compounds **14** or **15a,b** respectively.

The structures of all products were established by using elemental and spectral analyses.

## EXPERIMENTAL

### Synthesis of 3-Phenyl-5,6,7,8-tetrahydro[1,2,4]triazolo[3,4-*b*][1,3,4]thiadiazepine-6,8-dione (**1**)

To a solution of 4-amino-5-phenyl-1,2,4-triazole-3-thione (0.01 mol) in dry *p*-xylene (30 ml) were added triethylamine (0.02 mol) and malonyl dichloride (0.01 mol) dropwise. The reaction mixture was stirred for 8 h at room temperature. The reaction mixture was then filtered off, dried, and washed with water. The precipitate was dried and crystallized from dioxane (Table I).

## Synthesis of Compounds 2–4

### General Procedure

To a solution of compound **1** (0.01 mol) in diphenyl ether (15 ml), was added *o*-aminothiophenol, 2-aminoethanol, or cystamine (0.01 mol). The reaction mixture was refluxed for 3 h, concentrated, and cooled. The precipitate was filtered off and crystallized from suitable solvent (Table I).

## Synthesis of Compound 5

A mixture of compound **1** (0.005 mol) and dimethylthiomethylene malononitrile (0.005 mol) was refluxed in *n*-butanol (15 ml) until elimination of MeSH ceased ( $\approx 10$  h). The reaction mixture was concentrated and cooled. The separated solid was filtered off and crystallized from *n*-butanol (Table I).

## Synthesis of Compound 6

A mixture of 4 g anhydrous potassium carbonate, compound **1** (0.004 mol) in DMF (20 ml), and catalytic amount of TBAB was treated with 0.004 mol of CS<sub>2</sub> and stirred for 4 h at 60°C. After that, chloroacetonitrile (0.004 mol) was added, and the mixture was stirred for additional 16 h at 70°C. The reaction mixture was filtered, and the filtrate was added to ice-cold water. The separated solid was filtered off and crystallized from DMF/H<sub>2</sub>O to give compound **6** (Table I).

## Synthesis of Compound 7

To a solution of compound **1** (0.01 mol) in dry pyridine (30 ml), was added phosphorus pentasulphide (0.01 mol). The reaction mixture was refluxed for 7 h, filtered while hot, and concentrated. The precipitate was filtered off and crystallized from pyridine/H<sub>2</sub>O (Table I).

## Synthesis of Compound 8

A mixture of compound **7** (0.01 mol), and malononitrile (0.01 mol) in DMF (20 ml) was refluxed. The evolution of H<sub>2</sub>S was ceased ( $\approx 20$  h). The reaction mixture was cooled to room temperature and poured into ice-cold water (100 ml). The separated solid was filtered off and crystallized from DMF/H<sub>2</sub>O (Table I).

## Synthesis of Compounds 9a,b

### General Procedure

A mixture of compound **8** (0.004 mol), benzaldehyde or *p*-nitro-benzaldehyde (0.004 mol), and a catalytic amount of anhydrous sodium acetate in DMF (20 ml) was refluxed for 7 h. The reaction mixture was poured into ice-cold water (100 ml). The solid product was filtered off and crystallized from the suitable solvent (Table I).

## Synthesis of Compound 10

A mixture of 4 g anhydrous potassium carbonate, compound **8** (0.006 mol), and catalytic amount of TBAB in DMF (20 ml) was treated with 0.006 mol of CS<sub>2</sub> and stirred for 4 h at room temperature. Methyl iodide (0.012 mol) was added, and the mixture was stirred for an additional 4 h at room temperature. The reaction mixture was filtered off, and the filtered was added into ice-cold water. The product was filtered off and crystallized from DMF/H<sub>2</sub>O (Table I).

## Synthesis of Compound 11

A mixture of compound **10** (0.002 mol) and aniline (0.002 mol) in DMF (15 ml) was refluxed for 3 h. The reaction mixture was poured into ice-cold water (100 ml) containing a few drops of dil. HCl. The separated solid was filtered off, washed with water, dried, and crystallized from DMF/H<sub>2</sub>O (Table I).

## Synthesis of Compounds 12–15a,b

### General Procedure

To a solution of compound **8** (0.002 mol) in DMF (20 ml) was added in portions the amine or phenol (0.006 mol) in DMF (20 ml). The mixture was refluxed for 2 h. The reaction mixture was added into ice-cold water (100 ml). The precipitate was filtered off, dried, and crystallized from the proper solvent (Table I).

## REFERENCES

- [1] M. C. Hosur, M. B. Talawar, U. V. Laddi, R. S. Bennur, and S. C. Bennur, *Indian J. Chem.*, **34B**, 707 (1995).
- [2] R. P. Srivastava, V. V. Kumar, S. Bhatia, and S. Sharma, *Indian J. Chem.*, **34B**, 209 (1995).

- [3] S. Sudan, R. Gupta, R. Swara, G. B. Singh, S. Bani, and P. L. Kachroo, *Indian J. Chem.*, **34B**, 458 (1995).
- [4] M. H. Khan and S. Giri, *Indian J. Chem.*, **34B**, 1007 (1995).
- [5] J. Mohan and G. S. R. Anjaneyulug, *Curr. Sci.*, **58**, 1028 (1989).
- [6] A. M. M. E. Omar and O. M. Aboulwafa, *J. Heterocyclic. Chem.*, **23**, 1339 (1986).
- [7] A. R. Rasad, T. Ramalingamat, A. B. Rao, P. W. Diwan, and P. B. Sattur, *Indian J. Chem.*, **25B**, 556 (1986).
- [8] A. A. Deshmukh, M. K. Mody, T. Ramalingamt, and P. B. Sattur, *Indian J. Chem.*, **25B**, 793 (1985).
- [9] F. P. Invidiata, S. Grimaudo, P. Giammanco, and L. Giammanco, *Farmaco.*, **46**, 1489 (1991).
- [10] M. Kidwai, P. Sapra, P. Misra, R. K. Saxena, and M. Singh, *Bioorganic and Medicinal Chemistry*, **9**(2), 217 (2001).
- [11] K. M. Hussain and G. Somari, *Indian J. Chem. Sect. B*, **34**(11), 1007 (1995).
- [12] M. T. M. El-Wassimy, M. Abdel-Rahman, A. B. A. G. Ghattas, and O. A. A. Abd Allah, *Phosphorus, Sulfur, and Silicon*, **70**, 99 (1992).
- [13] A. B. A. G. Ghattas, H. M. Moustafa, O. A. Abd Allah, and A. A. Amer, *Synthetic Comm.*, **31**(16), 2447 (2001).
- [14] A. B. A. G. Ghattas, O. A. Abd Allah, H. M. Moustafa, and A. A. Amer, *Egypt. J. Chem.*, 2002 (under pub.).
- [15] S. A. M. Metwally, G. M. El-Naggar, M. I. Younis, T. I. El-Emary, and M. H. Elnagdi, *Liebigs Ann. Chem.*, 1037 (1989).