


This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Microwave Assisted Synthesis of 6-Aryl-3-substituted-5H-[1,2,4]-triazolo[4,3-b][1,2,4]triazoles: A Case for a Comparative Study

M. Rahimizadeh^a; A. Davoodnia^a; M. M. Heravi^b; M. Bakavoli^a

^a Ferdowsi University of Mashhad, Iran ^b Azzahra University, Tehran, Iran

Online publication date: 27 October 2010

To cite this Article Rahimizadeh, M. , Davoodnia, A. , Heravi, M. M. and Bakavoli, M.(2002) 'Microwave Assisted Synthesis of 6-Aryl-3-substituted-5H-[1,2,4]-triazolo[4,3-b][1,2,4]triazoles: A Case for a Comparative Study', Phosphorus, Sulfur, and Silicon and the Related Elements, 177: 12, 2923 — 2929

To link to this Article: DOI: 10.1080/10426500214890

URL: <http://dx.doi.org/10.1080/10426500214890>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.



MICROWAVE ASSISTED SYNTHESIS OF 6-ARYL-3-SUBSTITUTED-5H-[1,2,4]- TRIAZOLO[4,3-b][1,2,4]TRIAZOLES: A CASE FOR A COMPARATIVE STUDY

*M. Rahimizadeh,^a A. Davoodnia,^a M. M. Heravi,^b
and M. Bakavoli^a*
*Ferdowsi University of Mashhad, Iran^a and Azzahra University,
Vanak, Tehran, Iran^b*

(Received May 10, 2002; accepted July 14, 2002)

*A simple and fast synthesis of 6-aryl-3-substituted-5H-[1,2,4]-triazolo[4,3-b][1,2,4]triazoles **4** in high yields has been developed by microwave assisted heterocyclization of N-(3-methylthio-5-substituted-4H-1,2,4-triazol-4-yl)benzenecarboximidamides **3**. The effectiveness of the microwave irradiation and conventional heating for the formation of compounds **4** has been investigated.*

Keywords: 1,2,4-Triazolo[4,3-b][1,2,4]triazoles; conventional heating; heterocyclization; methyl iodide; microwave irradiation; photographic coupler

1,2,4-Triazolo[4,3-b][1,2,4]triazoles are very interesting compounds with diverse properties. These compounds have been considered as antifungal¹ and photographic couplers.^{2–4} The routes to 1,2,4-triazolo[4,3-b][1,2,4]triazoles mainly involve heterocyclization of suitably substituted-1,2,4-triazoles with appropriate reagents such as acid chlorides, hydroximoyl chloride, halonitriles, isocyanates, carbon disulfide, and diarylcarbodiimides in suitable solvents (see for example references^{5–10}). A literature survey disclosed that aryl nitriles have been relatively little explored for the synthesis of 1,2,4-triazolo[4,3-b][1,2,4]triazoles and there are only two references cited in the literature dealing with this matter.^{11,12} Thus, it seemed of interest to examine the reaction of aryl nitriles with substituted 1,2,4-triazoles as a


Address correspondence to M. Bakavoli, Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Mashhad, 91779, Iran. E-mail: mbakavoli@science1.um.ac.ir or mbakavoli@yahoo.com

convenient procedure for the synthesis of some 1,2,4-triazolo[4,3-b][1,2,4]triazoles **4**. In the present work we have investigated the reaction of aryl halides with N-(3-methylthio-5-substituted-4H-1,2,4-triazol-4-yl)benzenecarboximidamides **3** for the synthesis of some 6-aryl-3-substituted-5H-1,2,4-triazolo[4,3-b][1,2,4]triazoles **4** using microwave irradiation.

Microwave irradiation is a nonconventional energy source whose popularity and synthetic utility in organic chemistry has increased considerably in recent years.^{13–16} The rapid heating induced by such radiation avoids harsh classical conditions and the decomposition of the reagents, leading to the formation of products under mild reaction conditions, thus increasing the yield.

RESULTS AND DISCUSSIONS

Treatment of 4-amino-5-substituted-4H-[1,2,4]triazole-3-thiones **1** with aromatic nitriles in the presence of potassium t-butoxide in t-butanol under reflux gave products identified as N-(3-substituted-5-thioxo-1,5-dihydro-4H-1,2,4-triazol-4-yl)benzenecarboximidamide **2** (Scheme 1 and Table I). Furthermore, methylation of compounds **2** with methyl iodide in the presence of sodium hydroxide at room temperature



- a: R=Me, Ar=C₆H₅
 b: R=Me, Ar=4-Cl-C₆H₄
 c: R=Me, Ar=4-Br-C₆H₄
 d: R=Me, Ar=4-Me-C₆H₄
 e: R=H, Ar=C₆H₅

SCHEME 1

TABLE I N-(3-Substituted-5-thioxo-1,5-dihydro-4H-1,2,4-triazol-4-yl)benzenecarboximidamide (**2a-e**) and N-(3-methylthio-5-substituted-4H-[1,2,4]triazol-4-yl)benzenecarboximidamide (**3a-e**)

Entry	Reaction time (hours)	Yield (%)	m.p. (0°C)	C calcd. (found)	H calcd. (found)	N calcd. (found)	Spectral data
2a	6	88	284–286	51.48 (51.37)	4.75 (4.66)	30.02 (30.15)	¹ H NMR: δ (d ₆ -DMSO), 2.13 (s, 3H, Me), 7.2 (s, 2H, NH ₂), 7.3–8.1 (m, 5H, Ph), 13.14 (s, 1H, NH); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ⁻¹ ; C=N, 1645 cm ⁻¹ ; MS m/z, M ⁺ 233
2b	4	82	298–300	44.86 (44.99)	3.76 (3.62)	26.16 (26.04)	¹ H NMR: δ (d ₆ -DMSO), 2.10 (s, 3H, Me), 7.41 (s, 2H, NH ₂), 7.5–8.1 (m, 4H, P-Cl-C ₆ H ₄), 13.25 (broad, 1H, NH); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ⁻¹ ; C=N, 1643 cm ⁻¹ ; MS m/z, M ⁺ 267.5
2c	6	75	286–288	38.47 (38.31)	3.22 (3.14)	22.43 (22.54)	¹ H NMR: δ (d ₆ -DMSO), 2.09 (s, 3H, Me), 7.39 (s, 2H, NH ₂), 7.5–8.0 (m, 4H, P-Br-C ₆ H ₄), 13.25 (broad, 1H, NH); IR (KBr disc), ν, NH ₂ , 3300–3460 cm ⁻¹ ; C=N, 1643 cm ⁻¹ ; MS m/z, M ⁺ 312
2d	3	86	308–310	53.42 (53.29)	5.30 (5.43)	28.32 (28.21)	¹ H NMR: δ (d ₆ -DMSO), 2.10 (s, 3H, Me), 2.37 (s, 3H, Me), 7.1–8.0 (m, 6H, NH ₂ and aromatic ring), 13.20 (s, 1H, NH); IR (KBr disc), ν, NH ₂ , 3300–3450 cm ⁻¹ ; C=N, 1644 cm ⁻¹ ; MS m/z, M ⁺ 247
2e	6	78	230–232	49.30 (49.43)	4.14 (4.06)	31.94 (31.82)	¹ H NMR: δ (d ₆ -DMSO), 7.1–8.0 (m, 7H, NH ₂ and Ph), 8.30 (s, 1H, N=CH), 13.48 (broad, 1H, NH); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ⁻¹ ; C=N, 1641 cm ⁻¹ ; MS m/z, M ⁺ 219
3a	0.5	81	255–257	53.42 (53.31)	5.30 (5.23)	28.32 (28.43)	¹ H NMR: δ (d ₆ -DMSO), 2.17 (s, 3H, Me), 2.52 (s, 3H, S-Me), 7.30–8.0 (m, 7H, Ph and NH ₂); IR (KBr disc), ν, NH ₂ , 3200–3350 cm ⁻¹ ; C=N, 1665 cm ⁻¹ ; MS m/z, M ⁺ 247

(Continued on next page)

TABLE I N-(3-Substituted-5-thioxo-1,5-dihydro-4H-1,2,4-triazol-4-yl)benzenecarboximidamide (**2a-e**) and N-(3-methylthio-5-substituted-4H-[1,2,4]triazol-4-yl)benzenecarboximidamide (**3a-e**) (Continued)

Entry	Reaction time (hours)	Yield (%)	m.p. (°C)	C		H		N		Spectral data
				calcd. (found)	calcd. (found)	calcd. (found)	calcd. (found)	calcd. (found)	calcd. (found)	
3b	0.5	85	237–239	46.89 (46.75)	4.29 (4.38)	24.85 (24.96)				¹ H NMR: δ (d ₆ -DMSO), 2.16 (s, 3H, Me), 2.52 (s, 3H, S–Me), 7.3–8.1 (m, 6H, NH ₂ and P–Cl–C ₆ H ₄); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ^{–1} ; C≡N, 1656 cm ^{–1} ; MS m/z, M ⁺ 281.5
3c	0.5	71	232–234	40.50 (40.63)	3.71 (3.59)	21.47 (21.39)				¹ H NMR: δ (d ₆ -DMSO), 2.16 (s, 3H, Me), 2.52 (s, 3H, S–Me), 7.3–8.0 (m, 6H, NH ₂ and P–Br–C ₆ H ₄); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ^{–1} ; C≡N, 1654 cm ^{–1} ; MS m/z, M ⁺ 326
3d	0.5	78	251–253	55.15 (55.22)	5.78 (5.84)	26.80 (26.69)				¹ H NMR: δ (d ₆ -DMSO), 2.16 (s, 3H, Me), 2.37 (s, 3H, Me), 2.52 (s, 3H, S–Me), 7.15–7.95 (m, 6H, NH ₂ and aromatic ring); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ^{–1} ; C≡N, 1656 cm ^{–1} ; MS m/z, M ⁺ 261
3e	0.5	74	228–230	51.48 (51.36)	4.75 (4.91)	30.02 (30.11)				¹ H NMR: δ (d ₆ -DMSO), 2.57 (s, 3H, S–Me), 7.25–8.05 (m, 7H, Ph and NH ₂), 8.47 (s, 1H, N=CH); IR (KBr disc), ν, NH ₂ , 3300–3400 cm ^{–1} ; C≡N, 1654 cm ^{–1} ; MS m/z, M ⁺ 233

TABLE II Fast Heterocyclisation Reaction of N-(3-methylthio-5-substituted-4H-[1,2,4]triazol-4-yl)benzenecarboximidamides (**3a–e**) under Microwave Irradiation

Products	R	Ar	Yield (%)	m.p. °C	Lit ¹⁷
4a	CH ₃	C ₆ H ₅	72	227–229	226–228
4b	CH ₃	4-Cl-C ₆ H ₄	77	260–262	263–264
4c	CH ₃	4-Br-C ₆ H ₄	58	276–278	—
4d	CH ₃	4-Me-C ₆ H ₄	68	237–239	238–240
4e	H	C ₆ H ₅	56	215–216	216–218

afforded the corresponding N-(3-methylthio-5-substituted-4H-1,2,4-triazol-4-yl)benzenecarboximidamide **3**. The structures of the products were deduced from their spectra and elemental analysis (Table I). For example, the ¹H NMR spectrum of **3a** in d₆-DMSO shows two singlets at δ 2.17 ppm (3H, Me) and 2.52 ppm (3H, –SMe) and a multiplet at δ 7.30–8.0 ppm (7H, Ph, and NH₂). The IR spectrum of this compound shows two bands in the region 3350–3200 cm^{–1} due to the NH₂ group. The spectral data for compounds **3** are given in Table I.

When compounds **3** in a little amount of N,N-dimethylacetamide were subjected to microwave irradiation for the indicated time (Table III), the cyclic products **4** were obtained (Table II). The structure of the new product **4c** was established from its analytical and spectral data and for known compounds **4a**, **b**, **d**, and **e** by comparison with authentic samples.

It is noteworthy that reaction times required are longer in the absence of microwave irradiation. We investigated the effectiveness of the microwave irradiation and conventional heating for the formation of compounds **4a–e**. The results are summarized in Table III. It can be concluded that the synthesis of compounds **4a–e** under microwave irradiation were 36–72 times faster and the yields are higher than conventional heating methods. This ratio between the reaction time using

TABLE III Comparison of Time and Yields on the Formation of Compounds **4a–e** Using Microwave and Conventional Heating

Product	Conventional heating		Microwave heating			
	t/min (°C)	Yield (%)	Power/W	t/min	Yield (%)	t _c /t _{mw}
4a	360(120)	21	700	7	72	51
4b	360(110)	25	700	5	77	72
4c	360(130)	19	1000	10	58	36
4d	360(120)	23	700	10	68	36
4e	360(130)	14	1000	10	56	36

conventional heating and microwave irradiation (t_c/t_{mw}) shows the microwave heating effect.

In conclusion, we have developed a simple, efficient and fast practical method for one-pot conversion of N-(3-methylthio-5-substituted-4H-1,2,4-triazol-4-yl)benzencarboximidamides into 6-aryl-3-substituted-5H-1,2,4-triazolo[4,3-b][1,2,4]triazoles by applying microwave irradiation in solvent free conditions.

EXPERIMENTAL

Melting points were recorded on an Electrothermal type 9100 melting point apparatus. The IR spectra were obtained on a 4300 Shimadzu Spectrometer. The ^1H NMR (100 MHz) spectra were recorded on a Bruker AC 100 Spectrometer. Mass spectra were obtained from Varian CH-7 instrument at 70 eV. Elemental analyses were performed by Tarbiat Modarres University, Tehran, Iran.

Compounds **4a**, **b**, **d**, **e** were known compounds and their physical data, IR, and NMR spectra were essentially identical with those of authentic samples.

N-(3-Substituted-5-thioxo-1,5-dihydro-4H-1,2,4-triazol-4-yl)benzenecarboximidamide **2a–e**

Typical Procedure

A mixture of 4-amino-3-methyl-4H-[1,2,4]triazole-5-thione **1** (1.30 g, 0.01 mol), benzonitrile (1.23 g, 0.012 mol) and Potassium t-butoxide (2.24 g, 0.02 mol) in t-butanol (60 ml) was heated under reflux for 6 h. After the completion of the reaction, the precipitate was filtered off dissolved in water, and subsequently neutralized by 1N HCl. The crude product was collected and recrystallized from ethanol to give compounds **2a** in high yield (see Table I).

N-(3-Methylthio-5-substituted-4H-[1,2,4]triazol-4-yl)-benzenecarboximidamides **3a–e**

Typical Procedure

To a stirred solution of the foregoing compound **2a** (1.17 g, 0.005 mol) in sodium hydroxide (0.28 g, 0.007 mol) and water (30 ml), methyl iodide (0.71 g, 0.005 mol) was gradually added. The stirring was continued for 0.5 h at room temperature. The crude product was filtered and recrystallized from acetonitrile to give compound **3a** in good yield (see Table I).

6-Aryl-3-substituted-5H-1,2,4-triazolo[4,3-b][1,2,4]-triazoles 4a–e

Typical Procedure

The foregoing compound **3a** (0.49 g, 0.002 mol) in a little amount of N,N-dimethylacetamide were subjected to microwave irradiation for 7 min. The crude product was recrystallized from benzene-ethanol to give compound **4a** in good yield (see Table III).

4c, 6-(4-bromo-phenyl)-3-methyl-5H-[1,2,4]triazolo[4,3-b][1,2,4]triazole, Anal. Calcd for $C_{10}H_8BrN_5$: C, 43.19; H, 2.90; N, 25.18, Found: C, 43.08; H, 2.98; N, 25.09. 1H NMR: δ (d_6 -DMSO), 2.56 (s, 3H, Me), 7.6–8.2 (m, 4H, *p*-Br- C_6H_4), 13.4 (broad, 1H, NH); IR (KBr disc): ν , 3060, 2920, 1630 cm^{-1} ; MS: m/z , M^+ 278.

REFERENCES

- [1] M. T. M. El-Wassimy, M. Abdel-Rahman, A. B. A. G. Ghattas, and O. A. A. Abd Allah, *Phosphorous, Sulfur, Silicon Relat. Elem.*, **70**(1–2), 99 (1992).
- [2] D. Chapman, U.S. 3,598,602 (Cl. 96-109; C 07d, G 03C), 10 Aug 1971, Appl. 07 Oct 1969; C. A. 76: P106442e.
- [3] F. Ishii, Jpn. Kokai Tokkyo Koho JP 07,120,896 [95,120,896] (Cl. G03C7/38), 12 May 1995; C. A. 123; P213018 g.
- [4] H. Fukunaga, N. Furutachi, and G. Furusawa, Eur. Pat. Appl. EP342, 637 (Cl. G03C7/38), 23 Nov 1989, JP Appl. 88/118,075, 17 May 1988; 125PP. C. A. 113: P68234 m.
- [5] T. A. F. O'Mahony, R. N. Butler, and F. L. Scott, *J. Chem. Soc., Perkin Trans. 2*, **10**, 1319 (1972).
- [6] C. Parkanyi, Abdou O. Abdelhamid, and J. C. S. Cheng, *J. Heterocyclic Chem.*, **21**, 1029 (1984).
- [7] K. T. Potts and C. Hirsch, *J. Org. Chem.*, **1**, 143 (1968).
- [8] P. Molina, A. Lorenzo, R. Ma Claramunt, and J. Elguero, *Tetrahedron Lett.*, **25**, 5427 (1984).
- [9] P. Molina, M. Alajarin, A. Ferao, and M. J. Perez de Vega, *J. Chem. Soc., Perkin Trans. 1*, **12**, 2667 (1987).
- [10] P. Molina, M. Alajarin, A. Ferao, A. Lorenzo, and M. J. Vilaplana, *Heterocycles*, **27**, 161 (1988).
- [11] P. Molina, M. Alajarin, and M. J. Perez de Vega, *Heterocycles*, **23**, 2613 (1985).
- [12] P. Molina, M. Alajarin, and M. J. Vilaplana, *Synthesis*, 415 (1983).
- [13] S. Caddick, *Tetrahedron*, **52**, 1403 (1995).
- [14] C. R. Strauss and R. W. Trainor, *Aust. J. Chem.*, **48**, 1665 (1995).
- [15] F. Langa, P. de la Cruz, A. de la hoz, A. Diaz-Ortiz, and E. Diez-Barra, *Contemp. Org. Synth.*, **4**, 373 (1997).
- [16] D. M. P. Mingos and A. G. Whittaker, *Microwave Dielectric Heating Effects in Chemical Synthesis under Extreme or non-Classical Conditions*, edited by R. Van Eldik and C. P. Hubbard (John Wiley and Sons, New York, 1997).
- [17] H. H. Takimoto, G. C. Denault, and S. Hotta, *J. Heterocyclic Chem.*, **3**, 119 (1966).