

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Reactions with Hydrazonoyl Halides 34¹: Synthesis of Some New 2,3-Dihydro-1,3,4-thiadiazoles

Omer S. Abu-Team; Nora M. Rateb; Abdou O. Abdelhamid

Online publication date: 18 June 2010

To cite this Article Abu-Team, Omer S. , Rateb, Nora M. and Abdelhamid, Abdou O.(2003) 'Reactions with Hydrazonoyl Halides 34¹: Synthesis of Some New 2,3-Dihydro-1,3,4-thiadiazoles', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 178: 11, 2363 — 2371

To link to this Article: DOI: 10.1080/714040950

URL: <http://dx.doi.org/10.1080/714040950>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

REACTIONS WITH HYDRAZONOYL HALIDES 34¹: SYNTHESIS OF SOME NEW 2,3-DIHYDRO-1,3,4-THIADIAZOLES

Omer S. Abu-Team, Nora M. Rateb, and Abdou O. Abdelhamid
Cairo University, Giza, Egypt

(Received April 5, 2003; accepted May 21, 2003)

Hydrazonoyl halides 4a–h have been caused to react with each of alkyl carbodithioates (3, 13–17)a,b in the presence of triethylamine to give 2,3-dihydro-1,3,4-thiadiazoles in good yields. Structures of the new compounds were elucidated on the basis of elemental analyses, spectral data, and alternative methods of synthesis whenever possible.

Keywords: 1,3-dipolar cycloaddition; 1,3,4-thiadiazoles; carbodithioates; hydrazonoyl halides; unsymmetrical azines

1,3,4-Thiadiazole and its derivatives have become very useful compound in medicine, agriculture and in many fields of technology.² Although few patented compounds become marketed products, patents do give a rough indication of utility. Some of the technological uses involve dyes, lubricant compositions, optically active liquid crystals, photographic materials, epoxy resins and miner others. A large number of thiadiazoles have been patented in the agricultural filed as herbicides, insecticides, fungicides, bactericides, and anthelmintics. In the medical field they have been patents for almost every disease known to man, and quite a few of them have become commercial products. As an extension of our study^{3–5} and as a part of our program aiming at the synthesis of different 1,3,4-thiadiazoles, here we report on the reactivity of α -keto hydrazonoyl halides toward some alkyl carbodithioates.


RESULTS AND DISCUSSION

The reaction of equimolar amounts of C-ethoxycarbonyl-N-phenylhydrazonoyl chloride (**4a**) with [((1E,3E)-1-aza-4-phenylbuta-1,3-dienyl)amino]methylthiomethane-1-thione (**3a**) in ethanolic

Address correspondence to Abdou O. Abdelhamid, Department of Chemistry, Faculty of Science, Cairo University, Giza 12316, Egypt. E-mail: Abdou@main-scc.cairo.eun.eg



Two possible pathways can account for the formation of **8**: 1) 1,3-Addition of the thiol tautomer **3** to the nitrilium imide **10a**, (which prepared in situ by treatment of hydrazonoyl chloride **4a** with triethylamine), can give the thiohydrazonate ester **11a**, which undergoes nucleophilic cyclization to yield **12a**, which affords **8a** by loss of RSH; and 2) 1,3-cycloaddition of the nitrilium imide **10a** to the C=S of **3a** (or **3b**) can give **12a** directly (cf. Scheme 2).




SCHEME 2

Similarly, the appropriate hydrazonoyl halides **4b–h** react with each of alkyl carbodithioate **3a** (or **3b**), to afford 2,3-dihydro-1,3,4-thiadiazole derivatives **8b–h** respectively.

Hydrazonoyl chloride **4a** reacts with the appropriate ((1E)-1-aza-2-[4-(methylethyl)phenyl]vinyl)amino)methylthiomethane-1-thione (**13a**) or (methane{(1E)-1-aza-2-[4-(methylethyl)phenyl]vinyl}amino)-phenylmethylthio-1-thione (**13b**) in ethanolic triethylamine afforded, in each case, the same isolable product (m.p., mixed m.p., and spectra). Structure **18a** was assigned to the isolated products on the bases of their elemental analyses and spectral data. For example, IR spectra of **18a** revealed absorption band at 1710 cm^{-1} due to the carbonyl group. Its ^1H NMR spectrum showed signals at δ_{H} 1.25 (d, 6H, 2CH_3), 1.40 (t, 3H, CH_3CH_2), 2.94 (hept., 1H, $\text{CH}(\text{CH}_3)_2$), 4.45 (q, 2H, CH_2CH_3), 7.25–8.02 (m, 9H, ArH's) and 8.38 (s, 1H, CH).

By similar route, the appropriate alkyl carbodithioates **13(a, b)** react with the appropriate hydrazonoyl halides **4b–h** to afford unsymmetrical azines **18b–h**, respectively (cf. Scheme 3).

Hydrazonoyl chloride **4a** reacts with each of the appropriate [((1E)-2-(2H-benzo[3,4-d]1,3-dioxolan-5-yl)-1-azavinyl)amino]methylthiomethane-1-thione (**14a**), [((1E)-2-(2H-benzo[3,4-d]1,3-dioxolan-5-yl)-1-azavinyl)amino](phenylmethylthio)methane-1-thione (**14b**), [((1E)-1-aza-2-pyrrol-2-ylvinyl)amino]methylthiomethane-1-thione (**15a**), [((1E)-1-aza-2-pyrrol-2-ylvinyl)amino]phenylmethylthio)methane-1-thione (**15b**), [((1E)-1-aza-2-indol-3-ylvinyl)amino]methylthiomethane-1-thione (**16a**), [((1E)-1-aza-2-indol-3-ylvinyl)amino]phenylmethylthio)methane-1-thione (**16b**), 4-((1E)-2-aza-2-[(methylthiothioxomethyl)-amino]vinyl}benzene-1,2-diol (**17a**), 4-((1E)-2-aza-2-[(phenylmethylthio)thioxomethyl)-amino]vinyl}benzene-1,2-diol (**17b**), to give 2,3-dihydro-1,3,4-thiadiazoles (**19–22a**) respectively. Structures




SCHEME 3

19–22 were confirmed on the basis of elemental analyses, spectral data, and alternative synthesis (cf. Scheme 3 and Experimental).

Analogously, the appropriate hydrazonoyl halides **4b–h**, react with the appropriate **(14–17)a,b** in ethanol containing triethylamine at room temperature to give unsymmetrical azines **(18–22)b–h**, respectively, in a good yields.

Moreover, *C*-phenyl-*N*-phenylhydrazonoyl chloride (**23**) reacts with the appropriate methyl carbodithioates (**3**, **13–17**)a, in ethanolic triethylamine to give the corresponding 2,3-dihydro-1,3,4-thiadiazoles **24a–e**, respectively (cf. Scheme 4).



SCHEME 4

TABLE I Characterization Data of the Newly Synthesized Compounds

Compd. no.	m.p., °C solvent	Yield (%)	Mol. formula (mol. wt.)	% Analyses, calcd./found			
				C	H	N	S
3a	160–162 EtOH	89	C ₁₁ H ₁₂ N ₂ S ₂ (236.36)	55.90 56.20	5.12 5.30	11.85 12.00	27.13 27.30
3b	187–189 EtOH	87	C ₁₇ H ₁₆ N ₂ S ₂ (312.46)	65.35 65.50	5.16 5.30	8.97 9.00	20.52 20.70
8a	118–120 EtOH	85	C ₂₀ H ₁₈ N ₄ O ₂ S (378.46)	63.47 63.60	4.79 4.90	14.80 15.00	8.47 8.60
8b	135–138 ACOH	87	C ₁₉ H ₁₆ N ₄ O ₂ S (364.43)	62.62 62.40	4.43 4.40	15.37 15.10	8.80 8.60
8c	160–162 ACOH	87	C ₁₉ H ₁₆ N ₄ OS (348.43)	65.50 65.30	4.63 4.40	16.08 15.90	9.20 9.00
8d	175–176 ACOH	90	C ₂₄ H ₁₈ N ₄ OS (410.50)	70.22 70.40	4.42 4.60	13.65 13.80	7.81 8.00
8e	190–191 ACOH	80	C ₂₂ H ₁₆ N ₄ O ₂ S (400.46)	65.99 66.00	4.03 4.20	13.99 14.00	8.01 8.20
8f	198–200 ACOH	78	C ₂₂ H ₁₆ N ₄ OS ₂ (416.46)	63.45 63.20	3.87 3.60	13.45 13.20	15.40 15.10
8g	168–170 ACOH	75	C ₂₈ H ₂₀ N ₄ OS (460.56)	73.02 73.00	4.38 4.30	12.16 12.10	6.96 6.90
8h	200–202 ACOH	78	C ₂₄ H ₁₉ N ₅ OS (425.49)	67.75 67.50	4.50 4.30	16.46 16.20	7.54 7.20
13a	106–108 EtOH	80	C ₁₂ H ₁₆ N ₂ S ₂ (252.40)	57.10 57.30	6.39 6.50	11.10 11.30	25.41 25.60
13b	65–67 EtOH	55	C ₁₈ H ₂₀ N ₂ S ₂ (328.50)	65.81 66.00	6.14 6.30	8.53 8.70	19.52 19.70
14a	199–200 EtOH	90	C ₁₀ H ₁₀ N ₂ O ₂ S ₂ (254.33)	47.00 47.20	3.92 4.10	11.01 11.20	25.21 25.40
14b	180–182 ACOH	95	C ₁₆ H ₁₄ N ₂ O ₂ S ₂ (330.43)	58.16 58.30	4.27 4.40	8.48 8.60	19.41 19.60
15a	124–124 EtOH	95	C ₇ H ₉ N ₃ S ₂ (199.30)	42.19 42.40	4.55 4.80	21.08 21.30	32.18 32.30
15b	130–133 EtOH	90	C ₁₃ H ₁₃ N ₃ S ₂ (275.40)	56.70 56.90	4.76 4.90	15.26 15.40	23.29 23.40
16a	193–196 EtOH	95	C ₁₁ H ₁₁ N ₃ S ₂ (249.36)	52.99 53.10	4.45 4.60	16.85 17.00	25.72 25.90
16b	188–190 EtOH	91	C ₁₇ H ₁₅ N ₃ S ₂ (325.46)	62.74 62.50	4.65 4.40	12.91 12.70	19.70 19.50
17a	207–210 EtOH	86	C ₉ H ₁₀ N ₂ O ₂ S ₂ (242.32)	44.61 44.80	4.16 4.30	11.56 11.70	26.46 26.60
17b	175–177 EtOH	85	C ₁₅ H ₁₄ N ₂ O ₂ S ₂ (318.42)	56.58 56.80	4.43 4.30	8.80 8.90	20.14 19.80
18a	85–86 EtOH	85	C ₂₁ H ₂₂ N ₄ O ₂ S (394.50)	63.94 64.00	5.62 5.80	14.20 14.40	8.13 8.30
18b	98–100 EtOH	75	C ₂₀ H ₂₀ N ₄ O ₂ S (380.47)	63.14 63.30	5.30 5.50	14.73 14.90	8.43 8.60
18c	134–135 ACOH	87	C ₂₀ H ₂₀ N ₄ OS (364.47)	65.91 65.70	5.53 5.30	15.37 15.10	8.80 8.60

(Continued on next page)

TABLE I Characterization Data of the Newly Synthesized Compounds
(Continued)

Compd. no.	m.p., °C Solvent	Yield (%)	Mol. formula (mol. wt.)	% Analyses, calcd./found			
				C	H	N	S
18d	145–147	89	C ₂₅ H ₂₂ N ₄ OS (426.54)	70.40	5.20	13.14	7.52
	ACOH			70.60	5.50	13.30	7.70
18e	163–165	85	C ₂₃ H ₂₀ N ₄ O ₂ S (416.51)	66.33	4.84	13.45	7.60
	ACOH			66.10	4.60	13.10	7.40
18f	183–184	79	C ₂₃ H ₂₀ N ₄ OS ₂ (432.56)	63.86	4.66	12.95	14.82
	ACOH			63.80	4.60	12.90	14.80
18g	138–140	81	C ₂₉ H ₂₄ N ₄ OS (476.60)	73.08	5.08	11.76	6.73
	ACOH			73.30	5.20	11.80	6.90
18h	133–135	83	C ₂₅ H ₂₃ N ₅ OS (441.56)	68.00	5.25	15.86	7.26
	ACOH			68.20	5.40	16.00	7.40
19a	180–182	80	C ₁₉ H ₁₆ N ₄ O ₄ S (396.43)	57.57	4.07	14.13	8.09
	EtOH			57.80	4.00	14.30	8.30
19b	173–175	93	C ₁₈ H ₁₄ N ₄ O ₄ S (382.40)	56.54	3.69	14.65	8.34
	ACOH			56.70	3.90	14.90	8.50
19c	162–163	83	C ₁₈ H ₁₄ N ₄ O ₃ S (366.40)	59.01	3.85	15.29	8.75
	ACOH			59.30	4.00	15.40	8.90
19d	166–168	86	C ₂₃ H ₁₆ N ₄ O ₃ S (428.47)	64.47	3.76	13.08	7.48
	ACOH			64.40	3.70	13.00	7.40
19e	243–245	80	C ₂₁ H ₁₄ N ₄ O ₄ S (418.43)	60.25	3.37	13.39	7.66
	ACOH			60.40	3.50	13.50	7.80
19f	169–170	85	C ₂₁ H ₁₄ N ₄ O ₃ S ₂ (434.50)	58.05	3.25	12.98	14.76
	ACOH			58.30	3.40	13.00	15.00
19g	208–210	89	C ₂₇ H ₁₈ N ₄ O ₃ S (478.53)	67.77	3.79	11.71	6.70
	ACOH			67.70	3.70	11.60	6.50
19h	170–172	90	C ₂₃ H ₁₇ N ₅ O ₃ S (443.49)	62.29	3.86	15.79	7.23
	ACOH			62.40	4.00	16.00	7.50
20a	122–124	90	C ₁₆ H ₁₅ N ₅ O ₂ S (341.40)	56.29	4.43	20.51	9.39
	EtOH			56.10	4.40	20.40	9.30
20b	159–160	84	C ₁₅ H ₁₃ N ₅ O ₂ S (327.37)	55.04	4.00	21.39	9.79
	EtOH			55.10	4.10	21.50	10.00
20c	173	93	C ₁₅ H ₁₃ N ₅ OS (311.37)	57.86	4.21	22.49	10.30
	EtOH			57.80	4.20	22.40	10.30
20d	149–150	95	C ₂₀ H ₁₅ N ₅ OS (373.44)	64.33	4.05	18.75	8.59
	EtOH			64.30	4.20	18.90	8.70
20f	160	93	C ₁₈ H ₁₃ N ₅ OS ₂ (379.47)	56.98	3.45	18.46	16.90
	EtOH			67.00	3.40	18.40	16.90
20h	148–150	85	C ₂₀ H ₁₆ N ₆ OS (388.45)	61.84	4.15	21.63	8.25
	EtOH			61.60	4.00	21.40	8.00
21a	218–220	93	C ₂₀ H ₁₇ N ₅ O ₂ S (391.45)	61.37	4.38	17.89	8.19
	EtOH			61.50	4.50	17.90	8.30
21b	236–236	98	C ₁₉ H ₁₅ N ₅ O ₂ S (377.43)	60.46	4.01	18.56	8.49
	EtOH			60.40	4.00	18.50	8.50
21c	218–229	95	C ₁₉ H ₁₅ N ₅ OS (361.43)	63.14	4.18	19.38	8.87
	EtOH			63.00	4.00	19.10	8.60

(Continued on next page)

TABLE I Characterization Data of the Newly Synthesized Compounds
(Continued)

Compd. no.	m.p., °C Solvent	Yield (%)	Mol. formula (mol. wt.)	% Analyses, calcd./found			
				C	H	N	S
21d	248	91	C ₂₄ H ₁₇ N ₅ OS (423.50)	68.07	4.05	16.54	7.57
	ACOH			68.20	4.00	16.70	8.00
21f	305–307	92	C ₂₂ H ₁₅ N ₅ O ₂ S ₂ (429.53)	61.52	3.52	16.30	14.93
	ACOH			61.50	3.50	16.30	15.00
21h	277–280	85	C ₂₄ H ₁₈ N ₆ OS (438.51)	65.74	4.14	19.16	7.31
	ACOH			65.70	4.10	19.10	7.30
22a	200–203	87	C ₁₈ H ₁₆ N ₄ O ₄ S (384.42)	56.24	4.16	14.57	8.34
	EtOH			56.20	4.20	14.50	8.30
22b	195–198	86	C ₁₇ H ₁₄ N ₄ O ₄ S (370.39)	55.13	3.81	15.13	8.66
	EtOH			55.10	3.90	15.30	8.60
22c	217	90	C ₁₇ H ₁₄ N ₄ O ₃ S (354.39)	57.62	3.98	15.81	9.05
	EtOH			57.40	3.70	15.60	9.00
22d	218–220	89	C ₂₂ H ₁₆ N ₄ O ₃ S (416.46)	63.45	3.87	13.45	7.70
	EtOH			63.60	4.00	13.60	7.90
22f	239–241	92	C ₂₀ H ₁₄ N ₄ O ₃ S ₂ (422.49)	56.86	3.34	13.26	15.18
	EtOH			56.80	3.30	13.20	15.10
24a	138–140	85	C ₂₃ H ₁₈ N ₄ S (382.49)	72.23	4.74	14.65	8.38
	EtOH			72.00	4.50	14.40	8.10
24b	113–115	81	C ₂₄ H ₂₂ N ₄ S (398.53)	72.33	5.56	14.06	8.05
	EtOH			72.30	5.50	14.00	8.00
24c	195–197	90	C ₂₂ H ₁₆ N ₄ O ₂ S (400.46)	65.99	4.03	13.99	8.01
	EtOH			66.00	4.00	14.00	8.00
24d	210–212	85	C ₂₃ H ₁₇ N ₅ S (395.49)	69.84	4.33	17.71	8.11
	EtOH			69.90	4.50	17.90	8.30
24e	235–238	87	C ₂₁ H ₁₆ N ₄ O ₂ S (388.45)	64.93	4.15	14.42	8.25
	EtOH			64.90	4.10	14.40	8.20

Structure **24** was confirmed on the basis of elemental analyses, spectral data, and alternative route synthesis. Thus, **23** reacted with the appropriate benzyl carbodithioates (**3**, **13–17b**) in ethanolic triethylamine at room temperature gave identical product (m.p., mixed m.p., and spectra) with the corresponding **24a–e** which obtained before.

EXPERIMENTAL

All melting points were determined on an Electrothermal apparatus and are uncorrected. IR spectra were recorded (KBr discs) on a Shimadzu FT-IR 8201 PC spectrophotometer. ¹H NMR spectra were recorded in CDCl₃ and (CD₃)₂SO on a Varian Gemini 200 MHz spectrometer and chemical shifts were expressed in δ units using TMS as internal reference. Elemental analyses were carried out at the

TABLE II ^1H NMR Spectra of Some Selected Synthesized Compounds

Compd. no.	^1H NMR (δ ppm)
3a	2.66 (s, 3H), 6.94–7.79 (m, 8H) and 10.91 (s, br., 1H).
3b	4.47 (s, 2H), 6.88–8.10 (m, 13H) and 13.2 (s, 1H).
8b	3.92 (s, 3H), 7.95–8.2 (m, 13H).
8c	2.62 (s, 3H), 7.95–7.55 (m, 10H), 7.98–8.02 (d, 2H) and 8.19–8.24 (t, 1H).
8d	6.96–8.41 (m, aromatic protons).
8f	6.96–7.57 (m, 11H), 7.81 (d, 1H), 8.09 (d, 2H), 8.22 (d, 1H) and 8.38 (t, 1H).
13a	1.28 (d, 6H), 2.66 (s, 3H), 2.97–2.97 (sep., 1H), 7.26–7.30 (d, 2H), 7.64–7.68 (d, 2H), 7.90 (s, 1H) and 10.7 (s, br., 1H).
13b	1.27 (d, 6H), 2.90–2.99 (sept., 1H), 4.47 (s, 2H), 7.26–7.72 (m, 9H), 8.10 (s, 1H) and 13.26 (s, br., 1H).
14a	2.7 (s, 3H), 6.11 (s, 2H), 6.99–7.26 (m, 3H), 8.16 (s, 1H) and 13.25 (s, br., 1H).
14b	4.46 (s, 2H), 6.12 (s, 2H), 7.26–7.85 (m, 8H), 8.17 (s, 1H) and 13.26 (s, br. 1H).
18a	1.25 (d, 6H), 1.40 (t, 3H), 2.90–2.97 (sept., 1H), 4.45 (q, 2H), 7.25–7.71 (m, 5H), 7.91 (d, 2H), 7.97 (d, 2H) and 8.38 (s, 1H).
18f	1.25 (d, 6H), 2.94 (sep., 1H), 7.22–8.12 (m, 12H) and 8.41 (s, 1H).
19a	1.43 (t, 3H), 4.45 (q, 2H), 6.02 (s, 2H), 6.86–8.00 (m, 8H) and 8.30 (s, 1H).
19f	6.02 (s, 2H), 6.80–8.40 (m, 12H).
24a	7.11–7.51 (m, 15H), 7.96–8.00 (d, 2H) and 8.19 (t, 1H).
24b	1.28 (d, 6H), 2.94 (sep., 1H), 7.22–8.12 (m, 14H) and 8.31 (s, 1H).
24c	6.01 (s, 2H), 7.22–8.12 (m, 13H) and 8.35 (s, 1H).

Microanalytical Center of the University of Cairo, Giza, Egypt. Hydrazonoyl halides **4a–h** and **23** were prepared according to the literature.^{7–14}

Synthesis of Alkyl Carbodithioates **3a,b**, and **(13–17)a,b**

An appropriate cinnamaldehyde, cumenaldehyde, piperenal, pyrrole-2-carboxaldehyde, 3-formylindole, or 2,4-dihydroxybenzaldehyde (0.01 mol) was added to solution of the appropriate alkyl hydrazinecarbodithioates^{15,16} (0.01 mol) in 2-propanol (50 ml) while stirring at room temperature. The reaction mixture was stirred for 2 h. The solid so formed was collected and crystallized from ethanol to give **3a,b** and **(13–17)a,b**, yields (87–83%) respectively (cf. Tables I and II).

Synthesis of 2,3-Dihydro-1,3,4-thiadiazole Derivatives **8a–g**, **(18–22)a–h**, and **24a–e**

Method A

A mixture of the appropriate alkyl carbodithioates **3a,b** or **(13–17)a,b** (0.005 mol) and the appropriate hydrazonoyl halides **4a–h** or **23** and

triethylamine (0.75 ml, 0.005 mol) in ethanol (20 ml) was stirred for 30 min. The resulting product was collected and crystallized from the proper solvent to give 2,3-dihydro-1,3,4-thiadiazoles (**8**, **18–22**)**a–h** and **24a–e**, respectively, in good yield (cf. Tables I and II).

Method B

A mixture of 5-hydrazono-4-phenyl-4,5-dihydro-1,3,4-thiadiazole-2-carboxylic acid ethyl ester (**9**) (1.3 g, 0.005 mol) and the appropriate of aldehydes (cinnamaldehyde, piperenal, cumenaldehyde, pyrrole-2-carboxaldehyde, 3-formylindole, or 2,4-dihydroxybenzaldehyde) (0.005 mmol) in ethanol (30 ml) was stirred for 2 h. The solid was collected and crystallized from the proper solvent to give in all respects (m.p., mixed m.p., and spectra) with the corresponding prepared in method A.

REFERENCES

- [1] Part 33: A. O. Abdelhamid, S. A. Abdelgawad, and S. F. El-Sharnoby, *Phosphorus, Sulfur, Silicon. Relet. Elem.*, **177**, 2699 (2002).
- [2] D. I. Kornis, *Comprehensive Heterocyclic Chemistry*, edited by A. R. Katritzky, C. W. Rens, and E. F. V. Scriven (Pergamon, Oxford, 1996), Vol. 4.
- [3] A. O. Abdelhamid, M. M. M. Sallam, and S. A. Amer, *Heteroatom Chem.*, **12**, 486 (2001).
- [4] A. O. Abdelhamid, H. F. Zohdi, and N. A. Ali, *Molecules*, **5**, 961 (2001).
- [5] A. O. Abdelhamid, N. M. Rateb, and K. M. Dawood, *Phosphorus, Sulfur, Silicon Relet. Elem.*, **167**, 251 (2000).
- [6] H. F. Zohdi, N. M. Rateb, M. M. M. Sallam, and A. O. Abdelhamid, *J. Chem. Res.*, (S) 742; (M) 3329 (1998).
- [7] G. Fravel, *Bull. Soc. Chim. Fr.*, **31**, 150 (1904).
- [8] N. E. Eweiss and A. Osman, *Tetrahedron Lett.*, 1169 (1979).
- [9] A. S. Shawali and A. Osman, *Tetrahedron*, **27**, 2571 (1971).
- [10] A. S. Shawali and A. O. Abdelhamid, *Bull. Chem. Soc. Jpn.*, **49**, 321 (1976).
- [11] A. O. Abdelhamid and F. H. H. El-Shiatey, *Phosphorus, Sulfur, Silicon and Relat. Elem.*, **39**, 45 (1988).
- [12] A. O. Abdelhamid, F. A. Attaby, F. A. Khalifa, and S. S. Ghabrial, *Arch. Pharm. Res.*, **15**, 14 (1992).
- [13] H. M. Hassaneen, A. S. Shawali, N. E. Elwan, and N. M. Abounada, *Sulfur Lett.*, **14**, 41 (1992).
- [14] P. Wolkoff, *Can. J. Chem.*, **53**, 1333 (1975).
- [15] D. L. Klayman, J. F. Bartosevich, T. S. Griffin, C. J. Mason, and J. P. Scovill, *J. Med. Chem.*, **22**, 855 (1979).
- [16] J. Korosi, Ger. Offen. 1, 934, 899 29 Jan. (1970); *Chem. Abstr.*, **72**, 100334s (1970).