

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### Conversion of Thioamides into Their Corresponding Oxygen Analogues Using Silver Carbonate Supported on Celite

B. Movassagh<sup>a</sup>; M. M. Lakouraj<sup>a</sup>; A. Gholami<sup>a</sup>

<sup>a</sup> Razi University, Kermanshah, Iran

**To cite this Article** Movassagh, B. , Lakouraj, M. M. and Gholami, A.(2003) 'Conversion of Thioamides into Their Corresponding Oxygen Analogues Using Silver Carbonate Supported on Celite', Phosphorus, Sulfur, and Silicon and the Related Elements, 178: 9, 1943 — 1947

**To link to this Article:** DOI: 10.1080/10426500390228620

**URL:** <http://dx.doi.org/10.1080/10426500390228620>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## CONVERSION OF THIOAMIDES INTO THEIR CORRESPONDING OXYGEN ANALOGUES USING SILVER CARBONATE SUPPORTED ON CELITE

*B. Movassagh, M. M. Lakouraj, and A. Gholami*  
*Razi University, Kermanshah, Iran*

(Received August 1, 2001; accepted April 24, 2003)

*Silver carbonate supported on celite ( $\text{Ag}_2\text{CO}_3/\text{Celite}$ ) is used as a mild heterogeneous reagent for conversion of variety of thioamides into their corresponding amides in acetonitrile at room temperature.*

**Keywords:** Amides; celite; silver carbonate; thioamides

Amides are important chemical intermediates since they can be hydrolyzed to acids, dehydrated to nitriles, and degraded to amines containing one less carbon atom by the Hofmann reaction. They constitute a group of substances of increasing interest and applications in pharmacology and industry. These compounds are very prevalent in nature, since all peptides and proteins are polymers of the natural  $\alpha$ -amino acids. Several general methods are available for the preparation of amides;<sup>1</sup> among them, the chemical transformation of thioamides into their corresponding oxo-derivatives, i.e., amides, is highly valuable for the nucleic thiobases and thionucleosides.<sup>2</sup> Reported reagents used for the conversion of thiocarbonyl into carbonyl compounds include iodate or bromate in alkaline solutions,<sup>3</sup> sodium peroxide,<sup>4</sup> dimethyl sulfoxide-acids,<sup>5,6</sup> thiophosgene,<sup>7</sup> diaryl telluroxide,<sup>8</sup> dimethyl selenoxide,<sup>9</sup> dimethyl sulfoxide-iodine,<sup>10</sup> benzeneseleninic acid,<sup>11</sup> singlet oxygen,<sup>12</sup> nitric acid,<sup>13</sup> selenium dioxide,<sup>14</sup> soft  $\text{NO}^+$  species,<sup>15</sup> clayfen,<sup>16</sup> mercuric acetate,<sup>17</sup> and trifluoroacetic anhydride.<sup>18</sup> Very recently, a convenient and mild procedure was introduced for conversion of secondary and tertiary thioamides into the corresponding amides using Caro's acid supported on silica gel.<sup>19</sup> Oxidation of thioamides can take several courses, but all involve the sulfur

We thank the Razi University Research Council for financial assistance.

Address correspondence to B. Movassagh, Department of Chemistry, Razi University, Kermanshah 67149, Iran. E-mail: movassagh@hotmail.com

atom; sulfur is removed as sulfate and replaced by oxygen when an oxidant (e.g., iodine or hydrogen peroxide) is used in basic solution.<sup>20</sup>

## RESULTS AND DISCUSSION

Oxidations by  $\text{Ag}_2\text{CO}_3/\text{celite}$  are heterogeneous reactions that take place under essentially neutral conditions. This reagent is used for oxidation of different functionalities.<sup>21</sup> Here we report the use of silver carbonate supported on Celite,  $\text{Ag}_2\text{CO}_3/\text{celite}$ , as a viable new method for effecting this transformation. This supported reagent can be used for the generation of variety of amides from their corresponding thioamides in acetonitrile at room temperature under air atmosphere in high to excellent yields (Table I). Isolation of the product from this oxidation is remarkably simple and consists merely in filtration to remove the spent reagent system, followed by evaporation of the solvent.

This procedure is applicable to a variety of thioamides including N,N-disubstituted-, N-substituted- and unsubstituted thioamides, thiobenzamides, and thioureas by reaction with one molar equivalent of the reagent, except for 1,4-dithiobenzoyl piperazine (entry 13, Table I) for which two molar equivalents of the reagent is used. The highest yield is obtained with N,N-diethyl-4-nitrobenzamide (98%, entry 12, Table I), and the lowest with thioacetamide (77%, entry 1, Table I).

From the results above, it can be concluded that, by using silver carbonate supported on celite, different types of thioamides are converted smoothly to amides in high to excellent yields.

## EXPERIMENTAL

### General

All yields refer to pure isolated products. The amides were characterized by comparison of their spectral (IR,  $^1\text{H}$ -NMR) and physical data (melting point) with those of authentic samples.<sup>10,11,19,22</sup> The starting thioamides were either prepared from the corresponding amides using tetraphosphorous decasulfide ( $\text{P}_4\text{S}_{10}$ )<sup>23</sup> or the modified Willgerodt-Kindler reaction.<sup>24</sup>  $^1\text{H}$ -NMR spectra were recorded at 60 MHz in  $\text{CCl}_4$  and  $\text{CDCl}_3$  using tetramethylsilane (TMS) as internal standard.

### Preparation of Silver Carbonate Supported on Celite<sup>21</sup>

The celite was purified by washing it successively with methanol containing 10% concentrated hydrochloric acid and then with distilled

**TABLE I** Reaction of Thioamides with  $\text{Ag}_2\text{CO}_3/\text{Celite}$  in Acetonitrile

Entry	$\text{R}_1$	$\text{R}_2$	$\text{R}_3$	Ref. for educt.	Reaction time (h)	Yield <sup>a</sup> (%)	m.p. of product (°C)	Ref. for product
1	Me	H	H	25	7.5	77	80	25
2	$\text{NH}_2$	H	H	25	3	85	133	25
3	Ph	H	H	22	2.5	83	126	25
4	Me	H	Ph	26	4.5	83	114	26
5	Ph	H	4-ClC <sub>6</sub> H <sub>4</sub>	19	5.5	84	193	19,25
6	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	H	PhCH <sub>2</sub>	19	18.5	83	142	19
7	PhNH	H	Ph	25	3	96	240	25
8	Ph	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$		19	6.5	97	47	19,25
9	4-ClC <sub>6</sub> H <sub>4</sub>	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$		19	2	85	76	19
10	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Me	Ph	19	3	90	108	19,27
11	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Et	Et	19	4	78	68	19,28
12	4-Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub>	Et	Et	29	5	98	65	30
13		1,4-dithiobenzoyl piperazine		31	2.5	84 <sup>b</sup>	196	32

<sup>a</sup> Isolated yields.<sup>b</sup> Thioamide:reagent system ratio is 1:2.




FIGURE 1

water until neutral; it was dried at 120°C. Purified celite (3 g) was added to a stirred solution of silver nitrate (3.4 g, 20 mmol) in distilled water (20 ml). A solution of sodium carbonate (1.1 g, 10.4 mmol) in distilled water (30 ml) was then added slowly to the homogeneous suspension and stirring was continued for a further 15 min. The yellow-green precipitate, which was formed, was collected by filtration and dried in a vacuum oven for 3 h. The reagent system thus prepared contained about 0.35 g (1.3 mmol) of  $\text{Ag}_2\text{CO}_3$  per gram.

### General Procedure for Conversion of Thioamides Into Amides

A mixture of the thioamide (1 mmol) and  $\text{Ag}_2\text{CO}_3/\text{celite}$  (1 or 2 mmol) in acetonitrile (15 ml) was stirred at room temperature for the period indicated (Table I). Progress of the reaction was followed by TLC (eluent:  $\text{Et}_2\text{O}/\text{CCl}_4$ , 3/1). After completion of the reaction the solids were collected by filtration, washed with excess acetonitrile, and the combined filtrates evaporated. The crude amide was either recrystallized from n-heptane,  $\text{EtOH}/\text{H}_2\text{O}$ , or subjected to column chromatography using silica gel (n-heptane/ $\text{Et}_2\text{O}$ ).

### REFERENCES

- [1] R. C. Atkins and F. A. Carey, *Organic Chemistry* (McGraw-Hill, New York, 1997), p. 362.
- [2] a) R. H. Hall, *The Modified Nucleosides in Nucleic Acids* (Columbia University Press, New York and London, 1971), p. 347; b) N. K. Kochetkov and E. I. Budovskii, *Organic Chemistry of Nucleic Acids* (Plenum Press, London and New York, 1971), part B, p. 374.
- [3] H. H. Capps and W. M. Dehn, *J. Am. Chem. Soc.*, **54**, 4301 (1932).
- [4] M. J. Kalm, *J. Org. Chem.*, **26**, 2925 (1961).
- [5] M. Mikolajczyk and J. Luczak, *Synthesis*, 491 (1974).
- [6] M. Mikolajczyk and J. Luczak, *Chem. Ind. (London)*, 701 (1974).
- [7] A. Abugar, S. Sharma, and R. N. Iyer, *Indian J. Chem.*, **19B**, 211 (1980).
- [8] S. V. Ley, C. A. Meerholz, and D. H. R. Barton, *Tetrahedron Lett.*, 1785 (1980).
- [9] M. Mikolajczyk and J. Luczak, *J. Org. Chem.*, **43**, 2132 (1978).

- [10] M. Mikolajczyk and J. Luczak, *Synthesis*, 114 (1975).
- [11] N. J. Cussans, S. V. Ley, and D. H. R. Barton, *J. Chem. Soc., Perkin Trans. 1*, 1650 (1980).
- [12] J. E. Gano and S. Atik, *Tetrahedron Lett.*, 4635 (1979).
- [13] A. Husemann, *Liebigs Ann. Chem.*, **126**, 269 (1863).
- [14] R. Boudet, *Bull. Soc. Chim. France*, 846 (1951).
- [15] K. A. Jorgensen, A.-B. A. G. Ghattas, and S.-O. Lawesson, *Tetrahedron*, **38**, 1163 (1982).
- [16] S. Chalais, A. Cornelis, P. Laszlo, and M. Mathy, *Tetrahedron Lett.*, **26**, 2327 (1985).
- [17] F. Challenger, E. A. Mason, E. C. Holdsworth, and R. Emmott, *J. Chem. Soc.*, 292 (1953).
- [18] R. Masuda, M. Hoja, T. Ichi, S. Sasano, T. Kobayashi, and C. Kuroda, *Tetrahedron Lett.*, **32**, 1195 (1991).
- [19] B. Movassagh, M. M. Lakouraj, and K. Ghodrati, *Synth. Commun.*, **30**, 2353 (2000).
- [20] a) H. Wojahn and E. Wenepe, *Arch. Pharm.*, **285**, 375 (1952); b) P. A. S. Smith and R. O. Kan, *J. Org. Chem.*, **29**, 2261 (1964).
- [21] A. McKillop and D. W. Young, *Synthesis*, 401 (1979).
- [22] S. Scheibye, B. S. Pedersen, and S.-O. Lawesson, *Bull. Soc. Chim. Belg.*, **87**, 229 (1978).
- [23] J. W. Scheeren, P. H. J. Ooms, and R. J. F. Nivard, *Synthesis*, 149 (1973).
- [24] K. A. Jensen and C. Pedersen, *Acta Chem. Scand.*, **15**, 1087 (1961).
- [25] R. C. Weast (ed.), *CRC Handbook of Chemistry and Physics* (CRC Press Inc., FL, 1987).
- [26] S. Raucher and P. Klein, *Tetrahedron Lett.*, **31**, 4061 (1980).
- [27] S. I. Lur'e, *Zhur. Obshchei Khim. (J. Gen. Chem.)*, **18**, 1517 (1948).
- [28] H. Erlenmeyer, J. P. Jung, and E. Sorkin, *Helv. Chim. Acta*, **29**, 1960 (1946).
- [29] K. Ghodrati, M.Sc. Thesis, Razi University (2000).
- [30] D. Braun, W. Rettig, S. Delmond, J. -F. Letard, and R. Lapouyad, *J. Phys. Chem. A*, **101**, 6830 (1997); *C. A.*, **127**, 197625c (1997).
- [31] P. -Y. Ding, R. -T. Li, and M. -S. Cai, *Synth. Commun.*, **27**, 973 (1997).
- [32] H. S. Mosher, J. Cornell, Jr., O. L. Stafford, and T. Roe, Jr., *J. Amer. Chem. Soc.*, **75**, 4949 (1953).