

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

## Silica Chloride as a Mild and Efficient Reagent for Acetylation of Alcohols

Farhad Shirini<sup>a</sup>; Mohammad Ali Zolfigol<sup>b</sup>; Mahroo Khaleghi<sup>a</sup>

<sup>a</sup> Guilan University, Rasht, Iran <sup>b</sup> Bu-Ali Sina University, Hamadan, Iran

**To cite this Article** Shirini, Farhad , Zolfigol, Mohammad Ali and Khaleghi, Mahroo(2003) 'Silica Chloride as a Mild and Efficient Reagent for Acetylation of Alcohols', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 178: 9, 1999 — 2002

**To link to this Article:** DOI: 10.1080/10426500390228684

**URL:** <http://dx.doi.org/10.1080/10426500390228684>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SILICA CHLORIDE AS A MILD AND EFFICIENT REAGENT FOR ACETYLATION OF ALCOHOLS

Farhad Shirini,<sup>a</sup> Mohammad Ali Zolfigol,<sup>b</sup>  
and Mahroo Khaleghi<sup>a</sup>

Guilan University, Rasht<sup>a</sup> and Bu-Ali Sina University,  
Hamadan, Iran<sup>b</sup>

(Received February 18, 2003)

*A variety of alcohols underwent acetylation with acetic anhydride in the presence of silica chloride. All reactions were performed at room temperature and under completely heterogeneous conditions in good to high yields.*

**Keywords:** Acetylation; alcohol; esterification; heterogeneous conditions; silica chloride

Functional group protection and deprotection is important in synthetic organic chemistry.<sup>1</sup> Among protecting groups for alcohols, the esters are the most important with acetate being the simplest and easiest of all. Acetylation<sup>2</sup> most commonly is performed using<sup>3,4</sup> reagents such as acetic anhydride or acetyl chloride in the presence of triethyl amine or pyridine as a catalyst.<sup>5</sup> 4-(Dimethylamino)pyridine (DMAP) is known to increase the rate of acetylation when used as a cocatalyst.<sup>6</sup>


Tributylphosphine has been introduced as a less basic catalyst for acetylation of alcohols.<sup>7</sup> In addition to the above catalysts, protic or Lewis acids,<sup>8–14</sup> or some times solid acid catalysts<sup>15–17</sup> are known to catalyze the acetylation of alcohols with acid anhydrides. Recently, scandium triflate has been used for the efficient acetylation of alcohols.<sup>18</sup> Most recently, the use of bismuth as a milder acetylation catalyst than scandium triflate has been reported.<sup>19</sup> However most of the reported methods suffer from one or more of the following disadvantages: long reaction time, vigorous reaction conditions, the occurrence of side

We appreciate the financial support of this work by the Guilan University Research Council.

Address correspondence to Farhad Shirini, Department of Chemistry, Faculty of Science, Guilan University, Rasht, Iran. E-mail: shirini@guilan.ac.ir

reactions, unavailability of the reagents, as well as poor yields of the desired product. Thus, there still is a demand to develop new and mild methods for the acetylation of alcohols in the presence of inexpensive and bench top reagents.

Silica chloride has been used as a mild and efficient reagent for a number of organic reactions.<sup>20–22</sup> In this article we disclose a new method for the acetylation of alcohols using acetic anhydride in the presence of silica chloride in n-hexane (Scheme 1).



### SCHEME 1

As shown in Table I, primary and secondary alcohols are acetylated with acetic anhydride in excellent yields in the presence of silica chloride (entries 1–15). Dihydroxy compounds are converted into the corresponding diacetates efficiently (entry 16). Interestingly, hindered

**TABLE I** Acetylation of Alcohols Using Ac<sub>2</sub>O in the Presence of Silica Chloride<sup>a</sup>

Entry	Substrate	Product	Time (min.)	Yield % <sup>b</sup>
1	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> OH	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> OAc	5	90
2	2-BrC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	2-BrC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OAc	5	86
3	4-MeOC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	4-MeOC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OAc	5	85
4	4-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	4-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OAc	5	85
5	4-(Me) <sub>3</sub> CC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	4-(Me) <sub>3</sub> CC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OAc	3	90
6	3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OH	3-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> OAc	5	85
7	C <sub>6</sub> H <sub>5</sub> CH(OH)CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub> CH(OAc)CH <sub>3</sub>	5	92
8	C <sub>6</sub> H <sub>5</sub> CH(OH)C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> CH(OAc)C <sub>6</sub> H <sub>5</sub>	15	88
9	C <sub>6</sub> H <sub>5</sub> CH=CHCH <sub>2</sub> OH	C <sub>6</sub> H <sub>5</sub> CH=CHCH <sub>2</sub> OAc	5	85
10	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH(OH)CH <sub>3</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH(OAc)CH <sub>3</sub>	10	85
11	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OH	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> OAc	5	80
12	Cyclohexanol	Cyclohexyl acetate	5	85
13	1-octanol	1-octyl acetate	2	82
14	1-butanol	1-butyl acetate	10	80
15	(–) menthol	(–) menthyl acetate	10	86
16	C <sub>6</sub> H <sub>5</sub> CH(OH)CH <sub>2</sub> OH	C <sub>6</sub> H <sub>5</sub> CH(OAc)CH <sub>2</sub> OAc	12	82 <sup>c,d</sup>
17	<i>tert.</i> butyl alcohol	<i>tert.</i> butyl acetate	30	85
18	1-methylcyclohexanol	1-methylcyclohexyl acetate	10	80

<sup>a</sup>Products were characterized by their physical constants, comparison with authentic samples and IR and NMR spectroscopy.

<sup>b</sup>Isolated yield.

<sup>c</sup>Yield refers to isolated pure diacetate.

<sup>d</sup>2.5 mmol of acetic anhydride was used.

tertiary alcohols such as *tert*-butyl alcohol and 1-methylcyclohexanol (entries 17 and 18) also are acetylated in excellent yields in the presence of this reagent. The acetylation reactions were performed under mild and completely heterogeneous conditions at room temperature and take place with good to high yields.

In conclusion, we have shown that silica chloride is a very efficient and versatile reagent for acetylation of alcohols. One of the advantages of this method is that even hindered substrates can be acetylated in high yields under mild reaction conditions. The actual role of silica chloride is not clear and this subject is under investigation in our laboratory.

## EXPERIMENTAL

### General Procedure for Acetylation of Alcohols Using $\text{Ac}_2\text{O}$ in the Presence of Silica Chloride<sup>23</sup>

A mixture of alcohol (1 mmol), acetic anhydride (1.5 mmol), and silica chloride (0.1 g) in *n*-hexane (3 mL) was stirred at room temperature for the specified time. The progress of the reaction was monitored by TLC or GC. After completion of the reaction, the solvent was evaporated and water was added (10 mL). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  ( $2 \times 15$  mL). The organic layer was separated, washed with saturated  $\text{NaHCO}_3$  ( $2 \times 10$  mL) and water (10 mL), and dried over anhydrous  $\text{MgSO}_4$ . Evaporation of the solvent followed by column chromatography on silica gel afforded the pure acetate.

## REFERENCES

- [1] T. W. Green and P. G. M. Wuts, *Protective Groups in Organic Synthesis* (John Wiley Sons Inc., New York, 1991), 2nd ed.
- [2] E. Haslam, *Tetrahedron*, **36**, 2409 (1980).
- [3] J. Mulzer, *Comprehensive Organic Synthesis*, edited by B. M. Trost and I. Fleming (Oxford, Pergamon Press, 1991), vol. 6, p. 323.
- [4] B. M. Trost, *Angew. Chem., Int. Ed. Engl.*, **34**, 259 (1995).
- [5] G. Strok, T. Takahashi, I. Kawamoto, and T. Suzuki, *J. Am. Chem. Soc.*, **100**, 8272 (1978).
- [6] a) G. Höfle, W. Steglich, and H. Vorbrüggen, *Angew. Chem., Int. Ed. Engl.*, **17**, 569 (1978); b) W. Steglich and G. Höfle, *Angew. Chem., Int. Ed. Engl.*, **8**, 981 (1969).
- [7] a) E. Vedejs, N. S. Bennett, L. M. Conn, et al., *J. Org. Chem.*, **58**, 7286 (1993); b) E. Vedejs and S. T. Diver, *J. Am. Chem. Soc.*, **115**, 3358 (1993).
- [8] A. C. Cope and E. C. Herrick, *Org. Synth.*, **4**, 304 (1963).
- [9] T.-S. Jin, Y.-R. Ma, Z.-H. Zhang, and T.-S. Li, *Synth. Commun.*, **28**, 3173 (1998).
- [10] R. Kumareswaran, A. Gupta, and Y. D. Vankar, *Synth. Commun.*, **27**, 277 (1997).

- [11] P. A. Procopiou, S. P. D. Baugh, S. S. Flack, and G. G. A. Inglis, *J. Org. Chem.*, **63**, 2342 (1998).
- [12] P. Saravanan and V. K. Singh, *Tetrahedron Lett.*, **40**, 2611 (1999).
- [13] a) B. Karimi and H. Seradj, *Synlett*, 519 (2001); b) K. K. Chauhan, G. G. Frost, I. Love, and D. Waite, *Synlett*, 1743 (1999).
- [14] a) K. P. R. Kartha and R. A. Field, *Tetrahedron*, **53**, 11753 (1997); b) R. Borah, N. Deka, and J. C. Sarma, *J. Chem. Res.(S)*, 110 (1997).
- [15] T. S. Li and A. X. Li, *Chem. Soc., Perkin Trans. 1*, 1913 (1998).
- [16] R. Ballini, G. Bosica, L. Carloni, R. Maggi, and G. Sartori, *Tetrahedron Lett.*, **39**, 6049 (1998).
- [17] P. M. Bhaskar and D. Loganathan, *Tetrahedron Lett.*, **39**, 2215 (1998).
- [18] K. Ishihara, M. Kubota, H. Kurihara, and H. Yamamoto, *J. Org. Chem.*, **61**, 4560 (1996).
- [19] A. Orita, C. Tanahashi, A. Kakuda, and J. Otera, *Angew. Chem., Int. Ed. Engl.*, **39**, 2877 (2000).
- [20] H. Firouzabadi, N. Iranpoor, B. Karimi, and H. Hazarkhani, *Synlett*, 263 (2000).
- [21] M. A. Zolfigol, F. Shirini, and A. Ghorbani Chogamarani, *Synth. Commun.*, **32**, 1809 (2002).
- [22] F. Shirini, M. A. Zolfigol, M. Khaleghi, and I. Mohammadpoor-Baltork, *Synth. Commun.* (in press).
- [23] M. A. Zolfigol, M. Torabi, and S. E. Mallakpour, *Tetrahedron*, **57**, 8381 (2001).