

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

OXIDATIVE DEPROTECTION OF TRIMETHYLSILYL ETHERS WITH HEXAMETHYLENETETRAMINEBROMINE SUPPORTED ONTO SILICA GEL UNDER MICROWAVE IRRADIATION IN SOLVENTLESS SYSTEM

Mahboobeh Behnam Dehkordi^a; Majid M. Heravi^b; Parviz Khosrofar^a; Ahmad Ziafat^a; Mitra Ghassemzadeh^c

^a Department of Chemistry, Ferdowsi University of Mashhad, Mashhad, Iran ^b Department of Chemistry, School of Sciences, Azzahra University, Tehran, Iran ^c Chemistry & Chemical Engineering Research Center of Iran, Tehran, Iran

Online publication date: 16 August 2010

To cite this Article Dehkordi, Mahboobeh Behnam , Heravi, Majid M. , Khosrofar, Parviz , Ziafat, Ahmad and Ghassemzadeh, Mitra(2004) 'OXIDATIVE DEPROTECTION OF TRIMETHYLSILYL ETHERS WITH HEXAMETHYLENETETRAMINEBROMINE SUPPORTED ONTO SILICA GEL UNDER MICROWAVE IRRADIATION IN SOLVENTLESS SYSTEM', Phosphorus, Sulfur, and Silicon and the Related Elements, 179: 12, 2595 — 2598

To link to this Article: DOI: 10.1080/10426500490494697

URL: <http://dx.doi.org/10.1080/10426500490494697>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

OXIDATIVE DEPROTECTION OF TRIMETHYLSILYL ETHERS WITH HEXAMETHYLENETETRAMINE-BROMINE SUPPORTED ONTO SILICA GEL UNDER MICROWAVE IRRADIATION IN SOLVENTLESS SYSTEM

Mahboobeh Behnam Dehkordi,^a Majid M. Heravi,^b
Parviz Khosrofar,^a Ahmad Ziafat,^a and Mitra Ghassemzadeh^c
Department of Chemistry, Ferdowsi University of Mashhad,
Mashhad, Iran;^a Department of Chemistry, School of Sciences,
Azzahra University, Vanak, Tehran, Iran;^b and Chemistry &
Chemical Engineering Research Center of Iran, Tehran, Iran^c

(Received May 13, 2004; accepted May 13, 2004)

Hexamethylenetetramine-bromine supported onto silica gel rapidly converts trimethylsilyl ethers to their corresponding carbonyl compounds under microwave irradiation in solventless system.

Keywords: Deprotection; hexamethylenetetramine-bromine; microwave irradiation; oxidation

INTRODUCTION

The protection of certain functional groups and the deprotection of the protected derivatives constitute important processes in the multistage synthetic chemistry of polyfunctional molecules, including the total synthesis of natural products.¹ One of the most useful and convenient methods for protection of hydroxy groups is their transformation to trialkylsilyl ethers. Direct oxidation of trimethylsilyl ethers to the corresponding aldehydes and ketones is of great relevance to organic synthesis.

Although various methods have been reported for such a transformation,² some of them show limitations such as the requirement

Partial financial support from Research Council of Ferdowsi University Mashhad is gratefully acknowledged.

Address correspondence to Majid M. Heravi, Department of Chemistry, School of Sciences, Azzahra University, Vanak, Tehran, Iran. E-mail: mmheravi@azzahra.ac.ir

of aqueous reaction conditions, use of expensive reagents, long reaction times, tedious workup, and low yield of products. Therefore, presentation of new methods and inexpensive and environmentally friendly reagents for such functional group transformation is still in much demand. Hexamethylenetetramine-bromine is an inexpensive reagent, which has been reported as an oxidizing agent.³ We have recently successfully used this oxidant in deoximation and desemicarbonization reactions.⁴

In recent years, the organic reactions that are assisted by microwave,⁵ especially under solventless systems,⁶ have attracted attention because of their enhanced selectivity, mild reaction conditions, and associated ease of manipulation.

In continuation of our ongoing efforts in microwave-assisted reactions,⁷ herein we wish to report a simple oxidative deprotection of trimethylsilyl ethers with hexamethylenetetramine-bromine supported onto silica gel in solventless system, which is accelerated by microwave irradiations.

In the absence of solid support, the attempted oxidative deprotection of silyl ethers failed with hexamethylenetetramine-bromine under thermal as well as microwave irradiation even after prolonged reaction periods. In view of established beneficial effects of the reagents on minerals,⁸ we used silica gel as a support for hexamethylenetetramine-bromine. Hexamethylenetetramine-bromine was mixed intimately with an equivalent weight of silica gel. When this supported reagent was mixed thoroughly with benzylsilyl ether using a spatula in a beaker and placed in microwave oven the reaction was completed in 2 min as confirmed by thin layer chromatography (TLC; hexane:ethyl acetate, 8:2; see Table I Entry 2).

To establish generality of the method, trimethylsilyl ethers of various alcohols were used under the above condition.


TABLE I Oxidative Deprotection of Trimethylsilyl Ethers with Hexamethylene-tetramine-bromine Supported onto Silica Gel under Microwave Irradiation

Entry	Substrate	Time (s)	Product	Yield ^a (%)
1	PhCH ₂ OSiMe ₃	120	PhCHO	93
2	Cyclohexyltrimethylsilyl ether	120	Cyclohexanone	88
3	<i>n</i> -CH ₃ (CH ₂) ₆ CH ₂ OSiMe ₃	120	CH ₃ (CH ₂) ₆ CHO	89
4	CH ₃ (CH ₂) ₅ CH ₂ OSiMe ₃	150	CH ₃ (CH ₂) ₅ CHO	72
5	(-)-Menthylsilyl ether	180	Menthone	71
6	Cholesteryltrimethylsilyl ether	150	Cholostenone	67

^aYields refer to isolated product.

In conclusion, we have developed a solid-state method for facile direct oxidative deprotection of trimethylsilyl ethers using hexamethylenetetramine-bromine supported onto silica gel under microwave irradiation in solventless system.

The reagent could be readily prepared by adding bromine to a chloroform solution of commercially available amine. This homogenous non-hygroscopic solid is very stable at room temperature and is not affected by ordinary exposure to light, air, or water, and it has no offensive odor of bromine or amine (Scheme 1).



SCHEME 1

EXPERIMENTAL

All compounds are known and characterized with their physical data by comparison with those of authentic samples. In view of the hazard associated with an oxidant and evolution of gases during the reaction, caution is recommended for its use at elevated temperature. We suggest that the microwave oven be operated carefully and for a shorter time due to possibility of a high localized temperature.

Oxidative Deprotection of Trimethylsilyl Ethers: General Procedure

Hexamethylenetetramine-bromide supported on silica gel (60 mesh, 2 mmol) was mixed thoroughly with the appropriate trimethylsilyl ether (1 mmol) in a beaker. The beaker was placed in a microwave oven and irradiated for specific time (Table I). The progress of reaction was monitored by TLC. The crude CH_2Cl_2 (5 ml) was added and filtered off. The filtrate was evaporated to dryness to afford the corresponding carbonyl compound (Table I).

REFERENCES

- [1] T. G. Greene and P. G. M. Wuts, *Protection Groups in Organic Synthesis* (John Wiley & Sons Inc., New York, 1991) 2nd ed.

- [2] a) M. M. Heravi, D. Ajami, K. Tabar-Hydar, and M. M. Mojtahedi, *J. Chem. Res.*, 620 (1998); b) M. M. Heravi, D. Ajami, and M. Ghassemzadeh, *Synthesis*, **3**, 393 (1999); c) M. M. Heravi, D. Ajami, and K. Tabar-Hydar, *Synth. Commun.*, **29**, 1009 (1999); d) M. M. Heravi, Y. Sh. Beheshtiha, H. A. Oskooie, R. Hekmat-Shoar, and M. Khalilpoor, *Phosphorus, Sulfur, and Silicon*, **161**, 251 (2000) and references cited therein.
- [3] a) I. Yavari and A. Shaabani, *J. Chem. Res.*, 274 (1994); b) M. M. Heravi, D. Ajami, and M. A. Nooshabadi, *Iran J. Chem. Eng.*, **18(2)**, 88 (1999).
- [4] a) Y. Sh. Beheshtiha, M. M. Heravi, N. Sarmad, and R. Hekmat-Shoar, *J. Chem. Res.*, 76 (2001); b) M. A. Nooshabadi, K. Aghapoor, M. Tajbakhsh, and M. M. Heravi, *J. Sci. I. R. Iran*, **12**, 33 (1999).
- [5] a) R. A. Abramovich, *Org. Prep. Proceed. Int.*, **23**, 683 (1991); b) A. G. Whittaker and D. M. P. Mingas, *J. Microwave Power Electromagn. Energy*, **29**, 195 (1994).
- [6] R. S. Varma, *Green Chem.*, 43 (1999).
- [7] a) M. M. Heravi, D. Ajami, M. M. Mojtahedi, and M. Ghassemzadeh, *Tetrahedron Lett.*, **40**, 651 (1999); b) M. M. Heravi, D. Ajami, K. Aghapoor, and M. Ghassemzadeh, *J. Chem. Soc., Chem. Commun.*, 833 (1999).