

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### SYNTHESIS AND STRUCTURE OF A NOVEL SYMMETRICALLY SUBSTITUTED PHOSPHATRIPTYCENE OXIDE

Junji Kobayashi<sup>a</sup>; Tomohiro Agou<sup>a</sup>; Takayuki Kawashima<sup>a</sup>

<sup>a</sup> University of Tokyo, Tokyo, Japan

Online publication date: 12 August 2010

**To cite this Article** Kobayashi, Junji , Agou, Tomohiro and Kawashima, Takayuki(2004) 'SYNTHESIS AND STRUCTURE OF A NOVEL SYMMETRICALLY SUBSTITUTED PHOSPHATRIPTYCENE OXIDE', Phosphorus, Sulfur, and Silicon and the Related Elements, 179: 4, 959 — 960

**To link to this Article:** DOI: 10.1080/10426500490429086

**URL:** <http://dx.doi.org/10.1080/10426500490429086>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SYNTHESIS AND STRUCTURE OF A NOVEL SYMMETRICALLY SUBSTITUTED PHOSPHATRIPTYCENE OXIDE


*Junji Kobayashi, Tomohiro Agou, and Takayuki Kawashima*  
*University of Tokyo, Tokyo, Japan*

(Received August 17, 2003; accepted October 3, 2003)

*Keywords:* Derivatives; phosphatriptycene oxide

Heterotriptycenes bearing one or two heteroatoms at bridgehead positions have been widely investigated, and phosphatriptycenes are one of attractive targets because electronic and structural properties of the bridgehead can be studied by  $^{31}\text{P}$  NMR. Since the first synthesis of a phosphatriptycene in 1974,<sup>1</sup> there have been two other examples reported.<sup>2</sup> However, their synthetic routes were complicated and not suitable in the case of multifunctionalized substrates. Here we report the synthesis of phosphatriptycene oxide **1** from tris(3-methoxyphenyl)phosphine oxide **2** in moderate yield by two steps.

Direct lithiation of **2** with *t*-BuLi, followed by the reaction with diphenyl carbonate gave the corresponding phenoxycarbonyl derivative **3** in 29% yield. Phosphatriptycene oxide **1** was synthesized in 52% yield via intramolecular cyclization by treatment of **3** with 2 equivalents of lithium diisopropylamide.



SCHEME 1

The study was partially supported by the Grant in Aid for the 21st Century COE Program for Frontiers in Fundamental Chemistry.

Address correspondence to Junji Kobayashi, Department of Chemistry, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. E-mail: jkoba@chem.s.u-tokyo.ac.jp

$^{31}\text{P}$  NMR signal of **1** was observed at  $\delta_{\text{P}}$  2.0, which is 28 ppm upfield shifted compared to that of **2** ( $\delta_{\text{P}}$  30.3). Such an upfield shift attributable to the distortion around the phosphorus atom was also observed in the reported phosphatriptycenes. Its crystal structure was determined by x-ray crystallographic analysis.

Phosphatriptycene oxide **1** easily was converted to the corresponding phosphatripticene and other chalcogenides in good yields.

## REFERENCES

- [1] C. Jongsma, J. P. de Kleijn, and F. Bickelhaupt, *Tetrahedron*, **30**, 3465 (1974).
- [2] a) D. Hellwinkel, W. Schenk, and W. Blaicher, *Chem. Ber.*, **111**, 1798 (1978); b) K. G. Weinberg and E. B. Whipple, *J. Am. Chem. Soc.*, **93**, 1801 (1971).