

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### SYNTHESIS OF SOME DITHIOCARBAMATE DERIVATIVES AND THEIR ANTIMICROBIAL ACTIVITY

Zafer Asim Kaplancılı<sup>a</sup>; Gülhan Turan-Zitouni<sup>a</sup>; Gilbert Revial<sup>b</sup>; Gökalp Işcan<sup>a</sup>

<sup>a</sup> Anadolu University, Eskisehir, Turkey <sup>b</sup> CNRS, ESPCI, Paris, France

Online publication date: 16 August 2010

**To cite this Article** Kaplancılı, Zafer Asim , Turan-Zitouni, Gülhan , Revial, Gilbert and Işcan, Gökalp(2004) 'SYNTHESIS OF SOME DITHIOCARBAMATE DERIVATIVES AND THEIR ANTIMICROBIAL ACTIVITY', Phosphorus, Sulfur, and Silicon and the Related Elements, 179: 7, 1449 — 1454

**To link to this Article:** DOI: 10.1080/10426500490463664

**URL:** <http://dx.doi.org/10.1080/10426500490463664>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SYNTHESIS OF SOME DITHIOCARBAMATE DERIVATIVES AND THEIR ANTIMICROBIAL ACTIVITY

Zafer Asım Kaplancikli,<sup>a</sup> Gülhan Turan-Zitouni,<sup>a</sup>  
Gilbert Revial,<sup>b</sup> and Gökalp Işcan<sup>a</sup>  
Anadolu University, Eskisehir, Turkey,<sup>a</sup> and CNRS, ESPCI,  
Paris, France<sup>b</sup>

(Received October 28, 2003; in final form November 25, 2003)

*Some new, 2-[(N-substituted aminothiocabonylthio)acetyl]aminothiazole, N-substituted aminothiocabonylthioacetylaminodiphenylmethane and 9-[(N-substituted aminothiocabonylthio)acetyl]aminofluorene derivatives were synthesized by reacting 2-(chloroacetyl)aminothiazole, chloroacetylaminodiphenylmethane, and 9-(chloroacetyl)aminofluorene with secondary amine dithiocarbamate derivatives in acetone respectively. The structure elucidation of the compounds was performed by IR, <sup>1</sup>H-NMR, and FAB<sup>+</sup>-MS spectral data. The substances were tested for their antimicrobial activity.*

**Keywords:** Antimicrobial activity; diphenylmethane; dithiocarbamate; fluorene; thiazole

It is well known that, N-mono and N,N-di substituted dithiocarbamate derivatives show antibacterial, antiviral and antifungal activities.<sup>1–7</sup>

The structure-activity relationship study revealed that the antibacterial activity on thiocabonyl aromatic compounds was significantly affected by the lipophilicity, that is obtained by thiocabonyl moiety, especially the calculated log P value and the balance between hydrophilic substituent and hydrophobic substituent on the aromatic compounds.<sup>2</sup>

Some thiocabonyl aromatics that were synthesized inspired by the above mentioned rationale were found to possess good in vitro antibacterial activity against gram-positive bacteria.<sup>2</sup>

In view of these observations, some aromatic compounds, i.e., 2-(chloroacetyl)-aminothiazole, chloroacetylaminodiphenylmethane and

Address correspondence to Zafer Asım Kaplancikli, Anadolu University, Faculty of Pharmacy, 26470 Eskisehir, Turkey. E-mail: zakaplan@anadolu.edu.tr

9-(chloroacetyl)aminofluorene were reacted with dithiocarbamates and these compounds were tested for their antibacterial and antifungal activities.

## RESULTS AND DISCUSSION

### Chemistry

The present report deals with the synthesis of 2-[(N-substituted aminothiocabonylthio)-acetyl]aminothiazole (**5a–c**), N-substituted aminothiocabonylthioacetylaminodiphenyl-methane (**6a–c**) and 9-[(N-substituted aminothiocabonylthio)acetyl]aminofluorene (**7a–d**).

The 2-(chloroacetyl)aminothiazole (**2**), chloroacetylaminodiphenyl-methane (**3**) 9-(chloroacetyl)aminofluorene (**4**) were prepared as starting materials in accordance with the method described in the literature.<sup>1,7</sup>

The reaction of chloroacetylamine (**2**, **3**, **4**) and appropriate secondary amine-dithiocarbamatepotassium (**1**) in acetone, gave the 2-[(N-substituted aminothiocabonyl-thio)acetyl]aminothiazole (**5a–c**) N-substituted aminothiocabonylthioacetylaminodiphenyl-methane (**6a–c**) 9-[(N-substituted aminothiocabonylthio)acetyl]aminofluorene (**7a–d**) (Table I, Scheme 1).


Analytical and spectral data (IR, <sup>1</sup>H-NMR, FAB<sup>+</sup>-MS) confirmed the structure of the compounds (see also Table I).

In the IR spectra, some significant stretching bands due to C=O, C=S, N–H were observed at 1640–1680 cm<sup>−1</sup>, 1225–1255 cm<sup>−1</sup>, 3270–3080 cm<sup>−1</sup> respectively.

In the <sup>1</sup>H-NMR spectra, the signal due to COCH<sub>2</sub> methylene protons, present in all compounds, appeared at 4.15–4.40 ppm, as singlets.

**TABLE I** Some Characterizations of the Compounds

Comp.	R	m.p. (°C)	Yield (%)	Molecular formula	Mol. weight
<b>5a</b>	Pyrrolidine	194	80	C <sub>10</sub> H <sub>13</sub> N <sub>3</sub> OS <sub>3</sub>	287
<b>5b</b>	Piperidine	190	70	C <sub>11</sub> H <sub>15</sub> N <sub>3</sub> OS <sub>3</sub>	301
<b>5c</b>	Morpholine	198	65	C <sub>10</sub> H <sub>13</sub> N <sub>3</sub> O <sub>2</sub> S <sub>3</sub>	303
<b>6a</b>	Pyrrolidine	110	68	C <sub>20</sub> H <sub>22</sub> N <sub>2</sub> OS <sub>2</sub>	370
<b>6b</b>	Piperidine	94	73	C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> OS <sub>2</sub>	384
<b>6c</b>	Morpholine	154	85	C <sub>20</sub> H <sub>22</sub> N <sub>2</sub> O <sub>2</sub> S <sub>2</sub>	386
<b>7a</b>	Pyrrolidine	204	80	C <sub>20</sub> H <sub>20</sub> N <sub>2</sub> OS <sub>2</sub>	368
<b>7b</b>	Piperidine	208	60	C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> OS <sub>2</sub>	382
<b>7c</b>	Morpholine	228	55	C <sub>20</sub> H <sub>20</sub> N <sub>2</sub> O <sub>2</sub> S <sub>2</sub>	384
<b>7d</b>	Thiomorpholine	238	65	C <sub>20</sub> H <sub>20</sub> N <sub>2</sub> OS <sub>3</sub>	400



SCHEME 1

Aliphatic protons of pyrrolidine, piperidine, morpholine and thiomorpholine were observed at the 1.50–4.50 ppm region. Aromatic protons and NH proton were elucidated at expected regions.

## Microbiology

All compounds were evaluated for their antimicrobial properties. The compounds showed moderate inhibitor effects against human pathogenic microorganisms. Especially *E. coli*, *C. albicans* were inhibited by **7c**, with a MIC values of 62.5  $\mu\text{g/ml}$  which is equal to that of the standard antifungal agent. **5c** and **7d** showed significant antibacterial effects against *S. aureus* and *S. typhimurium* (Table II).

## EXPERIMENTAL

### Chemistry

Melting points were determined by using a Gallenkamp apparatus. Spectroscopic data were recorded by the following instruments. IR: Shimadzu IR-435 spectrophotometer;  $^1\text{H}$ -NMR: Bruker 250 MHz spectrometer; MS: fast atom bombardment mass spectra (FAB<sup>+</sup>-MS) were obtained by VG Quattro mass spectrometer.

**TABLE II** Antimicrobial Activities of the Compounds

	5a	5b	5c	6a	6b	6c	7a	7b	7c	7d	A	B
<i>E.coli</i>	500	250	125	250	250	125	250	250	62.5	500	31.25	.
<i>S. aureus</i>	250	250	62.5	125	250	125	250	250	125	500	3.9	.
<i>P. aeruginosa</i>	500	250	500	250	500	125	500	500	250	250	62.5	.
<i>E. aerogenes</i>	250	125	250	250	250	250	250	250	250	250	62.5	.
<i>P. vulgaris</i>	250	250	250	250	125	250	125	250	125	125	15.6	.
<i>S. typhimurium</i>	250	250	500	500	250	250	250	125	250	62.5	31.25	.
<i>C. albicans</i>	250	250	500	250	250	500	250	125	62.5	125	—	62.5

\*MIC ( $\mu\text{g/ml}$ ), A: Chloramphenicol, B: Ketoconazole.

### General Procedure for Synthesis of the Compounds

2-(Chloroacetyl)aminothiazole (**2**), chloroacetylaminodiphenylmethane (**3**), 9-(chloroacetyl)aminofluorene (**4**). The amine (2-thiazolylamine, diphenylmethylamine, and 9-fluorenylamine) (0.01 mmol) and triethylamine (0.01 mmol) were dissolved in benzene (50 ml) with constant stirring. Later, the mixture was cooled in an ice bath, and chloroacetylchloride (0.01 mmol) was added dropwise with stirring. The reaction mixture thus obtained was further agitated for 1 h at room temperature. The precipitate was filtrated, the solvent was evaporated to dryness under reduced pressure and the products were recrystallized from ethanol.<sup>1,7</sup>

2-[(N-Substituted aminothiocarbonylthio)acetyl]aminothiazole (**5a–c**) N-substituted aminothiocarbonylthioacetylaminodiphenylmethane (**6 a–c**), 9-[(N-substituted aminothiocarbonylthio)acetyl]aminofluorene (**7a–d**). A mixture of chloroacetylamine (**2**, **3**, **4**) (0.01 mmol) and appropriate secondary amine dithiocarbamate potassium (**1**) (0.01 mmol) was caused to react in acetone at room temperature for 4 h. The solvent was evaporated, washed with water, and the residue recrystallized from ethanol.

IR (KBr,  $\text{cm}^{-1}$ ): These compounds showed characteristic IR bands at 1640–1680  $\text{cm}^{-1}$  (C=O), 1225–1255  $\text{cm}^{-1}$  (C=S), and 3270–3080  $\text{cm}^{-1}$  (N–H).

**5a:**  $^1\text{H-NMR}$  ( $\delta$  ppm) (DMSO- $d_6$ ): 1,95–2,15 (4H, two p, C<sub>3</sub> and C<sub>4</sub> protons of pyrrolidine), 3,70 and 3,95 (4H, two t, C<sub>2</sub> and C<sub>5</sub> protons of pyrrolidine), 4,35 (2H, s, COCH<sub>2</sub>), 6,95 (1H, d, C<sub>5</sub> proton of thiazole), 7,50 (1H, d, C<sub>4</sub> proton of thiazole), 12,45 (1H, br, NH)

MS (FAB<sup>+</sup>): m/z: 288 [M + 1]

**5b:**  $^1\text{H-NMR}$  ( $\delta$  ppm)(DMSO- $d_6$ ): 1,60–1,80 (6H, br, C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub> protons of piperidine), 3,90 and 4,30 (4H, two br s, C<sub>2</sub> and C<sub>6</sub> protons of piperidine), 4,40 (2H, s, COCH<sub>2</sub>), 6,95 (1H, d, C<sub>5</sub> proton of thiazole), 7,55 (1H, d, C<sub>4</sub> proton of thiazole), 12,40 (1H, br, NH)

MS (FAB<sup>+</sup>): m/z: 302 [M + 1]

**6b:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 1,50–1,90 (6H, m, C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub> protons of piperidine), 3,85 and 4,25 (4H, two br s, C<sub>2</sub> and C<sub>6</sub> protons of piperidine), 4,20 (2H, s, COCH<sub>2</sub>), 6,15 (1H, d, CH), 7,10–7,35 (10H, m, aromatic protons), 9,10 (1H, d, NH)

MS (FAB<sup>+</sup>): m/z: 385 [M + 1]

**6c:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 3,60 (4H, br, C<sub>3</sub> and C<sub>5</sub> protons of morpholine), 3,90 (4H, br, C<sub>2</sub> and C<sub>6</sub> protons of morpholine), 4,15 (2H, s, COCH<sub>2</sub>), 6,10 (1H, d, CH), 7,20–7,35 (10H, m, aromatic protons), 9,10 (1H, d, NH)

MS (FAB<sup>+</sup>): m/z: 387 [M + 1]

**7a:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 1,85–2,15 (4H, two p, C<sub>3</sub> and C<sub>4</sub> protons of pyrrolidine), 3,65 and 3,75 (4H, two t, C<sub>2</sub> and C<sub>5</sub> protons of pyrrolidine), 4,15 (2H, s, COCH<sub>2</sub>), 6,00 (1H, d, C<sub>9</sub> proton of fluorene), 7,20–7,45 (6H, m, aromatic protons), 7,85 (2H, d, C<sub>1</sub> and C<sub>8</sub> protons of fluorene), 7,85 (1H, d, NH)

MS (FAB<sup>+</sup>): m/z: 369 [M + 1]

**7b:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 1,55–1,75 (6H, br, C<sub>3</sub>, C<sub>4</sub> and C<sub>5</sub> protons of piperidine), 3,90 and 4,30 (4H, two br, C<sub>2</sub> and C<sub>6</sub> protons of piperidine), 4,25 (2H, s, COCH<sub>2</sub>), 6,05 (1H, d, C<sub>9</sub> proton of fluorene), 7,25–7,40 (6H, m, aromatic protons), 7,90 (2H, d, C<sub>1</sub> and C<sub>8</sub> protons of fluorene), 8,80 (1H, d, NH)

MS (FAB<sup>+</sup>): m/z: 383 [M + 1]

**7c:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 3,70 (4H, t, C<sub>3</sub> and C<sub>5</sub> protons of morpholine), 4,00 (4H, br, C<sub>2</sub> and C<sub>6</sub> protons of morpholine), 4,25 (2H, s, COCH<sub>2</sub>), 6,10 (1H, d, C<sub>9</sub> proton of fluorene), 7,30–7,60 (6H, m, aromatic protons), 7,90 (2H, d, C<sub>1</sub> and C<sub>8</sub> protons of fluorene), 8,90 (1H, d, NH)

MS (FAB<sup>+</sup>): m/z: 385 [M + 1]

**7d:** <sup>1</sup>H-NMR (δ ppm)(DMSO-*d*<sub>6</sub>): 2,70 (4H, t, C<sub>3</sub> and C<sub>5</sub> protons of thiomorpholine), 4,30 and 4,50 (4H, two br, C<sub>2</sub> and C<sub>6</sub> protons of thiomorpholine), 4,20 (2H, s, COCH<sub>2</sub>), 6,10 (1H, d, C<sub>9</sub> proton of fluorene), 7,25–7,45 (6H, m, aromatic protons), 7,95 (2H, d, C<sub>1</sub> and C<sub>8</sub> protons of fluorene), 8,80 (1H, two d, NH)

MS (FAB<sup>+</sup>): m/z: 401 [M + 1]

## Microbiology

Antibacterial activities of compounds were determined using the tube dilution technique.<sup>8,9</sup> MIC values were calculated as μg/ml. The following were used as a test microorganisms; *Escherichia coli* (ATCC 25922), *Staphylococcus aureus* (ATCC 6538), *Pseudomonas aeruginosa* (ATCC 27853), *Proteus vulgaris* (NRRL B-123), *Enterobacter aerogenes* (NRRL 3567), *Salmonella typhimurium* (NRRL B-4420), and *Candida albicans* (University of Osmangazi, Faculty of Medicine, Eskişehir).

Microdilution broth susceptibility assay was used for the antimicrobial evaluation of the compounds. Stock solutions of the samples were prepared in dimethylsulfoxide. Dilution series using sterile distilled water were prepared from 4 mg/ml to 0.007 mg/ml in micro-test tubes that were transferred to 96-well microtiter plates. Overnight grown bacterial and *C. albicans* suspensions in double-strength Mueller-Hinton broth were standardised to  $10^8$  CFU/ml using McFarland No: 0.5 standard solution. 100  $\mu$ l of each microorganism suspension was then added into the wells. The last well-chain without microorganism was used as a negative control. Sterile distilled water and the medium served as a positive growth control. After incubation at 37°C for 18–24 h the first well without turbidity was determined as the minimal inhibitory concentration (MIC). Chloramphenicol was used as standard antibacterial agent whereas ketoconazole was used as antifungal agent.

## REFERENCES

- [1] G. Turan-Zitouni, P. Chevallet, Y. Robbe, and K. Güven, *Acta Pharm. Turcica*, **41**, 82 (1999).
- [2] R. Tokuyama, Y. Takahashi, M. Tsubouchi, et al., *Chemical and Pharmaceutical Bulletin*, **49**, 353 (2001).
- [3] E. Gaudernak, J. Seipelt, A. Trendl, A. Grassauer, and E. Kuechler, *J. Virology*, **76**, 6004 (2002).
- [4] S. Liao, S. Raung, and C. Chen, *Neuroscience Lett.*, **324**, 133 (2002).
- [5] C. Rafin, E. Veignie, M. Sancholle, et al., *J. Agric. Food Chem.*, **48**, 5283 (2000).
- [6] A. Gürsoy, Ö. Ateş, N. Karali, N. Cesur, and M. Kiraz, *Eur. J. Med. Chem.*, **31**, 643 (1996).
- [7] Ö. Ateş, N. Cesur, H. Guener, and M. Uzun, *Farmaco*, **50**, 361 (1995).
- [8] E. W. Koneman, S. D. Allen, and W. C. Winn, *Color Atlas and Textbook of Diagnostic Microbiology* (Lippincott Raven, Philadelphia, 1997), p. 785.
- [9] G. İşcan, N. Kırimer, M. Kürküoğlu, K. H. C. Başer, and F. Demirci, *J. Agric. Food Chem.*, **50**, 3943 (2002).