

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### CARO'S ACID SUPPORTED ON SILICA GEL, PART VIII: AN EFFICIENT AND SELECTIVE REAGENT FOR CONVERSION OF PHENYLHYDRAZONES AND SEMICARBAZONES TO THE CORRESPONDING CARBONYL COMPOUNDS

Mahmood Tajbakhsh<sup>a</sup>; Moslem-Mansor Lakouraj<sup>a</sup>; Mohammad-Hosein Gholami<sup>a</sup>; Farhad Ramzani-Lehmali<sup>a</sup>

<sup>a</sup> University of Mazandaran, Babolsar, Iran

Online publication date: 16 August 2010

**To cite this Article** Tajbakhsh, Mahmood , Lakouraj, Moslem-Mansor , Gholami, Mohammad-Hosein and Ramzani-Lehmali, Farhad(2004) 'CARO'S ACID SUPPORTED ON SILICA GEL, PART VIII: AN EFFICIENT AND SELECTIVE REAGENT FOR CONVERSION OF PHENYLHYDRAZONES AND SEMICARBAZONES TO THE CORRESPONDING CARBONYL COMPOUNDS', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 179: 9, 1731 – 1736

**To link to this Article:** DOI: 10.1080/10426500490466337

URL: <http://dx.doi.org/10.1080/10426500490466337>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## **CARO'S ACID SUPPORTED ON SILICA GEL, PART VIII: AN EFFICIENT AND SELECTIVE REAGENT FOR CONVERSION OF PHENYLHYDRAZONES AND SEMICARBAZONES TO THE CORRESPONDING CARBONYL COMPOUNDS**

*Mahmood Tajbakhsh, Moslem-Mansor Lakouraj,  
Mohammad-Hosein Gholami, and Farhad Ramzanian-Lehmali  
University of Mazandaran, Babolsar, Iran*

(Received January 6, 2004; accepted March 1, 2004)

*Deprotection of phenylhydrazones and semicarbazones to their parent aldehydes and ketones in high yields has been carried out using caro's acid supported on silica gel as a selective oxidant under mild conditions.*

**Keywords:** 2,4-Di-nitrophenylhydrazones; Caro's acid; deprotection; phenylhydrazones; semicarbazones

### **INTRODUCTION**

Derivatives of carbonyl compounds such as phenylhydrazones and semicarbazones not only are used for the characterization and purification of carbonyl compounds but also play an important role in the protection of carbonyl compounds, as they are highly crystalline and stable compounds. Thus, the regeneration of carbonyl compounds from their derivatives under mild condition is an important process in organic synthetic chemistry.

The classical method for the cleavage of phenylhydrazones and semicarbazones to aldehydes and ketones includes acid hydrolysis, which is not suitable for acid-sensitive compounds.<sup>1</sup>

We are thankful to the Mazandaran University Research Council for the partial support of this work.


Address correspondence to Mahmood Tajbakhsh, Department of Chemistry, School of Sciences, University of Mazandaran, P.O. Box 453, Babolsar, Iran. E-mail: tajbakhsh@umz.ac.ir

Several oxidative deprotection methods have been developed that have some advantages over the classical hydrolysis methods.<sup>2-13</sup> Little attention has been paid to the oxidative cleavage of phenylhydrazones and semicarbazones, and only a few reports are available dealing with the conversion of these derivatives to their corresponding carbonyl compounds.<sup>14-18</sup>

## RESULTS AND DISCUSSION

We have previously reported that Caro's acid supported on silica gel is an efficient reagent for oxidation of organic compounds.<sup>19,20</sup>

We now report a mild and convenient method for oxidative deprotection of semicarbazones and phenylhydrazones in high yields using Caro's acid/silica gel (Scheme 1).




SCHEME 1

Our experiments show that semicarbazones and phenylhydrazones are converted to their corresponding aldehydes and ketones in refluxing dichloromethane. Further oxidation of aldehydes to their carboxylic acid is not observed. This reagent has wide applicability for regeneration of carbonyl compounds from aliphatic and aromatic semicarbazones and phenyl hydrazones (Table I).

It is noteworthy that the 2,4-dinitrophenylhydrazones are not deprotected by this reagent in appropriate yield. These results suggest that this procedure will show considerable selectivity between semicarbazones and phenylhydrazones with 2,4-dinitrophenylhydrazones. This was established by an experiment in which a mixture of an equal amount of the derivatives, semicarbazone of benzaldehyde and 2,4-dinitrophenylhydrazone of *p*-chlorobenzaldehyde, was treated with Caro's acid on silica gel in CH<sub>2</sub>Cl<sub>2</sub> at reflux temperature for 20 min. Working up the reaction mixture showed that only the semicarbazone was selectively oxidized to benzaldehyde, and the 2,4-dinitrophenylhydrazone of *p*-chlorobenzaldehyde remained unchanged (Scheme 2).

Similar competitive reaction between phenylhydrazone of benzaldehyde and 2,4-dinitrophenylhydrazone of *p*-chlorobenzaldehyde




SCHEME 2

resulted only in the oxidation phenylhydrazone of benzaldehyde (Scheme 3).

When we treated  $\alpha$ ,  $\beta$ -unsaturated semicarbazone, only the C=N bond was selectively oxidized to the corresponding carbonyl compound, and the reagent was ineffective in oxidizing C=C double bond (Table I, Entry 8).

In order to show the efficiency of this method we have compared some of the results with relevant ones reported in the literature<sup>14-18</sup> (Table II). As indicated in Table II, when compared to other reagents Caro's acid on silica gel performs this transformation in higher yield, shorter reaction times, and milder conditions.

In conclusion the methodology described here for the regeneration of carbonyl compounds from their corresponding semicarbazones and phenylhydrazones with Caro's acid on silica gel is manipulatively simple, mild, highly selective, cheap, and it avoids disadvantages of other methods.



SCHEME 3

**TABLE I** Regeneration of Carbonyl Compounds from Semicarbazones and Phenylhydrazones with Caro's Acid/Silica Gel in Refluxing CH<sub>2</sub>Cl<sub>2</sub>

Entry	R1	R2	X	Time (min)	Yield <sup>a,b</sup> (%)
1	C <sub>6</sub> H <sub>5</sub>	H	NHCONH <sub>2</sub>	20	100
2	<i>p</i> -MeO C <sub>6</sub> H <sub>4</sub>	H	NHCONH <sub>2</sub>	35	100
3	<i>o,p</i> -(MeO) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	H	NHCONH <sub>2</sub>	5	100
4	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	H	NHCONH <sub>2</sub>	8	100
5	<i>p</i> -BrC <sub>6</sub> H <sub>4</sub>	H	NHCONH <sub>2</sub>	5	100
6	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	H	NHCONH <sub>2</sub>	50	90
7	<i>m</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	H	NHCONH <sub>2</sub>	80	90
8	C <sub>6</sub> H <sub>5</sub> CH=CH	H	NHCONH <sub>2</sub>	39	100
9	C <sub>6</sub> H <sub>5</sub>	CH <sub>3</sub>	NHCONH <sub>2</sub>	5	100
10	C <sub>6</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	NHCONH <sub>2</sub>	7	100
11	CH <sub>3</sub>	CH <sub>3</sub>	NHCONH <sub>2</sub>	4	100
12	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	NHCONH <sub>2</sub>	10	100
13	—(CH <sub>2</sub> ) <sub>4</sub> —		NHCONH <sub>2</sub>	45	90
14	—CH <sub>2</sub> CH <sub>2</sub> CH(C <sub>6</sub> H <sub>5</sub> )CH <sub>2</sub> CH <sub>2</sub> —		NHCONH <sub>2</sub>	25	100
15	Camphor		NHCONH <sub>2</sub>	15	100
16	C <sub>6</sub> H <sub>5</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	120	95
17	<i>p</i> -Me C <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	60	90
18	<i>o</i> -OHC <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	50	90
19	<i>o</i> -ClC <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	75	90
20	<i>m</i> -ClC <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	105	95
21	<i>p</i> -BrC <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	85	95
22	<i>p</i> -NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	H	NHC <sub>6</sub> H <sub>5</sub>	90	90
23	<i>o</i> -OHC <sub>6</sub> H <sub>4</sub>	CH <sub>3</sub>	NHC <sub>6</sub> H <sub>5</sub>	150	90
24	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>	NHC <sub>6</sub> H <sub>5</sub>	125	90

<sup>a</sup>Products were characterized by their physical constants, spectral characteristics (IR, <sup>1</sup>H NMR) and comparison with authentic samples.

<sup>b</sup>Yields are of pure isolated products.

## EXPERIMENTAL

Products were isolated, and their physical data were compared with those of known samples. Solvent was freshly distilled. Phenylhydrazones, 2,4-dinitrophenylhydrazones, and semicarbazones were prepared according to the described procedures.<sup>21</sup>

### General Procedure for Deprotection of Semicarbazones and Phenylhydrazones

In a round-bottomed flask (50 ml) equipped with a condensor and a magnetic stirrer, a mixture of the substrate (1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) and Caro's acid on silica gel (3 mmol) was placed. The reaction mixture was stirred at reflux temperature for the specified time (Table I). The

**TABLE II** Comparison of Some of the Results Obtained by the Desemicarbazonation with Caro's Acid Supported on Silica Gel with Some of Those Obtained by BTPPMS,<sup>a,17</sup> Dowex-50,<sup>16</sup> ACC/alumina,<sup>b,14</sup> CuCl<sub>2</sub>. 2H<sub>2</sub>O,<sup>15</sup> and QDC<sup>c,18</sup>

Entry	Reagent	Substrate	Time (min)	Yield (%)	Reaction condition
1	Caro's acid/ Silica gel	Acetophenone semicarbazone	5	100	CH <sub>2</sub> Cl <sub>2</sub> , Reflux
		Benzaldehyde semicardazone	20	100	
2	BTPPMS	Acetophenone semicarbazone	60	95	MeCN, Reflux
3	Dowex-50	Acetophenone semicarbazone	60	90	H <sub>2</sub> O, Reflux
		Benzaldehyde semicardazone	300	32	
4	ACC/alumina	Acetophenone semicarbazone	360	59	CH <sub>2</sub> Cl <sub>2</sub> , Reflux
		Benzaldehyde semicardazone	240	72	
5	CuCl <sub>2</sub> .2H <sub>2</sub> O	Acetophenone semicarbazone	30	81	MeCN, Reflux
		Benzaldehyde semicardazone	90	51	
6	QDC	Acetophenone semicarbazone	132	77	MeCN, Reflux
		Benzaldehyde semicardazone	135	85	

<sup>a</sup>BTPPMS, Benzyltriphenylphosphonium peroxymonosulfate.

<sup>b</sup>Ammonium chlorochromate.

<sup>c</sup>Quinolinium dichromate.

progress of the reaction was monitored by thin layer chromatography (TLC) or gas chromatography (GC). After completion of the reaction the solid material was filtered and washed with CH<sub>2</sub>Cl<sub>2</sub>, and dried by Na<sub>2</sub>SO<sub>4</sub>. The combined filtrates were evaporated on a rotary evaporator. The resulting crude material was purified on a silica gel plate (eluent: hexane/ethyl acetate, 5:1) to afford the pure carbonyl compound, yield 90–100% (Table I).

## REFERENCES

- [1] E. B. Hersberg, *J. Org. Chem.*, **13**, 542 (1984).
- [2] N. Kirkd and J. Sladec, *Tetrahedron Lett.*, **21**, 651 (1980).
- [3] E. J. Eisenbraun, R. P. Wesely, R. S. Budhram, and B. Dewprashad, *Chem. Ind.*, **14**, 584 (1989).
- [4] R. N. Ram and N. Varshak, *Tetrahedron Lett.*, **32**, 5829 (1991).
- [5] M. Curini, O. Rosati, and E. Pisani, *Synthesis*, **4**, 333 (1996).
- [6] D. Subhas and A. Venkant Narasaiah, *Synth. Commun.*, **30**, 1153 (2000).
- [7] I. Mohammadpoor-Baltork and Sh. Pouranshirvani, *Synth. Commun.*, **26**, 261 (1996).
- [8] H. Firouzabadi, M. Seddighi, A. Ahmadi, and A. R. Sardarian, *Synth. Commun.*, **19**, 3385 (1989).
- [9] G. S. Zhang, D. H. Yang, H. F. Chen, and K. Cai, *Synth. Commun.*, **28**, 2221 (1995).
- [10] G. S. Zhang, D. H. Yang, and M. F. Chen, *Synth. Commun.*, **28**, 3721 (1998).
- [11] H. Firouzabadi, N. Iranpoor, F. Kiaeezadeh, and J. Toofan, *Tetrahedron*, **42**, 719 (1986).

- [12] H. Firouzabadi and I. Mohammadpoor-Baltork, *Synth. Commun.*, **24**, 489 (1994).
- [13] H. Firouzabadi, E. Mottaghinejed, and M. Seddighi, *Synth. Commun.*, **19**, 469 (1989).
- [14] G. S. Zhang, H. Gong, D. H. Yung, and M. F. Chen, *Synth. Commun.*, **29**, 1165 (1999).
- [15] R. N. Ram and K. Varsha, *Tetrahedron Lett.*, **32**, 5829 (1991).
- [16] B. Ranu and D. C. Sarkar, *J. Org. Chem.*, **53**, 878 (1988).
- [17] A. R. Hajipour, S. E. Mallakpour, I. M. Baltork, and H. Adibi, *Synth. Commun.*, **31**, 3401 (2001).
- [18] M. Mirmohammad Sadeghi, I. M. Baltork, M. Azarm, and M. R. Mazidi, *Synth. Commun.*, **31**, 435 (2001).
- [19] B. Movassagh, M. M. Lakouraj, and K. Ghodrati, *Synth. Commun.*, **30**, 3597 (2000).
- [20] B. Movassagh, M. M. Lakouraj, and K. Ghodrati, *Synth. Commun.*, **32**, 847 (2002).
- [21] A. I. A. Vogel, *Textbook of Practical Organic Chemistry*, 4th ed. (Longmans London, 1978).