

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

A Convenient Synthesis of Thioacetates and Thiobenzoates Using Silica-Gel Supported Potassium Thioacetate and Thiobenzoate

Tadashi Aoyama^a; Toshio Takido^a; Mitsuo Kodomari^b

^a Department of Materials and Applied Chemistry, College of Science and Technology, Nihon University, Tokyo, Japan ^b Department of Applied Chemistry, Shibaura Institute of Technology, Tokyo, Japan

To cite this Article Aoyama, Tadashi, Takido, Toshio and Kodomari, Mitsuo(2005) 'A Convenient Synthesis of Thioacetates and Thiobenzoates Using Silica-Gel Supported Potassium Thioacetate and Thiobenzoate', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 180: 5, 1447 – 1448

To link to this Article: DOI: 10.1080/10426500590913023

URL: <http://dx.doi.org/10.1080/10426500590913023>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A Convenient Synthesis of Thioacetates and Thiobenzoates Using Silica-Gel Supported Potassium Thioacetate and Thiobenzoate

Tadashi Aoyama


Toshio Takido

Department of Materials and Applied Chemistry, College of Science and Technology, Nihon University, Tokyo, Japan

Mitsuo Kodomari

Department of Applied Chemistry, Shibaura Institute of Technology, Tokyo, Japan

Recently, many reactions using supported reagents have been reported. We have found that potassium thioacetate can be remarkably activated by support on silica gel, and silica gel-supported potassium thioacetate ($\text{CH}_3\text{COSK/SiO}_2$) reacts with alkyl halides in nonpolar solvents such as benzene to give alkyl thioacetates in good yields.



Benzyl bromide did not react with finely ground potassium thioacetate suspended in benzene at room temperature. In contrast, similar reactions with $\text{CH}_3\text{COSK/SiO}_2$ proceeded easily under similar conditions to afford benzyl thioacetate in 83% yield after 1 h. Silica gel was the most effective among the inorganic supports tested. Aluminas (neutral, acidic, and basic) were also effective but less than silica gel (33–58%). Molecular sieves 4A was much less effective. Kiesegelur, Montmorillonite K10, and charcoals were ineffective for activating potassium thioacetate. The reaction of a series of organic halides with $\text{CH}_3\text{COSK/SiO}_2$ in benzene gave the corresponding thioacetates in high yields. The reaction of benzyl chloride with $\text{CH}_3\text{COSK/SiO}_2$ required more strong conditions than that of benzyl bromide. Substituted benzyl bromides were converted into the corresponding thioacetates in excellent yields

Received July 9, 2004; accepted October 5, 2004.

Address correspondence to Tadashi Aoyama, Department of Materials and Applied Chemistry, College of Science and Technology, Nihon University, Kanda Surugadai, Chiyoda-ku, Tokyo 101-8308, Japan. E-mail: aoyama@chem.cst.nihon-u.ac.jp

at room temperature independent of the substituent. For instance, 4-methyl and 4-nitrobenzyl bromide were converted into the corresponding thioacetates in 98% and 97% yields. Xylylene dibromides produced only disubstituted compound. Mono-substituted compounds were not detected.

The reactivity of *n*-alkyl halides decreased as follows: I > Br > Cl. Octyl thioacetate was obtained quantitatively from octyl iodide at 80°C after 3 h, whereas octyl chloride gave only 32% yield of octyl thioacetate after 24 h. In the case of 1,4-dibromopentane, 4-bromopentyl thioacetate was formed along with a small amount of disubstituted compound, but 4-bromo-1-methylbutyl thioacetate was not detected. Using silica gel-supported potassium thiobenzoate (PhCOSK/SiO₂), the reaction rapidly occurred at room temperature to give the corresponding thiobenzoates in high yields. The reactivity of PhCOSK/SiO₂ was higher than CH₃COSK/SiO₂. Octyl bromide reacted with CH₃COSK/SiO₂ to afford the octyl thioacetate in 89% yield at 80°C after 8 h, whereas the reaction with PhCOSK/SiO₂ gave octyl thiobenzoate (**3a**) in 94% yield after 4 h. Moreover, octyl chloride gave **3a** in 52% yield after 24 h.

The present procedure using silica gel-supported potassium thioacetate and thiobenzoate allows for an efficient transformation of organic halides into thioesters. The notable advantages of this procedure are operational simplicity, good yield, and mild reaction conditions. Moreover, the products are easily isolated by filtering the used supported reagent and removing the solvent under vacuum, and are generally spectroscopically pure.