

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Nucleophilic Epoxidation of α -Sulfonyl- α,β -Unsaturated Esters with *m*-CPBA

J. L. Garc a Ruano^a; C. Fajardo^a; A. Fraile^a; M. R. Mart n^a

^a Departamento de Qu mica Org nica, Universidad Aut noma de Madrid, Madrid, Spain

To cite this Article Ruano, J. L. Garc a , Fajardo, C. , Fraile, A. and Mart n, M. R.(2005) 'Nucleophilic Epoxidation of α -Sulfonyl- α,β -Unsaturated Esters with *m*-CPBA', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 180: 5, 1489 — 1490

To link to this Article: DOI: 10.1080/10426500590913302

URL: <http://dx.doi.org/10.1080/10426500590913302>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Nucleophilic Epoxidation of α -Sulfonyl- α,β -Unsaturated Esters with *m*-CPBA

J. L. García Ruano

C. Fajardo

A. Fraile

M. R. Martín


Departamento de Química Orgánica, Universidad Autónoma de Madrid, Madrid, Spain

INTRODUCTION

m-CPBA is usually described as an electrophilic oxidant that transfers oxygen to non electron-poor alkenes. There are a few reports about the epoxidation with *m*-CPBA of deactivated olefins.^{1a–d} The reactions of 1-phenyl-(2-*p*-tolylsulfonyl)ethane^{1a,b} and 2-nitro-1-phenylpropene^{1c} using *m*-CPBA in basic aqueous media were described. We report herein the results obtained in the reaction of the title sulfones with *m*-CPBA.

RESULTS

The reactions of sulfinylfuranones **1a–c** with an excess of *m*-CPBA in CH₂Cl₂ at 0°C for 1.5–8 h afforded the corresponding sulfones **2a–c**, which were not isolated and *in situ* evolved into the sulfonyl oxiranes **3a–c** in good yields (75–89%). The reactions are stereoselective, affording only the products resulting from the *anti* approach of the reagent with respect to the OEt group. Under similar conditions, furanone **4** did not evolve into its epoxide derivative (Scheme 1).




SCHEME 1

Received July 9, 2004; accepted October 5, 2004.

Address correspondence to C. Fajardo, Departamento de Química Orgánica, Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain. E-mail: cristina.fajardo@uam.es

The above results suggest that *m*-CPBA is also able to act as a nucleophile on the proper substrates, which lead us to investigate the epoxidation of different α,β -unsaturated sulfones (**5–6**) with *m*-CPBA (Scheme 2 and Table I).



SCHEME 2

TABLE I

Eq. <i>m</i> -CPBA/eq. base	T (°C)	Time	Ratio 5:7
2.10/0	20	87 h	27:73
1.15/1.15 K₂CO₃	−20	40 min	0:100 (91)
1.15/1.15 KOH	−20	10 min	0:100 (88)

The reaction with *m*-CPBA in dichloromethane was complete to give oxirane only in the case of the furanones **2**. Cyclopentenone **5** afforded only 73% of oxirane **7** after 87 h and the open-chain compounds did not undergo epoxidation under these conditions. The addition of base to solution of *m*-CPBA allowed us to obtain the sulfonyl-oxiranes **8** in good yields; an increase in the reaction rate of **2** and **5** was also detected.

REFERENCE

- [1] a) R. Curci and F. DiFuria, *Tetrahedron Lett.*, 4085 (1974); b) Y. Apeloig, M. Karni, and Z. Rappoport, *J. Am. Chem. Soc.*, **105**, 2784 (1983); c) R. Curci, F. DiFuria, and M. Meneghin, *Gazz. Chim. Ital.*, **108**, 123 (1978); d) K. Ogura, S. Takahashi, Y. Kawamoto, M. Suzuki, M. Fujita, Y. Suzuki, and Y. Sugiyama, *Tetrahedron Lett.*, **34**, 2649–2652 (1993).