

This article was downloaded by:

On: 28 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### Studies on Pyrazine Derivatives, XLII: Synthesis and Tuberculostatic Activity of 6-(1,4-Dioxo-8-azaspiro- [4, 5]-decano)- and 6-(1-Ethoxycarbonylpiperazino)- pyrazinocarboxylic Acid Derivatives

Barbara Milczarska<sup>a</sup>; Henryk Foks<sup>a</sup>; Zofia Zwolska<sup>b</sup>

<sup>a</sup> Department of Organic Chemistry, Medical University of Gdańsk, Gdańsk, Poland <sup>b</sup> Department of Microbiology, Institute of Tuberculosis and Pulmonary Diseases, Warsaw, Poland

**To cite this Article** Milczarska, Barbara , Foks, Henryk and Zwolska, Zofia(2005) 'Studies on Pyrazine Derivatives, XLII: Synthesis and Tuberculostatic Activity of 6-(1,4-Dioxo-8-azaspiro- [4, 5]-decano)- and 6-(1-Ethoxycarbonylpiperazino)- pyrazinocarboxylic Acid Derivatives', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 180: 9, 1977 – 1992

**To link to this Article:** DOI: 10.1080/104265090902741

**URL:** <http://dx.doi.org/10.1080/104265090902741>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Studies on Pyrazine Derivatives, XLII: Synthesis and Tuberculostatic Activity of 6-(1,4-Dioxo-8-azaspiro-[4,5]-decano)- and 6-(1-Ethoxycarbonylpiperazino)-pyrazinocarboxylic Acid Derivatives

**Barbara Mileczarska**

**Henryk Foks**

Department of Organic Chemistry, Medical University of Gdańsk,  
Gdańsk, Poland

**Zofia Zwolska**

Department of Microbiology, Institute of Tuberculosis and Pulmonary  
Diseases, Warsaw, Poland

*The 2-cyano-6-chloropyrazine's chlorine atom reactivity was substituted with amines 1,4-dioxo-8-azaspiro-[4,5]-decane and 1-ethoxycarbonylpiperazine. The substrates thus obtained were used in the syntheses of the new derivatives, which were tested for their tuberculostatic activity. Minimum inhibitory concentration (MIC) value of the most active ones (5b, 12a, 14b, 16b, 21b) was 25 µg/mL.*

**Keywords** 2-Cyano-6-chloropyrazine; 6-aminosubstituted-2-cyanopyrazine; N<sup>1</sup>-substituted thioamidopyrazinocarboxyamidrazones; thioamides

## INTRODUCTION

The tuberculostatic activity of pyrazine and its derivatives has been, among other properties, a subject of intensive studies. Most of the active compounds made the pyrazino-carboxylic acid derivatives.<sup>1–5</sup> An introduction of the 1,3,4-oxathiazolino-2-one system into the pyrazine molecule increased the tuberculostatic activity 16 times in comparison with that of pyrazinamide.<sup>6</sup>


The earlier reports of Foks and colleagues<sup>7,8</sup> on some new 2-cyano-6-substituted pyrazine derivatives showed the considerable tuberculostatic activity of these compounds as well.

Received September 7, 2004; accepted September 16, 2004.

Address correspondence to Henryk Foks, Medical University of Gdańsk, Department of Organic Chemistry, Al Gen J Hallera 107, Gdańsk, PL 80-416, Poland. E-mail: hfoks@farmacja.amg.gda.pl

## RESULTS AND DISCUSSION

In continuation of our studies on the reactivity and applications of pyrazine derivatives, the new 2,6-disubstituted pyrazines have been synthesized (Scheme).<sup>1</sup>



**SCHEME 1**

The reaction of 2-cyano-6-chloropyrazine with amines 1,4-dioxaspiro-[4,5]-decane and 1-ethoxycarbonylpiperazine produced the initial pyrazinonitriles **1a** and **1b**, which were converted into thioamides (**2**), amidoximes (**3**), imidoesters (**4**), and amidrazones (**5**). The amidoximes obtained thus were allowed to react with aromatic isothiocyanates (*p*-chloro-, *p*-bromo-, and *p*-methyloisothiocyanate) the derivatives (**13–15**). The reaction course and time were controlled by means of thin layer chromatography.

In the reactions of imidoesters (**4**) with the substituted thiosemicarbazides **A–E** the N<sup>1</sup>-substituted thioamidopyrazincarboxyamidrazones (**7–12**) were formed (see Table I). The attempts at obtaining the same compounds by the earlier preparation of 3-carbonimidoyldithiocarbazoic acid S-methyl esters, followed by their reactions with the corresponding secondary amines, failed.

The condensation of amidrazones (**5a, b**) with aromatic aldehydes *p*-bromo-, *o*-, *m*-, and *p*-chloro-, *p*-nitro-, *o*-methoxy-, and 2,5-dimethoxybenzoic) produced the corresponding Schiff bases (**16–21**) (see Table II). The addition of piperidine increased the reaction yields.


TABLE I Characteristics of the Synthesized Compounds 1a, b–15a, b

| Compound no. | R <sup>2</sup> | R <sup>3</sup> | Formula molecular weight                                                 | M.p [°C] solvent for crystallization | Reaction yield [%] | IR (KBr) cm <sup>-1</sup>                                                                                   | <sup>1</sup> H NMR (500 MHz) δ [ppm] solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                                                                                       |
|--------------|----------------|----------------|--------------------------------------------------------------------------|--------------------------------------|--------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1a           | —              | —              | C <sub>12</sub> H <sub>14</sub> N <sub>4</sub> O <sub>2</sub><br>246.2   | 85–86<br>Methanol                    | 42                 | 3078 (=CH), 2062, 2232 (CN), 1579, 1514, 1100 (C—O—C)                                                       | <b>A</b> : 1.7(m, CH <sub>2</sub> CCH <sub>2</sub> ); 3.65(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.9(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 8.2, 8.4(s, 2H Py)<br><b>B</b> : 1.2(q, 3H CH <sub>3</sub> ); 3.55, 3.7(m, 8H piperazine); 4.05(q, 2H CH <sub>2</sub> ); 8.3, 8.7(s, 2H Py) |
| 1b           | —              | —              | C <sub>12</sub> H <sub>15</sub> N <sub>5</sub> O <sub>2</sub><br>261.0   | 104–106<br>Methanol                  | 40                 | 3075 (=CH), 2987, 2907, 2238 (CN), 1688 (C=O), 1588, 1520, 1251 (C—O)                                       | <b>B</b> : 1.2(q, 3H CH <sub>3</sub> ); 3.55, 3.7(m, 8H piperazine); 4.05(q, 2H CH <sub>2</sub> ); 8.3, 8.7(s, 2H Py)                                                                                                                                                                           |
| 2a           | —              | —              | C <sub>12</sub> H <sub>16</sub> N <sub>4</sub> O <sub>2</sub> S<br>280.0 | 182–183<br>Methanol                  | 80                 | 3347, 3147 (NH <sub>2</sub> ), 2998 (=CH), 2953, 2880, 1615, 1584, 1330, 1288, 1109 (C—O—C)                 | <b>A</b> : 1.8(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.02(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 7.8(s, NH); 8.4, 9.1(s, 2H Py); 9.0(s, NH)                                                                                               |
| 2b           | —              | —              | C <sub>12</sub> H <sub>17</sub> N <sub>5</sub> O <sub>2</sub> S<br>295.0 | 161–168<br>Methanol                  | 93                 | 3380, 3258, 3178 (NH <sub>2</sub> ), 3060 (=CH), 2981, 2908, 1701 (C=O), 1611, 1584, 1279 (C=S), 1225 (C—O) | <b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.67(m, 8H piperazine); 4.2(q, 2H CH <sub>2</sub> ); 8.4, 9.1(s, 2H Py); 7.9, 8.9 (s, 2H NH)                                                                                                                                                            |
| 3a           | —              | —              | C <sub>12</sub> H <sub>17</sub> N <sub>5</sub> O <sub>3</sub><br>279.0   | 194–200<br>Methanol                  | 49                 | 3493 (NOH), 3382, 3107 (NH <sub>2</sub> ), 2964, 2847, 1655, 1579, 1464, 1099 (C—O—C)                       | <b>B</b> : 1.65(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.7(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.9(m, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 5.8 (2H NH <sub>2</sub> )+ D <sub>2</sub> O decay; 8.21, 8.27(s, 2H Py); 9.95(s, 1H NOH) + D <sub>2</sub> O decay                        |
| 3b           | —              | —              | C <sub>12</sub> H <sub>18</sub> N <sub>6</sub> O <sub>3</sub><br>294.0   | 183–191<br>Methanol                  | 73                 | 3495, 3374, 3179 (NH <sub>2</sub> , NOH), 3060(=CH), 2980, 2868, 1694 (C=O), 1648, 1578, 1250 (C—O)         | <b>B</b> : 1.19(t, 3H CH <sub>3</sub> ); 3.47, 3.64(m, 8H piperazine); 4.07(q, 2H CH <sub>2</sub> ); 5.85(s, 2H NH); 8.25(d, 2H Py); 9.97(s, 1H NOH)                                                                                                                                            |

(Continued on next page)




TABLE I Characteristics of the Synthesized Compounds 1a, b–15a, b (Continued)



| Compound no. | R <sup>2</sup> | R <sup>3</sup> | Formula molecular weight                                                              | M.p [°C] solvent for crystallization     | Reaction yield [%] | IR (KBr) cm <sup>-1</sup>                                                                   | <sup>1</sup> H NMR (500 MHz) δ [ppm] solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                              |
|--------------|----------------|----------------|---------------------------------------------------------------------------------------|------------------------------------------|--------------------|---------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 4a           | —              | —              | C <sub>13</sub> H <sub>18</sub> N <sub>4</sub> O <sub>3</sub><br>278.0                | 99–105<br>Benzene                        | 86                 | 3232 (NH), 3059 (=CH), 3000, 2954, 1653, 1568, 1525, 1256, 1122 (C—O)                       | <b>B</b> : 1.8(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.0-4.03(m, 4H OCH <sub>2</sub> CH <sub>2</sub> O + 3H CH <sub>3</sub> ); 8.28, 8.3(s, 2H Py; 8.8–9.1(NH)                       |
| 4b           | —              | —              | C <sub>13</sub> H <sub>19</sub> N <sub>5</sub> O <sub>3</sub><br>293.0                | 106–110<br>Benzene                       | 59                 | 3271 (NH), 2962, 2905, 1719 (C=O), 1648, 1577, 1250, 1111 (C—O)                             | <b>B</b> : 1.15(t, 3H CH <sub>3</sub> ); 3.25(s, 3H CH <sub>3</sub> O); 3.3, 3.7(m, 8H piperazine); 4.05(q, 2H CH <sub>2</sub> ); 8.2, 8.5(s, 2H Py); 9.2(s, 1H NH)                                                                    |
| 5a           | —              | —              | C <sub>12</sub> H <sub>18</sub> N <sub>6</sub> O <sub>2</sub><br>278.0                | 84–87<br>Methanol/<br>H <sub>2</sub> O   | 64                 | 3404, 3355, 3300, 3200 (NH <sub>2</sub> ), 3050 (=CH), 2955, 2915, 1639, 1573, 1106 (C—O—C) | <b>A</b> : 1.78(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.78(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.0(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 8.07, 8.3(s, 2H Py)                                                            |
| 5b           | —              | —              | C <sub>12</sub> H <sub>19</sub> N <sub>7</sub> O <sub>2</sub><br>293.0                | 121–122<br>Methanol/<br>H <sub>2</sub> O | 92                 | 3419, 3321, 3199 (NH <sub>2</sub> ), 2986, 2917, 1698 (C=O), 1645, 1575, 1248 (C—O)         | <b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.65(m, 8H piperazine); 4.17(q, 2H CH <sub>2</sub> ); 5.0(s, 2H NH <sub>2</sub> ); 8.1, 8.6(s, 2H Py)                                                                                          |
| 6a           | —              | —              | C <sub>14</sub> H <sub>20</sub> N <sub>6</sub> O <sub>2</sub> S <sub>2</sub><br>368.0 | 153–158<br>Benzene                       | 33                 | 3392, 3145 (NH), 2954, 2918, 1670 (HN—C=S), 1617, 1580, 1107                                | <b>B</b> : 1.67(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 2.44(d, 3H CH <sub>3</sub> ); 3.75, 3.8(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.92(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 7.14, 9.2, 12.0(s, NH); 8.4, 8.6(s, 2H Py) |
| 6b           | —              | —              | C <sub>14</sub> H <sub>21</sub> N <sub>7</sub> O <sub>2</sub> S <sub>2</sub><br>383.0 | 130–131<br>Methanol/<br>H <sub>2</sub> O | 34                 | 3373, 3202 (NH), 3060 (=CH), 2983, 2920, 1669 (C=O), 1628, 1583, 1286, 1247                 | <b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 1.65(s, 3H CH <sub>3</sub> ); 3.65 (m, 8H piperazine); 4.2(q, 2H CH <sub>2</sub> ); 5.95, 8.8(s, NH); 8.2, 8.4(s, 2H Py)                                                                       |

|           |                                                                                     |                                                                            |                                          |    |                                                                                                     |                                                                                                                                                                                                                                                                 |
|-----------|-------------------------------------------------------------------------------------|----------------------------------------------------------------------------|------------------------------------------|----|-----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>7a</b> |  | — C <sub>17</sub> H <sub>25</sub> N <sub>7</sub> O <sub>2</sub> S<br>391.0 | 170–172<br>Methanol                      | 83 | 3408, 3324, 3170 (NH),<br>3060 (=CH), 2955,<br>2875, 1647, 1562,<br>1506, 1356, 1105                | <b>A:</b> 1.68(m, 4H CH <sub>2</sub> CCH <sub>2</sub> );<br>1.91(m, 4H pyrrolidine);<br>3.62(m, 4H pyrrolidine);<br>3.82 (m, 4H CH <sub>2</sub> NCH <sub>2</sub> );<br>3.92(m, 4H OCH <sub>2</sub> CH <sub>2</sub> O);<br>8.49, 8.67(s, 2H Py);<br>9.8–10.0(NH) |
| <b>7b</b> |  | — C <sub>17</sub> H <sub>26</sub> N <sub>8</sub> O <sub>2</sub> S<br>406.0 | 166–170<br>Methanol/<br>H <sub>2</sub> O | 54 | 3434, 3299, 3156 (NH),<br>3070 (=CH), 2960,<br>2860, 1697 (C=O),<br>1666, 1600, 1573,<br>1286, 1249 | <b>A:</b> 1.29(t, 3H CH <sub>3</sub> ); 1.99(m<br>4H pyrrolidine); 3.67(m,<br>8H pyrrolidine +<br>piperazine); 3.82(m, 4H<br>CH <sub>2</sub> NCH <sub>2</sub> ); 4.19(q, 2H<br>CH <sub>2</sub> ); 6.3(NH); 8.2, 8.5(s,<br>2H Py)                                |
| <b>8a</b> |  | — C <sub>17</sub> H <sub>25</sub> N <sub>7</sub> O <sub>3</sub> S<br>407.0 | 160–166<br>Acetone                       | 34 | 3418, 3257, 3150 (NH),<br>3088 (=CH), 2950,<br>2892, 1668, 1601,<br>1575, 1368, 1343,<br>1247, 1101 | <b>A:</b> 1.8(m, 4H CH <sub>2</sub> CCH <sub>2</sub> );<br>3.84(m, 4H CH <sub>2</sub> NCH <sub>2</sub> );<br>3.74, 4.0(m, 8H<br>morpholine + 4H<br>OCH <sub>2</sub> CH <sub>2</sub> O); 6.9–<br>7.0(NH); 8.3, 8.42(s, 2H<br>Py)                                 |

(Continued on next page)




TABLE I Characteristics of the Synthesized Compounds 1a, b–15a, b (Continued)


| Compound no. | R <sup>2</sup>                                                                      | R <sup>3</sup> | Formula molecular weight                                                 | M.p [°C] solvent for crystallization | Reaction yield [%] | IR (KBr) cm <sup>-1</sup>                                                               | <sup>1</sup> H NMR (500 MHz) δ [ppm] solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                                                                                     |
|--------------|-------------------------------------------------------------------------------------|----------------|--------------------------------------------------------------------------|--------------------------------------|--------------------|-----------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 8b           |  | —              | C <sub>17</sub> H <sub>26</sub> N <sub>8</sub> O <sub>3</sub> S<br>422.0 | 170–175 Methanol/<br>Acetone         | 69                 | 3461, 3348 (NH), 3050 (=CH), 2958, 2897, 1696 (C=O), 1666, 1568, 1526, 1248, 1116       | <b>B</b> : 1.19(t, 3H CH <sub>3</sub> ); 3.48(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.56, 3.81(t, 8H morpholine); 3.75(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.06(q, 2H CH <sub>2</sub> ); 8.51, 8.57(s, 2H Py); 12.66(NHC=S)                                                         |
| 9a           |  | —              | C <sub>20</sub> H <sub>29</sub> N <sub>7</sub> O <sub>4</sub> S<br>463.0 | 175–178 Ethyl acetate/<br>Acetone    | 59                 | 3362, 3291 (NH), 2951, 2884, 1677, 1570, 1526, 1312, 1260, 1131                         | <b>B</b> : 1.54(m, 4H CH <sub>2</sub> CCH <sub>2</sub> -a); 1.67(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> -a); 3.9 (m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.95(m, 8H OCH <sub>2</sub> CH <sub>2</sub> O); 8.47, 8.6(s, 2H Py); 12.6(1H NHC=S) |
| 9b           |  | —              | C <sub>20</sub> H <sub>30</sub> N <sub>8</sub> O <sub>4</sub> S<br>478.0 | 180–182 Acetone                      | 77                 | 3379, 3300, 3129 (NH), 3040 (=CH), 2956, 2892, 1698 (C=O), 1585, 1531, 1360, 1232, 1131 | <b>B</b> : 1.19(t, 3H CH <sub>3</sub> ); 1.55(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.48(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.75, 3.93(m, 8H piperazine); 3.9(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 4.05(q, 2H CH <sub>2</sub> ); 8.51, 8.56(s, 2H Py); 12.7 (NHC=S)          |

|                                                                                                   |                                                                            |                     |    |                                                                                                     |                                                                                                                                                                                                                                                                                                                                                                                    |
|---------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|---------------------|----|-----------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>10a</b><br> | — C <sub>23</sub> H <sub>30</sub> N <sub>9</sub> O <sub>2</sub> S<br>482.0 | 165–168<br>Methanol | 31 | 3427, 3284, 3143 (NH),<br>3059 (=CH), 2952,<br>2885, 1670, 1598,<br>1224                            | <b>B</b> : 1.67(m, 4H CH <sub>2</sub> CCH <sub>2</sub> );<br>3.1(m, 4H piperazine);<br>3.82(m, 4H piperazine);<br>3.82(m, 4H piperidine);<br>3.92(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O);<br>3.98(m, 4H piperazine);<br>6.78, 6.97, 7.2(m, 5H Ph);<br>7.7(s, NH <sub>2</sub> + D <sub>2</sub> O-decay);<br>8.49, 8.6(s, 2H Py);<br>12.67(s, NH—C=S +<br>D <sub>2</sub> O-decay) |
| <b>10b</b><br> | — C <sub>23</sub> H <sub>31</sub> N <sub>9</sub> O <sub>2</sub> S<br>497.0 | 173–175<br>Methanol | 29 | 3448, 3345, 3166 (NH),<br>3065 (=CH), 2976,<br>2921, 1673 (C=O),<br>1599, 1577, 1529,<br>1347, 1227 | <b>B</b> : 1.1(t, 3H CH <sub>3</sub> ); 3.1(m,<br>8H piperazine- <b>a</b> );<br>3.49(m, 8H piperazine);<br>4.06(q, 2H CH <sub>2</sub> ); 6.78,<br>6.98, 7.2(m, 5H Ph); 8.52,<br>8.57(s, 2H Py);<br>12.7(NH—C=S)                                                                                                                                                                    |

(Continued on next page)

TABLE I Characteristics of the Synthesized Compounds 1a, b–15a, b (Continued)

| Compound no. | R <sup>2</sup>                                                                      | R <sup>3</sup> | Formula molecular weight                                                 | M.p [°C] solvent for crystallization | Reaction yield [%] | IR (KBr) cm <sup>-1</sup>                                                   | <sup>1</sup> H NMR (500 MHz) δ [ppm] solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                                                                                                                   |
|--------------|-------------------------------------------------------------------------------------|----------------|--------------------------------------------------------------------------|--------------------------------------|--------------------|-----------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>11a</b>   |  | —              | C <sub>24</sub> H <sub>32</sub> N <sub>8</sub> O <sub>2</sub> S<br>496.0 | 157–159<br>Benzene                   | 55                 | 3290, 3143 (NH), 3030 (=CH), 2928, 2887, 1683, 1578, 1421, 1303, 1243, 1108 | <b>B</b> : 1.67(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 2.33(m, 4H piperazine); 3.47(s, 2H CH <sub>2</sub> ); 3.8–3.83(m, 8H CH <sub>2</sub> NCH <sub>2</sub> piperidine and piperazine); 3.92 (s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 7.3–7.35(m, 5H Ph); 8.47, 8.6(s, 2H Py); 12.7(NH—C≡S + D <sub>2</sub> O-decay) |
| <b>11b</b>   |  | —              | C <sub>24</sub> H <sub>32</sub> N <sub>8</sub> O <sub>2</sub> S<br>511.0 | 132–138<br>Benzene                   | 43                 | 3419, 3290 (NH), 3062 (=CH), 2923, 1700 (C=O), 1635, 1600, 1578, 1233, 1115 | <b>A</b> : 1.19(t, 3H CH <sub>3</sub> ); 2.33(s, 2H CH <sub>2</sub> ); 3.47 (m, 8H piperazine- <b>a</b> ); 3.74, 3.84(m, 8H piperazine); 4.07(q, 2H CH <sub>2</sub> ); 7.25, 7.31(m, 5H Ph); 8.5, 8.56(s, 2H Py); 12.7(NH—C≡S)                                                                                              |
| <b>12a</b>   |  | —              | C <sub>22</sub> H <sub>27</sub> N <sub>7</sub> O <sub>2</sub> S<br>453.0 | 154–158<br>Benzene                   | 55                 | 3302, 3155 (NH), 2955, 2884, 1685, 1579, 1530, 1350, 1239, 1108, 750        | <b>B</b> : 1.67(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 2.79 and 4.11(t, 4H CH <sub>2</sub> CH <sub>2</sub> ); 3.81(s, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.92 (s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 5.0(s, 2H CH <sub>2</sub> ); 7.14 (m, 4H Ar); 8.48, 8.60(s, 2H Py); 12.7(s, 1H NH—C≡S)                     |

|            |                                                                                     |      |                                   |                                   |    |                                                                                                     |                                                                                                                                                                                                                                                     |
|------------|-------------------------------------------------------------------------------------|------|-----------------------------------|-----------------------------------|----|-----------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>12b</b> |  | —    | $C_{22}H_{28}N_8O_2S$<br>468.0    | 140–146<br>Benzene                | 40 | 3350, 3084 (NH), 2926,<br>2859, 1692 (C=O),<br>1601, 1573, 1353,<br>1280, 1231 (C-O)                | <b>B:</b> 1.19(t, 3H CH <sub>3</sub> ); 2.79<br>and 4.11(t, 4H CH <sub>2</sub> CH <sub>2</sub> );<br>3.48 and 3.75(s, 8H<br>piperazine); 4.07(q, 2H<br>CH <sub>2</sub> ); 7.15(m, 4H Ar);<br>8.5, 8.57 (s, 2H Py);<br>12.7(s, 1H NH—C=S)            |
| <b>13a</b> | —                                                                                   | p-Cl | $C_{19}H_{21}N_6O_3SCl$<br>448.45 | 204–207.5<br>Acetone/<br>Methanol | 74 | 3377, 3257 (NH), 3050<br>(=CH), 2952, 2920,<br>1641, 1614, 1526,<br>1307, 1105                      | <b>A:</b> 1.82(m, 4H CH <sub>2</sub> CCH <sub>2</sub> );<br>3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> );<br>4.02(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O);<br>7.3, 7.6(m, 4H Ar); 8.55,<br>8.9(s, 2H Py); 9.6(s, 1H<br>NOH)                       |
| <b>13b</b> | —                                                                                   | p-Cl | $C_{19}H_{22}N_7O_3SCl$<br>463.45 | 207–210<br>Methanol/<br>Acetone   | 70 | 3390, 3250 (NH), 3050<br>(=CH), 2985, 2920,<br>1689 (C=O), 1650,<br>1575, 1305, 1250                | <b>A:</b> 1.3(t, 3H CH <sub>3</sub> ); 3.67(m,<br>8H piperazine); 4.2(q, 2H<br>CH <sub>2</sub> ); 7.3, 7.55(m, 4H<br>Ar); 8.5 (m, 2H Py); 8.8(s,<br>NOH)                                                                                            |
| <b>14a</b> | —                                                                                   | p-Br | $C_{19}H_{21}N_6O_3SBr$<br>492.9  | 207–211<br>Acetone                | 52 | 3375, 3251 (NH), 3050<br>(=CH), 2952, 2919,<br>1640, 1613, 1574,<br>1307, 1250, 1102                | <b>B:</b> 1.68(m, 4H CH <sub>2</sub> CCH <sub>2</sub> );<br>3.79(m, 4H CH <sub>2</sub> NCH <sub>2</sub> );<br>3.81(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O);<br>7.4, 7.6(m, 4H Ar); 8.5,<br>8.6(s, 2H Py); 8.58, 9.3 (s,<br>2H NH); 9.7(s, 1H NOH) |
| <b>14b</b> | —                                                                                   | p-Br | $C_{19}H_{22}N_7O_3SBr$<br>508.0  | 193–194<br>Methanol/<br>acetone   | 16 | 3389, 3267 (NH), 3055<br>(=CH), 2980, 2924,<br>1674 (C=O), 1654,<br>1577, 1515, 1437,<br>1248, 1133 | <b>A:</b> 1.2(t, 3H CH <sub>3</sub> ); 3.49,<br>3.72(m, 8H piperazine);<br>4.07(q, 2H CH <sub>2</sub> ); 7.44,<br>7.64(m, 4H Ar); 8.5,<br>8.7(s, 2H Py); 8.6, 9.3(s,<br>2H NH); 9.69(s, NOH)                                                        |

(Continued on next page)

TABLE I Characteristics of the Synthesized Compounds 1a, b–15a, b (Continued)

| Compound no. | R <sup>2</sup> | R <sup>3</sup>    | Formula molecular weight                                                 | M.p [°C] solvent for crystallization | Reaction yield [%] | IR (KBr) cm <sup>-1</sup>                                                         | <sup>1</sup> H NMR (500 MHz) δ [ppm] solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                                                                                                                                                                                                                                                                                                |
|--------------|----------------|-------------------|--------------------------------------------------------------------------|--------------------------------------|--------------------|-----------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>15a</b>   | —              | p-CH <sub>3</sub> | C <sub>20</sub> H <sub>24</sub> N <sub>6</sub> O <sub>3</sub> S<br>428.0 | 202–205<br>Acetone/<br>Methanol      | 59                 | 3367, 3251 (NH), 3050 (=CH), 2952, 2920, 1640, 1614, 1574, 1525, 1340, 1257, 1107 | <b>B</b> : 1.68(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 2.2(s, 3H CH <sub>3</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.92(s, 4H OCH <sub>2</sub> CH <sub>2</sub> O); 7.05, 7.5(m, 4H Ar); 8.45, 9.3(s, 2H NH); 8.5, 8.65(s, 2H Py); 9.45(s, NOH)<br><b>B</b> : 1.19(t, 3H CH <sub>3</sub> ); 2.2(s, 3H CH <sub>3</sub> ); 3.5(m, 4H piperazine); 3.7(m, 4H piperazine); 4.06(q, 2H CH <sub>2</sub> ); 7.05, 7.55(m, 4H Ar); 8.5, 8.7(m, 2H Py); 9.45(s, 1H NOH + D <sub>2</sub> O-decay) |
| <b>15b</b>   | —              | p-CH <sub>3</sub> | C <sub>20</sub> H <sub>25</sub> N <sub>7</sub> O <sub>3</sub> S<br>443.0 | 198–200<br>Methanol/<br>Acetone      | 55                 | 3295, 2984, 2918, 1697 (C=O), 1635, 1576, 1434, 1402, 1249                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |

Ar—aromatic, Ph—phenyl, Py—pyrazine.

TABLE II Physicochemical Data of Condensation Products of Amidrazones with Aromatic Aldehydes

| Compound no. | R <sup>1</sup> | Formula molecular weight                                                  | M.p [°C] solvent for crystallization    | Yield [%] | IR (KBr) cm <sup>-1</sup>                                                                                                | <sup>1</sup> H NMR δ [ppm] 500 MHz solvent: <b>A</b> —CDCl <sub>3</sub> ; <b>B</b> —DMSO-d <sub>6</sub>                                                                                                                                                                                                                                                                                                |
|--------------|----------------|---------------------------------------------------------------------------|-----------------------------------------|-----------|--------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>16a</b>   | p-Cl           | C <sub>19</sub> H <sub>21</sub> N <sub>6</sub> O <sub>2</sub> Cl<br>400.4 | 175–177<br>Methanol                     | 65        | 3454, 3334 (NH <sub>2</sub> ), 3050 (=CH), 2984, 2941, 2880, 1621, 1602, 1560, 1520, 1108 (C—O—C)                        | <b>B</b> : 1.68(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.78(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.9 (4H CH <sub>2</sub> CH <sub>2</sub> ); 7.0–7.2(s, 2H NH); 7.5–7.9(m, 4H Ar); 8.4, 8.46(s, 2H Py); 8.54(s, 1H CH)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.5–3.75(m, 8H piperazine); 4.2 (q, 2H CH <sub>2</sub> ); 6.4(s, NH); 7.4, 7.75(d, 4H Ar); 8.3, 8.6 (s, 2H Py); 8.9(s, 1H CH) |
| <b>16b</b>   | p-Cl           | C <sub>19</sub> H <sub>22</sub> N <sub>7</sub> O <sub>2</sub> Cl<br>415.4 | 209–210<br>Methanol                     | 26        | 3493, 3378 (NH <sub>2</sub> ), 3061 (=CH), 2983, 2960, 2853, 1697 (C=O), 1626, 1603, 1560, 1437, 1250 (C—O)              | <b>B</b> : 1.68(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.78(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.93(4H CH <sub>2</sub> CH <sub>2</sub> ); 7.0–7.2(s, 2H NH); 7.5–7.9(m, 4H Ar); 8.4, 8.46(s, 2H Py); 8.54(s, 1H CH)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.65(m, 8H piperazine); 4.19(q, 2H CH <sub>2</sub> ); 6.4(s, NH); 7.4–7.8(m, 4H Ar); 8.2, 8.5(s, 2H Py)                       |
| <b>17a</b>   | o-Cl           | C <sub>19</sub> H <sub>21</sub> N <sub>6</sub> O <sub>2</sub> Cl<br>400.4 | 160–163<br>Acetone/<br>H <sub>2</sub> O | 96        | 3480, 3290 (NH <sub>2</sub> ), 3067 (=CH), 2954, 2887, 1616, 1568, 1105 (C—O—C)                                          | <b>B</b> : 1.68(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.78(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.93(4H CH <sub>2</sub> CH <sub>2</sub> ); 7.0–7.2(s, 2H NH); 7.5–7.9(m, 4H Ar); 8.4, 8.46(s, 2H Py); 8.54(s, 1H CH)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.65(m, 8H piperazine); 4.19(q, 2H CH <sub>2</sub> ); 6.4(s, NH); 7.4–7.8(m, 4H Ar); 8.2, 8.5(s, 2H Py)                       |
| <b>17b</b>   | o-Cl           | C <sub>19</sub> H <sub>22</sub> N <sub>7</sub> O <sub>2</sub> Cl<br>415.4 | 173–176<br>Benzene                      | 19        | 3489, 3373 (NH <sub>2</sub> ), 3062 (=CH), 2983, 2961, 2855, 1696 (C=O), 1625, 1603, 1560, 1437, 1250 (C—O)              | <b>B</b> : 1.68(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.78(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.93(4H CH <sub>2</sub> CH <sub>2</sub> ); 7.0–7.2(s, 2H NH); 7.5–7.9(m, 4H Ar); 8.4, 8.46(s, 2H Py); 8.54(s, 1H CH)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.65(m, 8H piperazine); 4.19(q, 2H CH <sub>2</sub> ); 6.4(s, NH); 7.4–7.8(m, 4H Ar); 8.2, 8.5(s, 2H Py)                       |
| <b>18a</b>   | p-Br           | C <sub>19</sub> H <sub>21</sub> N <sub>6</sub> O <sub>2</sub> Br<br>445.0 | 188–189<br>Dioxane                      | 80        | 3454, 3336 (NH <sub>2</sub> ), 3050 (=CH), 2942, 1620, 1559, 1525, 1465, 1107 (C—O—C)                                    | <b>B</b> : 1.8(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.03(s, 4H CH <sub>2</sub> CH <sub>2</sub> ); 6.3(s, NH); 7.5–7.7(m, 4H Ar); 8.3(s, 1H CH); 8.5, 8.8(s, 2H Py)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.67(m, 8H piperazine); 4.19(q, 2H CH <sub>2</sub> ); 6.4(s, 2H NH); 7.96, 8.27(d, 4H Ar); 8.3, 8.6(s, 2H Py); 8.9(s, 1H CH)          |
| <b>18b</b>   | p-Br           | C <sub>19</sub> H <sub>22</sub> N <sub>7</sub> O <sub>2</sub> Br<br>460.0 | 256–258<br>Dioxane                      | 59        | 3493, 3372 (NH <sub>2</sub> ), 3061 (=CH), 2975, 2961 (CH <sub>2</sub> ), 1697 (C=O), 1625, 1605, 1560, 1439, 1251 (C—O) | <b>B</b> : 1.8(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 4.03(s, 4H CH <sub>2</sub> CH <sub>2</sub> ); 6.3(s, NH); 7.5–7.7(m, 4H Ar); 8.3(s, 1H CH); 8.5, 8.8(s, 2H Py)<br><b>A</b> : 1.3(t, 3H CH <sub>3</sub> ); 3.67(m, 8H piperazine); 4.19(q, 2H CH <sub>2</sub> ); 6.4(s, 2H NH); 7.96, 8.27(d, 4H Ar); 8.3, 8.6(s, 2H Py); 8.9(s, 1H CH)          |

(Continued on next page)

TABLE II Physicochemical Data of Condensation Products of Amidrazones with Aromatic Aldehydes  
(Continued)

| Compound no. | R <sup>1</sup>                       | Formula molecular weight                                               | M.p [°C]<br>solvent for crystallization | Yield [%] | IR (KBr) cm <sup>-1</sup>                                                                                                      | <sup>1</sup> H NMR δ [ppm] 500 MHz<br>solvent: A—CDCl <sub>3</sub> ; B—DMSO-d <sub>6</sub>                                                                                                                                                                                         |
|--------------|--------------------------------------|------------------------------------------------------------------------|-----------------------------------------|-----------|--------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 19a          | p-NO <sub>2</sub>                    | C <sub>19</sub> H <sub>21</sub> N <sub>7</sub> O <sub>4</sub><br>411.0 | 222–226<br>Acetone/<br>Methanol         | 92        | 3458, 3335 (NH <sub>2</sub> ), 3050 (=CH), 2961, 2933, 2878, 1630, 1608, 1594, 1473, 1366 (NO <sub>2</sub> ), 1107 (C—O—C)     | B: 1.7(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.8(t, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.92 (s, 4H CH <sub>2</sub> CH <sub>2</sub> ); 7.2–7.4(s, NH <sub>2</sub> ); 8.2, 8.26(m, 4H Ar); 8.45(s, 1H CH); 8.55, 8.58(s, 2H Py)                                              |
| 19b          | p-NO <sub>2</sub>                    | C <sub>19</sub> H <sub>22</sub> N <sub>8</sub> O <sub>4</sub><br>426.0 | 245–251<br>Dioxane                      | 70        | 3505, 3391 (NH <sub>2</sub> ), 2890, 2913, 2857, 1696 (C=O), 1625, 1608, 1592, 1527, 1433, 1335 (NO <sub>2</sub> ); 1250 (C—O) | A: 1.28(t, 3H CH <sub>3</sub> ); 3.65(m, 8H piperazine); 4.18(q, 2H CH <sub>2</sub> ); 6.4(s, 2H NH); 7.9, 8.25(s, 4H Ar); 8.26, 8.6(s, 2H Py); 8.9(s, 1H CH)                                                                                                                      |
| 20a          | o-CH <sub>3</sub> O                  | C <sub>20</sub> H <sub>24</sub> N <sub>6</sub> O <sub>3</sub><br>396.0 | 187–192<br>Acetone/<br>H <sub>2</sub> O | 52        | 3501, 3386 (NH <sub>2</sub> ), 3067 (=CH), 2957, 2889, 1622, 1563, 1254, 1112                                                  | B: 1.68(t, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.77(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.85(s, 3H CH <sub>3</sub> O); 3.93(s, 4H CH <sub>2</sub> CH <sub>2</sub> ); 6.98(s, NH <sub>2</sub> ); 7–7.1, 7.4–8.1(m, 4H Ar); 8.4(s, 1H CH); 8.55, 8.7(s, 2H Py)              |
| 20b          | o-CH <sub>3</sub> O                  | C <sub>20</sub> H <sub>25</sub> N <sub>7</sub> O <sub>3</sub><br>411.0 | 185–187<br>Methanol                     | 73        | 3451, 3249 (NH <sub>2</sub> ), 3051 (=CH), 2984, 2908, 1695 (C=O), 1607, 1566, 1279 (C—O)                                      | B: 1.2(t, 3H CH <sub>3</sub> ); 3.5–3.7(m, 8H piperazine); 3.9(s, 3H CH <sub>3</sub> O); 4.07(q, 2H CH <sub>2</sub> ); 7.0(s, NH); 7.0–7.1, 7.4–8.2(m, 4H Ar); 8.38(s, 1H CH); 8.6, 8.7(s, 2H Py)                                                                                  |
| 21a          | 2,5-(CH <sub>3</sub> O) <sub>2</sub> | C <sub>21</sub> H <sub>27</sub> N <sub>6</sub> O <sub>4</sub><br>427.0 | 180–181<br>Acetone                      | 74        | 3469, 3268 (NH <sub>2</sub> ), 3072 (=CH), 2955, 2870, 1611, 1565, 1521, 1258, 1193, 1101                                      | B: 1.68(m, 4H CH <sub>2</sub> CCH <sub>2</sub> ); 3.77(s, 6H (CH <sub>3</sub> O) <sub>2</sub> ); 3.8(m, 4H CH <sub>2</sub> NCH <sub>2</sub> ); 3.93(s, 4H CH <sub>2</sub> CH <sub>2</sub> ); 7.0, 7.7(m, 3H Ar); 6.9–7.1(s, NH <sub>2</sub> ); 8.4(s, 1H CH); 8.55, 8.67(s, 2H Py) |
| 21b          | 2,5-(CH <sub>3</sub> O) <sub>2</sub> | C <sub>21</sub> H <sub>27</sub> N <sub>7</sub> O <sub>4</sub><br>441.0 | 112–117<br>Benzene                      | 43        | 3469, 3261 (NH <sub>2</sub> ), 3052 (=CH), 2990, 2937, 1697 (C=O), 1667, 1613, 1587, 1245 (C—O)                                | A: 1.3(t, 3H CH <sub>3</sub> ); 3.66, 3.85(m, 8H piperazine + 6H CH <sub>3</sub> O); 4.2(q, 2H CH <sub>2</sub> ); 6.2(1H NH); 7.0–7.4(m, 3H Ar); 8.25, 8.45(s, 2H Py); 9.0(s, 1H CH)                                                                                               |

Ar—aromatic, Py—pyrazine.

The structure of the newly obtained compounds was established by the analyses of IR and  $^1\text{H}$  NMR spectra. The spectral data are given in the Tables I and II.

## MICROBIOLOGY

The newly obtained derivatives were tested for their tuberculostatic activity towards the standard *Mycobacterium tuberculosis* H<sub>37</sub>Rv strain, as well as two strains isolated from tuberculous patients (Table III). One strain, *Myc. spec.* 210, was resistant to *P*-Aminosalicylic Acid (PAS), Isonicotinic Acid Hydrazide (INH), Ethambutol (EMB) and Rifampycine (RFP); the other strain, *Myc. spec.* 192, was fully susceptible to the drugs administered.

Tuberculostatic activity was tested in vitro with the classical test tube method on Youman's liquid medium containing 10% of a bovine serum.<sup>9</sup> The Minimum Growth Inhibiting Concentration (MIC) values obtained showed that some compounds were noteworthy. MIC values of the compounds **5b**, **12a**, **14b**, **16b**, and **21b** were within the limits of 25–50  $\mu\text{g/mL}$ . The MIC values obtained for pyrazinamide were within 30–60  $\mu\text{g/mL}$  using the same method.

TABLE III Tuberculostatic Activity [ $\mu\text{g/mL}$ ]

| Compound no. | Myc. tbc. H <sub>37</sub> Rv | Myc. spec. 192 susceptible | Myc. spec. 210 resistant | Compound no. | Myc. tbc. H <sub>37</sub> Rv | Myc. spec. 192 susceptible | Myc. spec. 210 resistant |
|--------------|------------------------------|----------------------------|--------------------------|--------------|------------------------------|----------------------------|--------------------------|
| <b>1a</b>    | 50                           | 100                        | 50                       | <b>12a</b>   | 25                           | 50                         | 25                       |
| <b>1b</b>    | 50                           | 100                        | 50                       | <b>12b</b>   | 50                           | 50                         | 50                       |
| <b>2a</b>    | 100                          | 50                         | 50                       | <b>13a</b>   | 50                           | 100                        | 100                      |
| <b>2b</b>    | 50                           | 50                         | 50                       | <b>13b</b>   | 50                           | 50                         | 50                       |
| <b>3a</b>    | 50                           | 100                        | 50                       | <b>14a</b>   | 100                          | 100                        | 100                      |
| <b>3b</b>    | 50                           | 50                         | 50                       | <b>14b</b>   | 50                           | 25                         | 25                       |
| <b>5a</b>    | 50                           | 100                        | 50                       | <b>15a</b>   | 100                          | 50                         | 50                       |
| <b>5b</b>    | 50                           | 50                         | 25                       | <b>15b</b>   | 50                           | 50                         | 50                       |
| <b>6a</b>    | 50                           | 50                         | 50                       | <b>16a</b>   | 50                           | 50                         | 50                       |
| <b>6b</b>    | 50                           | 50                         | 50                       | <b>16b</b>   | 50                           | 50                         | 25                       |
| <b>7a</b>    | 50                           | 50                         | 50                       | <b>18a</b>   | 100                          | 100                        | 100                      |
| <b>7b</b>    | 50                           | 50                         | 50                       | <b>18b</b>   | 50                           | 50                         | 50                       |
| <b>8a</b>    | 50                           | 100                        | 50                       | <b>19a</b>   | 50                           | 100                        | 50                       |
| <b>8b</b>    | 50                           | 50                         | 50                       | <b>19b</b>   | 50                           | 50                         | 50                       |
| <b>9b</b>    | 50                           | 50                         | 50                       | <b>20a</b>   | 50                           | 50                         | 50                       |
| <b>10a</b>   | 50                           | 50                         | 100                      | <b>20b</b>   | 50                           | 50                         | 50                       |
| <b>10b</b>   | 50                           | 50                         | 50                       | <b>21a</b>   | 50                           | 50                         | 50                       |
| <b>11a</b>   | 50                           | 50                         | 50                       | <b>21b</b>   | 50                           | 50                         | 25                       |
| <b>11b</b>   | 50                           | 50                         | 50                       |              |                              |                            |                          |

## EXPERIMENTAL

All melting points were obtained with a Boëtius apparatus and are uncorrected. The elemental analysis results for C and H of all the compounds obtained were in good agreement with the data calculated.

The IR spectra were taken with a Satellite spectrophotometer, and the  $^1\text{H}$  NMR spectra were taken with a Varian Unity 500 MHz apparatus. The reaction yields and physical constants of the new compounds are given in the Tables I and II.

### Syntheses of Substituted 2-Cyanopyrazine Derivatives (1a, 1b)

To a solution of 2-cyano-6-chloropyrazine (0.1 mol) in benzene (150 cm<sup>3</sup>), triethylamine (0.15 mol) and corresponding amine (0.1 mol), i.e., 1,4-dioxo-8-azaspiro-[4,5]-decane or 1-ethoxycarbonylpiperazine, was added. The mixture was refluxed for 2 h. On cooling down, water (60 cm<sup>3</sup>) was added and the mixture was extracted with benzene. The benzene extracts were dried over anhydrous MgSO<sub>4</sub>. The solution obtained was thickened by evaporation under vacuum and allowed to stand for crystallization. The precipitate was filtered off and recrystallized.

### Syntheses of Thioamides (2a, 2b)

The corresponding nitrile (2 mmol) was dissolved in ethanol (8 cm<sup>3</sup>) and ammonium polysulfide was added until turbidity appeared. After 12 h, the precipitate was filtered off and crystallized.

### Syntheses of Amidoximes (3a, 3b)

To a solution of hydroxylamine hydrochloride (17.0 mmol) in absolute methanol (10 cm<sup>3</sup>) the solution of KOH (22 mmol) in absolute methanol (10 cm<sup>3</sup>) was added. The precipitated KCl was filtered off and the filtrate was treated with the corresponding nitrile **1a** or **1b** (2.5 mmol). The whole was refluxed for 2 h. The solvent was then evaporated under reduced pressure and the residue treated with acetic acid (0.8 cm<sup>3</sup>) in water (10 cm<sup>3</sup>). The precipitate was filtered off and recrystallized.

### Syntheses of Imidoesters (4a, 4b)

To the solution of nitrile **1a** or **1b** (4 mmol) in benzene (60 cm<sup>3</sup>) the solution of NaOH (10 g) in water (10 cm<sup>3</sup>) was added, followed by methanol (10 cm<sup>3</sup>) and a small amount of triethylbenzylamine (TEBA). The whole

was stirred for 2 h at ambient temperature. The water (70 cm<sup>3</sup>) was then added and the aqueous layer after separation was extracted with benzene. The combined benzene extracts were dried over anhydrous MgSO<sub>4</sub>. The benzene solution was thickened under reduced pressure and the oily residue was treated with petroleum ether. The precipitates were filtered off and recrystallized.

### Syntheses of Amidrazones (5a, 5b)

To the solution of nitrile **1a** or **1b** (2.5 mmol) in methanol (8 cm<sup>3</sup>), 80%—hydrazine (1.3 cm<sup>3</sup>) was added. The solution was heated to 50°C for 1 h. The solvent and an excess of hydrazine were evaporated under reduced pressure. The residue was cooled and treated with a small amount of water. The precipitate obtained was filtered off and crystallized.

### The Reactions of Amidoximes with Aryloisothiocyanates (13a, b–15a, b)

Amidoxime **3a** or **3b** (0.5 mmol) and corresponding isothiocyanate (0.5 mmol) were suspended in DMF (5 cm<sup>3</sup>) and stirred at ambient temperature for 3–5 days. The mixture was ice-cooled and treated with water. The precipitate was filtered off and crystallized.

### Condensation of Amidrazones with Aromatic Aldehydes (16a, b–21a, b)

To the solution of the corresponding amidrazone (0.64 mmol) in absolute methanol (8 cm<sup>3</sup>) aromatic aldehyde (0.64 mmol) and a few drops of piperidine were added. The whole was refluxed for 4 h. On cooling down, the precipitates obtained were filtered off and crystallized.

### Syntheses of 3-Carbonimidoyldithiocarbazonic Acid S-Methyl Esters (6a, 6b)

Imidoester **4a** or **4b** (1 mmol) in absolute methanol (8 cm<sup>3</sup>) was treated with methyl dithiocarbazones (1 mmol) dissolved in methanol (3 cm<sup>3</sup>). The whole was stirred at ambient temperature for 24 h. The precipitate was filtered off and crystallized.

### Syntheses of Thiosemicarbazides A–E

A suspension of methyl dithiocarbazonate (2 mmol) in water (5 cm<sup>3</sup>) was treated with the corresponding amine (6 mmol), i.e., pyrrolidine

(**A**), morpholine (**B**), phenylpiperazine (**C**), 1,4-dioxo-8-azaspiro-[4,5]-decane (**D**), or 1-benzylpiperazine (**E**), dissolved in water (3 cm<sup>3</sup>). The whole was heated with a water bath for 6 h. On cooling down, the precipitate was filtered off and washed with water.

Thiosemicarbazide: **A** M.p. 165–9°C (45% yield); **B** M.p. 186–6°C (23%); **C** M.p. 170–3°C (42%); **D** M.p. 130–6°C (15%); **E** M.p. 161–5°C (34%).

### Syntheses of N<sup>1</sup>-Substituted Thioamidopyrazincarboxyamidrazones (7a, b–12a, b)

A solution of imidoester **4a** or **4b** (1 mmol) and the corresponding thiosemicarbazide **A–E** (1 mmol) in methanol (20 mcm<sup>3</sup>) was stirred at ambient temperature for 4 days. The solvent excess was distilled off. On cooling down, the oily residue was treated with benzene and petroleum ether. The precipitate obtained was filtered off and crystallized.

### REFERENCES

- [1] M. Dolezal, M. Miletin, J. Kunes, and K. Kriukova, *Molecules*, **7**, 363 (2002).
- [2] J. Hartl, M. Dolezal, J. Krinkova, A. Lycka, and Z. Odlerova, *Collect. Czech. Chem. Commun.*, **61**, 1109 (1996).
- [3] M. Ch. Davis, *Diss. Abstr. Int. B*, 58, 4230 (1998); C.A. **128**, 57096b (1998).
- [4] J. Krinkova, V. Buchta, M. Dolezal, P. Milan, and J. Hartl, *Il Farmaco*, **57**, 71 (2002); C.A. **137**, 152209k (2002).
- [5] M. Cynamon, R. Gimi, F. Gyenes, C. Sharpe, K. Bergmann, J. Han Hye, L. B. Gregor, R. Rapuolu, G. Luciano, J. T. Welch, *J. Med. Chem.*, **38**, 3902 (1995); C.A. **123**, 285904r (1995).
- [6] M. Gezginci, A. Martin, and S. Franzblau, *J. Med. Chem.*, **44**, 1560 (2001).
- [7] H. Foks, W. Manowska, and J. Sienicka, *Acta Polon. Pharm.*, **6**, 687 (1976).
- [8] H. Foks, M. Janowiec, and B. Pilarski, *Pol. J. Pharmacol. Pharm.*, **30**, 105 (1978).
- [9] D. Pancechowska-Ksepko, H. Foks, M. Janowiec, and Z. Zwolska-Kwiek, *Acta Polon. Pharm.*, **49**, 193 (1988).