

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

Applications of the $(\text{PhO})_3\text{PCl}_2$ Reagent: A New Protocol for Mild Cleavage of Sulfinamides and Sulfonamides

Alberto Spaggiari^a; Fabio Prati^a

^a Dipartimento di Chimica, Università di Modena e Reggio Emilia, Modena, Italy

To cite this Article Spaggiari, Alberto and Prati, Fabio(2006) 'Applications of the $(\text{PhO})_3\text{PCl}_2$ Reagent: A New Protocol for Mild Cleavage of Sulfinamides and Sulfonamides', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 181: 8, 1825 — 1830

To link to this Article: DOI: 10.1080/10426500500542927

URL: <http://dx.doi.org/10.1080/10426500500542927>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Applications of the $(\text{PhO})_3\text{PCl}_2$ Reagent: A New Protocol for Mild Cleavage of Sulfinamides and Sulfonamides

Alberto Spaggiari

Fabio Prati

Università di Modena e Reggio Emilia, Dipartimento di Chimica,
Modena, Italy

The triphenyl phosphite-chlorine reagent, freshly in situ prepared by the action of chlorine on a solution of triphenyl phosphite in dichloromethane, is successfully used in the cleavage of both sulfinamides and sulfonamides under extremely mild conditions (-15°C to r.t.). Final amines are easily recovered by a simple acid/base extraction, and no further purification is required.

Keywords Amines; chlorine; cleavage; sulfinamides; sulfonamides; triphenyl phosphite

INTRODUCTION

Sulfinamides and sulfonamides represent two classes of highly versatile intermediates in chemical synthesis.^{1–3} Along with their usefulness, however, they are strictly endowed with a remarkably strong chemical stability, which makes them either resistant and, conversely, quite difficult to cleave off,⁴ thus often discouraging their application as protective groups. In the course of our recent investigations into the chemistry of the triphenyl phosphite-chlorine reagent,^{5,6} we had already faced *N*-monosubstituted amides and had reported a highly efficient, and so far the mildest, method for the deacylation of such compounds.⁵ Thus, spurred by the gratifying success obtained in those circumstances and encouraged by the intrinsic structural resemblance, we decided to make a preliminary study aimed at exploring whether

Received March 31, 2005; accepted September 27, 2005.

We thank Riccardo Rubbiani for preliminary experiments and Prof. Franco Ghelfi (Dipartimento di Chimica, Università di Modena e Reggio Emilia, Modena, Italy) for kindly providing the chlorine delivery plant. The authors are also grateful to the Fondazione Cassa di Risparmio di Modena (Modena, Italy) for financial support.


Address correspondence to Alberto Spaggiari, Università di Modena e Reggio Emilia, Dipartimento di Chimica, via Campi, 183, Modena, 41100, Italy. E-mail: spaggiari.alberto@unimore.it

the same reagent could be conveniently employed also in the cleavage of more resistant sulfinamides.

Available methods described in the literature for carrying out the removal of the sulfur-containing fragment in sulfinamides often deal with severe reaction conditions, like acidic^{7,8} or basic⁹ hydrolysis. Acid-catalyzed alcoholysis to sulfinates¹⁰ and, more recently, the use of a perridinane reagent reported by Davis et al.¹¹ as well as the reaction with thiols associated to Lewis' acid settled by Li et al.¹ represent the only attempts to provide milder methods for cleavage. Also, sulfonamides usually require severe conditions to be cleaved, such as, for instance, the action of organolithium reagents,¹² alkali metals,¹³ arene radical anions,¹⁴ and hydrobromic acid.¹⁵ Thus, the quest for a general cleavage protocol under mild conditions would be a desirable issue in organic sulfur chemistry, and this communication therefore aims to propose a novel route that can possibly be applied to all kinds of amides by simply exploiting the high versatility of the same phosphorus-based halogenating reagent.

RESULTS AND DISCUSSION

In this context, we describe here our preliminary results when sulfinamides **1–3** were treated with $(\text{PhO})_3\text{PCl}_2$ in dry dichloromethane, according to the reaction pathway, as depicted in Scheme 1.



SCHEME 1

The reagent was prepared in situ, by bubbling chlorine in a solution of triphenyl phosphite in anhydrous dichloromethane at -15°C . A substrate was added, quickly followed by triethylamine, and the reaction mixture was left to stir at this temperature over a three-hour period.¹⁶ A large excess of ethylene glycol was then added, and the mixture was stirred for an additional twelve-hour period. Dichloromethane was then removed under reduced pressure and replaced by water. Vigorous


Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011




Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011

Downloaded At: 18:23 27 January 2011

sulfinamides also proceed to amines through an iminosulfinyl- or iminosulfonyl chloride analogue (Scheme 3).



SCHEME 3

This initial transformation is achieved by a means of the halogenating action of the oxophilic phosphorus reagent $(\text{PhO})_3\text{PCl}_2$ which, in turn, is converted to triphenyl phosphate, according to its already explored reactivity.⁵ A final alcoholysis of the iminochloride with ethylene glycol and water finally reveals the expected amine, along with either *p*-toluenesulfonyl or *p*-toluenesulfinyl acid, which remains, as sodium salt, in the aqueous layer at the end of the extraction.¹⁷ To the best of our knowledge, although this method does not represent the most efficient protocol for the cleavage of sulfur-based amides, and despite its limitation to the *N*-monosubstituted amides, it involves, however, the mildest reaction conditions, avoiding both harsh reagents and high temperatures, thus becoming a potentially intriguing choice for performing these deprotections when sensitive functionalities are present everywhere in the substrates. This circumstance seriously blocks the route of the most common (and severe) procedures of cleavage. We are at present undertaking further investigations into this reaction, as well as into mechanistic studies, and will report fully in due course.

EXPERIMENTAL

General

All glassware used in the reaction was oven dried and argon flushed. Dichloromethane was dried according to the standard procedure. Triphenyl phosphite was purchased from Aldrich and used without further purification. Chlorine used in the synthesis of the $(\text{PhO})_3\text{PCl}_2$ was checked, and it had a 99.8% title. Triethylamine was purchased from Acros Organics and distilled and stored over KOH before using. Sulfonylamides and sulfinamides **1–7** were prepared according to standard procedures.

General Procedure for the Cleavage of Sulfinamides and Sulfonamides

In a two-necked 50 mL round-bottom flask, anhydrous dichloromethane (15 mL) was cooled to -15°C under an argon atmosphere, and triphenyl phosphite (1100 μL , 4.2 mmol) was added. Chlorine was then bubbled in until the solution became intensely yellow. The color was discharged by the addition of a few drops of triphenyl phosphite, and sulfonamide **4** (1000 mg, 3.8 mmol) was added, quickly followed by triethylamine (610 μL , 4.4 mmol). The reaction was stirred at this temperature over a two-hour period, and then the cold bath was removed, and the mixture was slowly warmed to r.t. After two hours, ethylene glycol (4000 μL , 49 mmol) was added, and the stirring continued overnight. Dichloromethane was removed under reduced pressure and replaced by water (10 mL). After two hours, the mixture was acidified to pH of about 4 by 10% HCl, and the aqueous phase was then basified to pH 11 by 20% NaOH. A final extraction with dichloromethane yielded the expected *p*-toluidine in 61%.

REFERENCES

- [1] B. F. Li, K. Yuan, L. X. Dai, and X. L. Hou, *Synlett.*, **3**, 535 (2005).
- [2] I. V. Koval', *Russ. J. Org. Chem.*, **35**, 475 (1999).
- [3] K. A. Beaver, A. C. Siegmund, and K. L. Spear, *Tetrahedron Lett.*, **37**, 1145 (1996).
- [4] T. W. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, 3rd ed., pp. 603–615, (John Wiley and Sons, New York, 1999).
- [5] A. Spaggiari, L. C. Blaszcak, and F. Prati, *Org. Lett.*, **6**, 3885 (2004). For a recent synthetic application of this protocol, see also A. Spaggiari, L. C. Blaszcak, and F. Prati, *Ars Pharm.*, **46**, 167 (2005).
- [6] A. Spaggiari, P. Davoli, L. C. Blaszcak, and F. Prati, *Synlett.*, **4**, 661 (2005).
- [7] D. H. Hua, S. W. Miao, J. S. Chen, and S. Iguchi, *J. Org. Chem.*, **56**, 4 (1991).
- [8] F. A. Davis, S. Lee, H. M. Zhang, and D. L. Fanelli, *J. Org. Chem.*, **65**, 8704 (2000).
- [9] M. Harmata, M. Kahraman, D. E. Jones, N. Pavri, and S. E. Weatherwax, *Tetrahedron*, **54**, 9995 (1998).
- [10] M. Mikolajczyk, J. Drabowicz, and B. Bujnicki, *J. Chem. Soc. Chem. Commun.*, **14**, 568 (1976).
- [11] F. A. Davis, T. Ramachandar, and H. Liu, *Org. Lett.*, **6**, 3393 (2004).
- [12] I. Fleming, J. Frackenpohl, and I. Hiriyakkanavar, *J. Chem. Soc. Perkin Trans. 1*, **7**, 1229 (1998).
- [13] T. Oshawa, T. Takagaki, F. Ikehara, Y. Takahashi, and T. Oishi, *Chem. Pharm. Bull.*, **30**, 3178 (1982).
- [14] W. D. Closson, S. Ji, and S. Schulenberg, *J. Am. Chem. Soc.*, **92**, 650 (1970).
- [15] H. R. Snyder and H. C. Geller, *J. Am. Chem. Soc.*, **74**, 4864 (1952).
- [16] For more detailed information on the nature of this reagent and related experimental insights, see Ref. 5 and also L. D. Hatfield, L. C. Blaszcak, and J. W. Fisher, US Patent 4,230,644, (1980); *Chem. Abstr.*, **94**, 156523 (1981); L. D. Hatfield,

- L. C. Blaszczak, and J. W. Fisher, US Patent 4,226,986, (1980): *Chem. Abstr.*, **94**, 103348 (1981).
- [17] A GC-MS analysis of the crude reaction mixture in the course of desulfonylation of amide **4** revealed the presence of *p*-toluenesulfonyl chloride.